

Étale homological stability and arithmetic statistics

Benson Farb and Jesse Wolfson ^{*}

August 4, 2016

Abstract

We relate asymptotic point counts and arithmetic statistics over finite fields to homological stability and representation stability over \mathbb{C} in the example of configuration spaces of n points on smooth varieties. In order to accomplish this we must establish subexponential bounds on the growth of the unstable cohomology of such spaces. We apply this and étale homological stability to compute the large n limits of various arithmetic statistics of configuration spaces of varieties over \mathbb{F}_q .

1 Introduction

Let X be a scheme defined over \mathbb{Z} . The Weil conjectures provide a fundamental link between the topology of $X(\mathbb{C})$ and the arithmetic of $X(\mathbb{F}_q)$. As first indicated by work of Ellenberg-Venkatesh-Westerland [EVW], followed by Vakil-Wood [VW], Church-Ellenberg-Farb [CEF2] and others, this correspondence should convert homological stability phenomena in topology to asymptotic point counts on the arithmetic side. We summarize this in the following table, with the rows going from least to most general.

Topology	Arithmetic
$H^*(X(\mathbb{C}))$	$ X(\mathbb{F}_q) $
homological stability of X_n	asymptotics of $ X_n(\mathbb{F}_q) $ as $n \rightarrow \infty$
representation stability	asymptotics of arithmetic statistics on $X_n(\mathbb{F}_q)$

One of the main goals of the present paper is to realize the bottom rows of this table for varieties $P\text{Conf}_n(X)$ (resp. $U\text{Conf}_n(X)$) of configurations of ordered (resp. unordered) n -tuples of points on a smooth variety X . We first state our main theorem, after which we will define all of the terms appearing in its statement.

Theorem A (Arithmetic statistics of configuration spaces): *Let X be a scheme, smooth over $\mathbb{Z}[1/N]$ for some N , with geometrically connected fibers of finite type. Let p be a prime not dividing N , and let q be a power of p . Let Frob_q denote the q^{th} power Frobenius. Let P be any character polynomial, and denote by $H_{et}^i(P\text{Conf}(X))_P$ the stable P -isotypic*

^{*}B.F. is supported in part by NSF Grant Nos. DMS-1105643 and DMS-1406209. J.W. is supported in part by NSF Grant No. DMS-1400349.

part of the étale cohomology of the co-FI-scheme $\mathrm{PConf}_\bullet(X)_{\overline{\mathbb{F}}_q}$ (see §2 below for a precise definition). Denote by $H_{et}^i(\mathrm{UConf}(X))$ the stable étale cohomology of $\mathrm{UConf}_n(X)_{\overline{\mathbb{F}}_q}$. Then

$$\lim_{n \rightarrow \infty} q^{-n \dim X} \sum_{y \in \mathrm{UConf}_n(X)(\mathbb{F}_q)} P(y) = \sum_{i=0}^{\infty} (-1)^i \mathrm{Tr}(\mathrm{Frob}_q \odot H_{et}^i(\mathrm{PConf}(X))^*),$$

in particular, both sides of the above converge. Specializing to $P = 1$, we obtain

$$\lim_{n \rightarrow \infty} q^{-n \dim X} |\mathrm{UConf}_n(X)(\mathbb{F}_q)| = \sum_{i=0}^{\infty} (-1)^i \mathrm{Tr}(\mathrm{Frob}_q \odot H_{et}^i(\mathrm{UConf}(X))^*).$$

The proof of Theorem A has two main steps:

1. We prove what we call *étale homological (and representation) stability* for $H_{et}^*(X_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$. This allows us to break up $H_{et}^*(X_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$ into two parts: stable and unstable.
2. We obtain sub-exponential bounds on the growth of the unstable part of $H_{et}^*(X_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$. This allows us to prove that this unstable part does not contribute, via the Grothendieck-Lefschetz trace formula, to the limiting density of $|X_n(\mathbb{F}_q)|$. As we explain below, the absence of such bounds is a significant obstruction to understanding the asymptotic point counts of many families of interest.

In the étale context, stability of each $H_{et}^i(X_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$ as a Galois representation, not just as a vector space, is crucial. It is the difference between proving that limits such as $\lim_{n \rightarrow \infty} q^{-\dim X_n} |X_n(\mathbb{F}_q)|$ exist, and actually computing the limiting answer. We now discuss these two steps in more detail.

Homological stability. A sequence $\{X_n\}$ of spaces or groups is said to satisfy *homological stability* over a ring R if $H_i(X_n; R)$ or $H^i(X_n; R)$ is independent of n for $n \geq D(i)$; the number $D(i)$ is called the *stable range*. Typically, but not always, there are maps $\psi_n : X_n \rightarrow X_{n+1}$ or $\phi_n : X_{n+1} \rightarrow X_n$ inducing isomorphisms

$$(\psi_n)_* : H_i(X_n; R) \rightarrow H_i(X_{n+1}; R) \quad \text{or} \quad \phi_n^* : H^i(X_n; R) \rightarrow H^i(X_{n+1}; R).$$

Examples of X_n satisfying homological stability include classifying spaces of symmetric groups S_n (Nakaoka), arithmetic groups like $\mathrm{SL}_n \mathbb{Z}$ (Borel), the moduli spaces \mathcal{A}_g (Borel) and \mathcal{M}_g (Harer), and also configuration spaces $\mathrm{UConf}_n(M)$ of unordered n -tuples of distinct points on a manifold M (Arnol'd, McDuff, Segal, Church). Homological stability has been a powerful tool in topology. It converts an *a priori* infinite computation to a finite one. Further, the stable answer, $H_i(X_n; R)$ for $n \geq D(i)$, can often be computed explicitly.

Many natural sequences X_n come equipped with actions of groups G_n by automorphisms. A basic example is the space $\mathrm{PConf}_n(M)$ of ordered n -tuples of distinct points on a manifold M , on which the symmetric group S_n acts by permuting the ordering. Such spaces almost never satisfy homological stability, but they instead often satisfy *representation stability*: the decomposition of $H^i(X_n; \mathbb{Q})$ into a sum of irreducible S_n -representations stabilizes in a precise sense (see [CF, CEF1], §2.1 below, and [Fa] for a survey). When $R = \mathbb{Q}$, plugging the

trivial representation into this theory gives classical homological stability for the sequence X_n/S_n . So for example representation stability for $\mathrm{PConf}_n(M)$ gives classical homological stability for the space $\mathrm{UConf}_n(M) = \mathrm{PConf}_n(M)/S_n$ of unordered n -tuples of distinct points on M ; see [Ch]. The theory of representation stability, initiated by Church, Ellenberg and Farb, is currently undergoing a rapid development.

Étale homological stability. Consider a scheme Y , smooth over $\mathbb{Z}[1/N]$ for some N . We can extend scalars to \mathbb{C} and consider the complex points $Y(\mathbb{C})$, and we can also reduce modulo p for any prime $p \nmid N$. This gives a variety defined over \mathbb{F}_p , and for any positive power $q = p^d$ we can consider both the \mathbb{F}_q -points as well as the $\overline{\mathbb{F}}_q$ -points of Y , where $\overline{\mathbb{F}}_q$ is the algebraic closure of \mathbb{F}_q .

One of the most fundamental arithmetic invariants attached to Y is its *étale cohomology* $H_{et}^*(Y_{/\overline{K}}; \mathbb{Q}_\ell)$, where K is a number field or finite field of characteristic prime to N , and where $\ell \neq p$ is prime. The Galois action on $Y_{/\overline{K}}$ induces a Galois action on each \mathbb{Q}_ℓ -vector space $H_{et}^i(Y_{/\overline{K}}; \mathbb{Q}_\ell)$, and this action is a crucial part of the data.

Now let X_n be a sequence of schemes that are smooth over $\mathbb{Z}[1/N]$ for some N , for example $X_n = \mathrm{PConf}_n(Y)$ or $X_n = \mathrm{UConf}_n(Y)$ for Y smooth over $\mathbb{Z}[1/N]$. Given the usefulness of homological stability in topology, one wants to prove such stability for $H_{et}^i(X_n_{/\overline{K}}; \mathbb{Q}_\ell)$ for K a number field or a finite field with characteristic $p \nmid N$. There are a number of different notions of what “stability” means in this context (see §2 below). We adopt the strongest of these possibilities.

Definition 1.1 (Étale homological stability). *We say that a sequence X_n of schemes satisfies étale homological stability over a field K if for each $i \geq 0$ there exists $D = D(i)$ so that the isomorphism type of $H_{et}^i(X_n_{/\overline{K}}; \mathbb{Q}_\ell)$ as a $\mathrm{Gal}(\overline{K}/K)$ -representation does not depend on n for $n \geq D$. The function $D(i)$ is called the stable range.*

When each X_n in addition admits an S_n -action, such as $X_n = \mathrm{PConf}_n(Y)$ or Y^n , we have a corresponding notion of *étale representation stability* over K . This definition is a bit more involved; see §2.2. It implies étale homological stability for the sequence of varieties X_n/S_n . To state our results in this direction, we need two different descriptions of representations of the symmetric groups S_n .

Let X_i be the class function on all symmetric groups $S_n, n \geq 1$ given by setting $X_i(\sigma)$ to be the number of i -cycles in the cycle decomposition of σ . A *character polynomial* is any polynomial $P \in \mathbb{Q}[X_1, X_2, \dots]$; it is a class function on each $S_n, n \geq 1$. The *degree* of a character polynomial is defined by setting $\deg X_i := i$. See §2.1 below for more details. As shown by Church-Ellenberg-Farb [CEF1], character polynomials give a compact and uniform way of describing the character of certain infinite sequences of S_n -representations for $n = 1, 2, \dots$

A *partition* of n is a sequence $\lambda = (\lambda_1 \geq \dots \geq \lambda_r \geq 0)$ with $\sum_i \lambda_i = n$. The irreducible representations $V(\lambda)$ of S_n are classified by partitions $\lambda \vdash n$. A partition $\lambda \vdash k$ gives a sequence $V(\lambda)_n$ of irreducible S_n -representations for $n \geq k + \lambda_1$ by defining $V(\lambda)_n$ to be the irreducible representation of S_n corresponding to the partition $(n - k, \lambda_1, \dots, \lambda_r)$. Every irreducible representation of S_n is of the form $V(\lambda)_n$ for a unique partition λ ; for example the trivial and standard representations of S_n are $V(0)_n$ and $V(1)_n$, respectively.

Theorem B (Étale representation stability): *Let Y be a scheme, smooth over $\mathbb{Z}[1/N]$ for some N , with geometrically connected fibers of finite type. Let K be either a number field or a finite field of characteristic $p \nmid N$. For each $i \geq 0$, the sequence $H_{et}^i(\mathrm{PConf}_n(Y)_{/\bar{K}}; \mathbb{Q}_\ell)$ of $\mathrm{Gal}(\bar{K}/K)$ -modules satisfies étale representation stability¹ over K with stable range $D(i) = 2i$ for $\dim Y \geq 2$ and $D(i) = 4i$ for $\dim Y = 1$. In particular:*

1. Inductive description: *For all $n \geq 0$, there is an isomorphism of $\mathrm{Gal}(\bar{K}/K)$ -representations:*

$$H_{et}^i(\mathrm{PConf}_n(Y)_{/\bar{K}}; \mathbb{Q}_\ell) \cong \mathrm{colim}_S H_{et}^i(\mathrm{PConf}_{|S|}(Y)_{/\bar{K}}; \mathbb{Q}_\ell) \quad (1.1)$$

where the colimit is taken over the poset of all subsets $S \subset \{1, \dots, n\}$ such that $|S| \leq D(i)$. This gives, for each $n \geq D(i)$, a recipe for building the $\mathrm{Gal}(\bar{K}/K)$ -representation $H_{et}^i(\mathrm{PConf}_n(Y)_{/\bar{K}}; \mathbb{Q}_\ell)$ from a fixed finite collection of $\mathrm{Gal}(\bar{K}/K)$ -representations.

2. Stability of isotropics: *For each character polynomial P , there exists a unique virtual $\mathrm{Gal}(\bar{K}/K)$ -representation $H_{et}^i(\mathrm{PConf}(Y))_P$ over \mathbb{Q}_ℓ , linear in P , so that when $P = \chi_{V(\lambda)}$ for some $\lambda \vdash k$, there exists D such that for all $n \geq D$:*

$$H_{et}^i(\mathrm{PConf}(Y))_{\chi_{V(\lambda)}} = H_{et}^i(\mathrm{PConf}_n(Y)_{/\bar{K}}; \mathbb{Q}_\ell) \otimes_{\mathbb{Q}_\ell[S_n]} V(\lambda)_n$$

and the right hand side is independent of n as a $\mathrm{Gal}(\bar{K}/K)$ -representation.

3. Polynomial characters: *There exists a character polynomial $Q(X_1, \dots, X_r)$ so that for all $n \geq D(i)$:*

$$\chi_{H_{et}^i(\mathrm{PConf}_n(Y)_{/\bar{K}}; \mathbb{Q}_\ell)}(\sigma) = Q(X_1(\sigma), \dots, X_r(\sigma)) \quad \text{for all } \sigma \in S_n.$$

where $\deg(Q) \leq i$ if $\dim Y > 1$ and $\deg(Q) \leq 2i$ if $\dim Y = 1$.

Remarks:

1. Theorems A and B in the special case $Y = \mathbb{A}^1$ was proved by Church-Ellenberg-Farb [CEF2]. This special case is much simpler since the eigenvalues of Frob_q on $H_{et}^i(\mathrm{PConf}_n(\mathbb{A}^1)_{/\mathbb{F}_q}; \mathbb{Q}_\ell)$ are known explicitly; they equal q^i .
2. Our proof of Theorem B uses the theory of FI-modules, developed in [CEF1]. What we actually prove is that for each $i \geq 0$, the FI- $\mathrm{Gal}(\bar{K}/K)$ -module (see §2.2 below for the precise definition) $H_{et}^i(\mathrm{PConf}_\bullet(Y); \mathbb{Q}_\ell)$ is finitely generated. Items (1)–(3) of Theorem B then follow from the general theory of FI-modules, in particular theorems from [CEF1] and [CEFN]. As one consequence, the proof shows that Item (1) of Theorem B holds with \mathbb{Q}_ℓ replaced by \mathbb{Z}_ℓ or $\mathbb{Z}/\ell^n\mathbb{Z}$.
3. Nir Gadish [Ga] has recently isolated a concept of finitely generated I -poset, for a wide class of categories I , and has used this to prove étale representation stability for a rich class of sequences of complements of linear subspace arrangements.

¹See §2.2 for the precise definition of étale representation stability.

Plugging in $P = 1$ into Item (2) of Theorem B gives the following.

Corollary B' (Étale homological stability): *With terminology as in Theorem B, the sequence $H_{et}^i(\mathrm{UConf}_n(Y)_{/\overline{K}}; \mathbb{Q}_\ell)$ satisfies étale homological stability over K : these $\mathrm{Gal}(\overline{K}/K)$ -representations do not depend on n for $n \geq D(i)$.*

Remarks:

1. Quoc Ho [Ho] has recently given an independent proof of Corollary B' for Y smooth over any ground field. His method is based on factorization homology, and is quite different from the methods of this paper.
2. Dan Petersen [Pe] has recently extended Theorem B to a wider class of configuration-like spaces, also dropping the smoothness assumption.

Stability of arithmetic statistics. The application of homological stability to arithmetic statistics was pioneered by Ellenberg-Venkatesh-Westerland in [EVW]. The fundamental link is provided by the *Grothendieck-Lefschetz Trace Formula*²:

$$|Z(\mathbb{F}_q)| = q^{\dim(Z)} \sum_{i \geq 0} (-1)^i \mathrm{Tr}(\mathrm{Frob}_q : H_{et}^i(Z_{/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)^* \longrightarrow H_{et}^i(Z_{/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)^*) \quad (1.2)$$

and its twisted version (see (4.1) below). Given Deligne's theorem [De1, Theorem 1.6] that any eigenvalue λ of Frob_q on $H_{et}^i(Z_{/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)^*$ satisfies $|\lambda| \leq q^{-i/2}$, one can bound the number $|Z(\mathbb{F}_q)|$ of \mathbb{F}_q -points via

$$|Z(\mathbb{F}_q)| \leq q^{\dim Z} \sum_{i=0}^{2 \dim Z} b_i q^{-i/2}$$

where $b_i := \dim H_{et}^i(Z_{/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)^*$. Applying this reasoning to a sequence Z_n of smooth varieties gives

$$q^{-\dim Z_n} |Z_n(\mathbb{F}_q)| \leq \sum_{i=0}^{2 \dim Z_n} b_i(n) q^{-i/2} \quad (1.3)$$

where we have emphasized via notation that b_i is a function of n . It seems that étale homological stability, namely the fact that $b_i(n)$ is constant for $n \geq D(i)$, should imply that the limit as $n \rightarrow \infty$ of the left-hand side of (1.3) exists. However, it could be that $\dim(Z_n)$ goes to ∞ with n and that $b_i(n)$ grows more quickly than $q^{i/2}$, even for any q ; this would imply the divergence of the right-hand side of (1.3). This super-exponential growth is known to occur in natural examples, for example for Z_n the moduli space of genus n smooth algebraic curves, and also for Z_n the moduli space of n -dimensional principally polarized abelian varieties. In the latter example, recent work of Lipnowski-Tsimerman [LT] shows that this growth actually does change the point count $|Z_n(\mathbb{F}_q)|$, as they show this number grows more quickly than the expected $q^{\dim Z_n}$.

Thus, in order to apply étale homological stability to obtain the existence of asymptotic point counts in a given example, it is necessary to prove sub-exponential (in i) bounds

²Here we have assumed that Z is smooth and applied Poincaré Duality to the usual Grothendieck-Lefschetz Formula.

on $b_i(n)$, independent of n . In other words, control of the *unstable* étale cohomology $H_{et}^i(Z_{n/\bar{\mathbb{F}}_q}; \mathbb{Q}_\ell)^*$ is needed.

Proving such bounds is a major obstruction for arithmetic applications; see §3 for a discussion. This problem is a very special case (namely the case $P \equiv 1$) of more general arithmetic statistics, where one needs a twisted version of the Grothendieck-Lefschetz formula, and where the control on the “representation unstable cohomology” is even more difficult to prove; see §3 below. A significant part of this paper, §3, is devoted to overcoming this problem for the examples X^n and $\mathrm{PConf}_n(X)$.

Theorem C (Bounding the representation unstable cohomology): *Let X be either a smooth, orientable manifold with $\dim(H^*(X; \mathbb{Q})) < \infty$ (e.g. X compact), or a scheme, smooth over $\mathbb{Z}[1/N]$ for some N , with geometrically connected fibers of finite type. In the first case let H^i denote singular cohomology with \mathbb{Q} coefficients; in the second, let H^i denote étale cohomology with \mathbb{Q}_ℓ coefficients. Then for any character polynomial P there exists a function $F_P(i)$, subexponential in i , such that for all $n \geq 1$:*

$$\langle P, (H^i(\mathrm{PConf}_n(X))) \rangle_{S_n} \leq F_P(i).$$

In §4, we apply Theorems B and C to obtain Theorem A.

A different description of the left-hand side of Theorem C, established using analytic methods, will appear in forthcoming work of Weiyan Chen [Che]. En route to proving Theorems C and A, we also prove the analogous statements for $\mathrm{Sym}^n(X)$; see §4. However, we note that Theorem C requires more than just bounding the betti numbers $\mathrm{Sym}^n(X)$.

Acknowledgements. It is a pleasure to thank George Andrews, Kathrin Bringmann, Mark Kisin, and Ken Ono for useful discussions. We thank Nir Gadish and Brian Conrad for very helpful conversations about twisted ℓ -adic sheaves and transfer. We thank Weiyan Chen, Jordan Ellenberg and Burt Totaro for numerous helpful comments on an earlier draft. It is a pleasure to thank Matt Emerton for his careful reading and many detailed comments on an earlier draft of this paper. Finally, we thank an anonymous referee for making several suggestions that helped to simplify the proofs in this paper.

2 Étale Representation stability

In this section, we briefly summarize the theory of representation stability and FI-modules, as it is used in topology, as well as some of its consequences. This theory was developed by Church, Ellenberg and Farb [CF, CEF1], and later with Nagpal [CEFN]; see [Fa] for a survey. We refer the reader to these references for details. We then give a general setup for proving similar stability theorems in étale cohomology.

2.1 Quick summary of representation stability and FI-modules

An *FI-module* V over a Noetherian ring R is a functor from the category FI of finite sets and injections to the category of R -modules. Thus to each natural number n we have

associated an R -module V_n with an S_n action, with a map $V_m \rightarrow V_n$ for each injection $\{1, \dots, m\} \rightarrow \{1, \dots, n\}$. Recall that the opposite category FI^{op} is the same as FI but with arrows reversed. A *co-FI module* over R is a functor from FI^{op} to R -modules. We also have the associated notions of *FI-space*, *FI-scheme*, etc., and the associated co-FI versions.

An FI-module V is *finitely generated* if there is a finite set S of elements in $\coprod_i V_i$ so that no proper sub-FI-module of V contains S . One of the reasons that we care about finitely-generated FI-modules is the following theorem.

Theorem 2.1 (Structural properties of finitely-generated FI-modules). *Let V be an FI-module over a commutative Noetherian ring R . If V is finitely-generated then:*

Representation stability ([CEF1]): *When R is a field of characteristic 0, finite generation of V implies representation stability in the sense of [CF] for the sequence $\{V_n\}$ of S_n -representations.*

Inductive description ([CEFN]): *Let V_{\bullet} be a finitely-generated FI-module over a Noetherian ring R . Then there exists some $N \geq 0$ such that for all $n \in \mathbb{N}$, there is a natural isomorphism*

$$V_n \cong \operatorname{colim}_{S \subset [n], |S| \leq N} V_S,$$

i.e. these isomorphisms commute with homomorphisms of FI-modules. By definition, the stable range of V_{\bullet} $N(V)$ is the minimal such N .

Isomorphism of trivial isotopies ([Ch]): *Let V_{\bullet} be a finitely-generated FI-module over a Noetherian ring R with stable range $N(V)$. Then for all $n \geq N(V)$ the map $V_n \rightarrow V_{n+1}$, given by averaging the structure maps, induces an isomorphism*

$$V_n^{S_n} \xrightarrow{\cong} V_{n+1}^{S_{n+1}}.$$

We remark that the isomorphism of trivial isotopies illustrates one of the key advantages of considering (co-)FI-spaces: while stabilization maps for many natural sequences of spaces or schemes do not naively exist, a (co-)FI-space Z_{\bullet} comes equipped with canonical rational correspondences from Z_{n+1}/S_{n+1} to Z_n/S_n .

We will also need the following.

Lemma 2.2. *Let V_{\bullet} and W_{\bullet} be finitely generated FI-modules, and let N be the sum of their stable ranges. Then for all $n \geq N$, the maps $V_n \rightarrow V_{n+1}$ and $W_n \rightarrow W_{n+1}$ (associated to $\{1, \dots, n\} \subset \{1, \dots, n+1\}$) induce isomorphisms*

$$V_n \otimes_{\mathbb{Q}[S_n]} W_n \xrightarrow{\cong} V_{n+1} \otimes_{\mathbb{Q}[S_{n+1}]} W_{n+1}$$

that are natural in both variables with respect to homomorphisms of FI-modules.

Proof. By [CEF1, Proposition 2.3.6], the tensor product $V_{\bullet} \otimes_{\mathbb{Q}} W_{\bullet}$ is finitely generated since V_{\bullet} and W_{\bullet} are. Applying the co-invariants functor, we obtain the functor

$$n \mapsto V_n \otimes_{\mathbb{Q}[S_n]} W_n.$$

Because stable ranges add under tensor product ([CEF1, Proposition 2.3.6]), and because the stability degree (cf. [CEF1, Definition 3.1.3]) is less than or equal to the stable range ([CEF1, Proposition 3.3.3]), the map

$$V_n \otimes_{\mathbb{Q}[S_n]} W_n \rightarrow V_{n+1} \otimes_{\mathbb{Q}[S_{n+1}]} W_{n+1}$$

is an isomorphism for $n \geq N$. □

Character polynomials. Character polynomials and their degree were defined in the introduction. Let $\langle P, Q \rangle$ denote the inner product of S_n -characters. The expectations of character polynomials

$$\mathbb{E}_{\sigma \in S_n} P_n(\sigma) := \frac{1}{n!} \sum_{\sigma \in S_n} P_n(\sigma) = \langle P_n, 1 \rangle$$

compute the averages of natural combinatorial statistics with respect to the uniform distribution on S_n . As shown in Proposition 2.2 of [CEF2], the inner product $\langle P_n, Q_n \rangle$ of character polynomials $P, Q \in \mathbb{Q}[X_1, X_2, \dots]$ is independent of n once $n \geq \deg P + \deg Q$.

One remarkable property of finitely-generated FI-modules V is that the characters of the S_n -representations V_n are, for large enough n , given by a single polynomial.

Theorem 2.3 (Polynomiality of characters [CEFN]). *Let V be an FI-module over a field of characteristic 0. If V is finitely-generated then the characters χ_{V_n} of the S_n -representations V_n are eventually polynomial: there exists $N \geq 0$ and a polynomial $P(X_1, \dots, X_r)$, for some $r > 0$, so that*

$$\chi_{V_n} = P(X_1, \dots, X_r) \quad \text{for all } n \geq N. \quad (2.1)$$

In particular, if Q is any character polynomial then $\langle \chi_{V_n}, Q \rangle$ is independent of $n \geq \deg P + \deg Q$.

We note that evaluating (2.1) on the identity permutation gives a polynomial $P(T) \in \mathbb{Q}[T]$ so that

$$\dim_k V_n = P(n)$$

for all $n \geq N$.

Étale Representation Stability Given a co-FI-scheme Z_\bullet defined over \mathbb{F}_q , its étale cohomology $H_{et}^i(Z_{\bullet/\bar{\mathbb{F}}_q}; \mathbb{Q}_\ell)$ has additional structure beyond that of an FI-module over \mathbb{Q}_ℓ . The geometric Frobenius Frob_q gives a natural endomorphism of $Z_{\bullet/\bar{\mathbb{F}}_q}$, and this gives rise to an action of $\text{Gal}(\bar{\mathbb{F}}_q/\mathbb{F}_q)$ on the FI-module $H_{et}^i(Z_{\bullet/\bar{\mathbb{F}}_q}; \mathbb{Q}_\ell)$. As noted in the introduction, the eigenvalues of Frob_q and the action of $\text{Gal}(\bar{\mathbb{F}}_q/\mathbb{F}_q)$ are crucial parts³ of the data here. Weaker than knowing an eigenvalue λ of Frob_q on $H_{et}^j(Z_{n/\bar{\mathbb{F}}_q}; \mathbb{Q}_\ell)$ is knowing its weight. Deligne proved that λ is an algebraic number with $|\lambda| = q^{r/2}$ for some $j \leq r \leq 2j$, with $r = j$ if Z_n is smooth and proper. The number r is the *weight* of the eigenvalue λ . Similarly, for Z_\bullet defined over a number field K , the action of $\text{Gal}(\bar{K}/K)$ on $Z_{\bullet/\bar{K}}$ induces an action on $H_{et}^i(Z_{\bullet/\bar{K}}; \mathbb{Q}_\ell)$, and this action is a fundamental part of the data.

In increasing order of strength, we could ask that for each i there exists D so that for all $n \geq D$:

1. The isomorphism type of $H_{et}^i(Z_{n/\bar{K}}; \mathbb{Q}_\ell)$ as a \mathbb{Q}_ℓ -vector space does not depend on n ;
2. in addition, the list of weights of Frob_q on $H_{et}^i(Z_{n/\bar{K}}; \mathbb{Q}_\ell)$ does not depend on n ;
3. in addition, the list of eigenvalues of Frob_q on $H_{et}^i(Z_{n/\bar{K}}; \mathbb{Q}_\ell)$ does not depend on n .

³As observed e.g. by Milne [Mi2], the Tate conjecture implies that the eigenvalues of Frob_q determine the $\text{Gal}(\bar{\mathbb{F}}_q/\mathbb{F}_q)$ -action. But, this is not known at present.

4. The isomorphism type of $H_{et}^i(Z_{n/\bar{K}}; \mathbb{Q}_\ell)$ as a $\text{Gal}(\bar{K}/K)$ -representation does not depend on n .

We have adopted the strongest of these as our definition of étale homological stability.

2.2 Étale representation stability

Let Z be a co-FI scheme smooth over $\mathbb{Z}[1/N]$ for some fixed N , with geometrically connected fibers. Let $p \nmid N$ be prime, and let $\ell \neq p$ be a prime. For each $i \geq 0$, the étale cohomology $H_{et}^i(Z_{/\bar{\mathbb{F}}_p}; \mathbb{Z}_\ell)$ is an FI-module. In addition, for each $q = p^d$, the Frobenius Frob_q acts on each $H_{et}^i(Z_{n/\bar{\mathbb{F}}_p}; \mathbb{Z}_\ell)$, endowing it with the structure of a $\text{Gal}(\bar{\mathbb{F}}_p/\mathbb{F}_q)$ -module. The S_n -action on $H_{et}^i(Z_{n/\bar{\mathbb{F}}_p}; \mathbb{Z}_\ell)$ coming from its structure as an FI-module commutes with the action of $\text{Gal}(\bar{\mathbb{F}}_p/\mathbb{F}_q)$, as do all automorphisms of Z_n . Similarly, for any number field K , the action of $\text{Gal}(\bar{K}/K)$ on $H_{et}^i(Z_{/\bar{K}}; \mathbb{Q}_\ell)$ commutes with the FI-structure.

This discussion shows that, for K a number field or a finite field of characteristic prime to N , $H_{et}^i(Z_{/\bar{K}}; \mathbb{Q}_\ell)$ is an $\text{Gal}(\bar{K}/K)$ -FI-module; that is, an FI-module equipped with an action of $\text{Gal}(\bar{K}/K)$ by FI-automorphisms. We have the corresponding notions of finitely generated $\text{Gal}(\bar{K}/K)$ -FI-module: there is a finite set $S \subset \coprod_n H_{et}^i(Z_{n/\bar{K}}; \mathbb{Q}_\ell)$ so that no proper sub-FI-module of $H_{et}^i(Z_{/\bar{K}}; \mathbb{Q}_\ell)$ contains S .

Definition 2.4 (Étale representation stability). *We say that a sequence Z_n of $\text{Gal}(\bar{K}/K)$ -modules satisfies étale representation stability if $\{1, \dots, n\} \mapsto Z_n$ is a finitely generated $\text{Gal}(\bar{K}/K)$ -FI-module.*

Theorem C in [CEFN] gives an inductive description of finitely-generated FI-modules V over any Noetherian ring R . Namely, there exists $D \geq 0$ such that for all $n \in \mathbb{N}$, there is a natural isomorphism

$$V_n \cong \text{colim}_S V_S \tag{2.2}$$

where the colimit is taken over the poset of all subsets $S \subset \{1, \dots, n\}$ such that $|S| \leq D$. If V is a finitely-generated $\text{Gal}(\bar{K}/K)$ -FI-module then (2.2) gives an isomorphism of $\text{Gal}(\bar{K}/K)$ -modules. Thus (2.2) gives, for each $n \geq D$, a recipe for building the $\text{Gal}(\bar{K}/K)$ -representation V_n from a fixed finite collection of $\text{Gal}(\bar{K}/K)$ -representations.

Étale representation stability for products and configuration spaces. Attached to any scheme X there is an associated *configuration space* $\text{PConf}_n(X)$ of ordered n -tuples in X , defined by:

$$\text{PConf}_n(X) := \{(x_1, \dots, x_n) \in X^n : x_i \neq x_j \ \forall i \neq j\} = X^n \setminus \Delta$$

where Δ is the fat diagonal and where we write $x \in X$ to denote an arbitrary R -point of X . The group S_n acts freely on PConf_n by permuting the coordinates. The quotient $\text{UConf}_n(X) := \text{PConf}_n(X)/S_n$ is the *configuration space of unordered n -tuples* of points in X .

For any scheme X , denote by X^\bullet the co-FI scheme that sends $\{1, \dots, n\}$ to the cartesian product X^n , and associates to any injection between finite sets the natural projection

maps between cartesian powers. The open subsets $\mathrm{PConf}_n(X) \subset X^n$ are preserved under the co-FI structure maps, and so we obtain a co-FI scheme $\mathrm{PConf}_\bullet(X)$. In particular, $H^i(\mathrm{PConf}_\bullet(X); \mathbb{Q}_\ell)$ is an FI-module for any $i \geq 0$.

With this setup, we can now prove Theorem B from the introduction. The proof also gives the following result.

Theorem 2.5. *Theorem B with $\mathrm{PConf}_n(Y)$ (resp. $\mathrm{UConf}_n(Y)$) replaced by Y^n (resp. $\mathrm{Sym}^n(Y)$) holds.*

Proof of Theorem B and Theorem 2.5. This theorem follows from the proofs of [CEF1, Theorem 6.1.2] and [CEF1, Theorem 6.2.1]. The only difference is that we now work with étale rather than Betti cohomology.

To wit, the co-FI-schemes $\mathrm{PConf}_\bullet(Y)$ and Y^\bullet give rise to $FI\text{-Gal}(\bar{K}/K)$ -modules by taking ℓ -adic cohomology. The proof of [CEF1, Theorem 6.1.2] carries over verbatim to the étale setting to show that $H_{et}^i(Y^\bullet; \mathbb{Q}_\ell)$ is a finitely generated $FI\text{-Gal}(\bar{K}/K)$ -module for all $i \geq 0$; see Theorem 4.1.7 and Remark 6.1.3 of [CEF1] for the stable range and degree of the character polynomial. Note here that this improved stable range comes from the fact that X^\bullet is a so-called FI#-module.

For $\mathrm{PConf}_\bullet(Y)$, the proof of [CEF1, Theorem 6.2.1] applies in étale cohomology just as for singular cohomology. Indeed, as Totaro discusses in [To, p. 1064], for any scheme Y smooth over $\mathbb{Z}[1/N]$, the Leray spectral sequence in ℓ -adic cohomology for the inclusion

$$\mathrm{PConf}_n(Y) \hookrightarrow Y^n$$

has E_2 -page isomorphic to

$$E_2^{p,q} = \bigoplus_{\{J \vdash n \mid |J|=n-q/(2\dim(Y)-1)\}} H_{et}^p(Y_J; \mathbb{Q}_\ell^{\mathrm{c}_J}(-\frac{q\dim(Y)}{2\dim(Y)-1}))$$

where $Y_J \subset Y^n$ denotes the diagonal where points coincide according to the partition J , $\dim(Y)$ denotes the dimension of the scheme, and if J consists of pieces of size $j_1, \dots, j_{n-q/(2\dim(Y)-1)}$, then

$$c_J := (j_1 - 1)! \cdots (j_{n-q/(2\dim(Y)-1)} - 1)!.$$

See also [We] for a detailed proof of this description of the Leray spectral sequence in the étale setting. The proof of [CEF1, Theorem 6.2.1] now carries over verbatim to show that for all i , the $FI\text{-Gal}(\bar{K}/K)$ -module $H_{et}^i(\mathrm{PConf}_\bullet(Y); \mathbb{Q}_\ell)$ is finitely generated for all $i \geq 0$. For the claimed stable ranges and degree of character polynomial, see [CEF1, Theorem 6.3.1]. \square

3 Convergent Cohomology

In this section we provide the necessary bounds for the “representation unstable cohomology” of X^n and of $\mathrm{PConf}_n(X)$ that will be necessary for the arithmetic applications in Section 4.

3.1 Definition of convergent cohomology

A function $F : \mathbb{N} \longrightarrow \mathbb{N}$ has exponential growth rate λ if

$$\lim_{n \xrightarrow{\longrightarrow} \infty} \frac{\log f(n)}{n} = \lambda. \quad (3.1)$$

If (3.1) holds with $\lambda = 0$, we say that F has *sub-exponential growth*.

Let Z be a co-FI-scheme over $\mathbb{Z}[1/N]$. For each $i \geq 0$, let $H^i(Z_n)$ denote either the singular cohomology $H^i(Z_n(\mathbb{C}); \mathbb{Q})$ or the étale cohomology $H_{et}^i(Z_n/\overline{K}; \mathbb{Q}_\ell)$ for K a number field or finite field of characteristic prime to N . In each case $H^i(Z_\bullet)$ is an FI-module (over \mathbb{Q} and \mathbb{Q}_ℓ , respectively). For any class function P on S_n , denote by $\langle P, H^i(Z_n) \rangle$ the inner product of (the character of) $H^i(Z_n)$ with P .

In order to compute arithmetic statistics for a co-FI scheme Z , one needs to control the “representation unstable” cohomology of Z ; see §4. More precisely, one needs to prove one of the following two properties, which were shown to be equivalent in [CEF2, §3]:

1. For each $0 \leq a \leq n$ there is a function $F_a(i)$, subexponential in i and not depending on n , so that:

$$\dim H^i(Z_n)^{S_{n-a}} \leq F_a(i) \text{ for all } n \text{ and } i. \quad (3.2)$$

2. For each character polynomial $P \in \mathbb{Q}[X_1, X_2, \dots]$ there exists a function $F_P(i)$, subexponential in i and not depending on n , such that:

$$|\langle P, H^i(Z_n) \rangle| \leq F_P(i) \text{ for all } n \text{ and } i. \quad (3.3)$$

It is crucial that these bounds hold independently of n . While the second condition is the one that applies to arithmetic statistics (see §4 below), it is quite difficult to check. Thus the equivalence with the first condition is quite useful.

Definition 3.1 (Convergent cohomology). *We say that the co-FI scheme (or space) Z has convergent (singular or étale) cohomology if either of the two equivalent properties 1 or 2 in equations (3.2) or (3.3) holds. If these properties hold with $F_P(i)$ having exponential growth $0 < \lambda < \infty$, we say that Z has weakly convergent cohomology with convergence rate λ .*

These kinds of bounds are typically not easy to prove. In [EVW] this is accomplished⁴ for the cohomology of certain Hurwitz spaces by obtaining an exponential upper bound for the number of i -cells, via an explicit cell decomposition. In [CEF2] such bounds for the example $H^i(PConf_n(\mathbb{C}); \mathbb{Q})$ are obtained by a detailed knowledge of these S_n -representations. The rest of this section is devoted to giving such bounds for two natural classes of co-FI schemes. We then apply this in §4 to arithmetic statistics for \mathbb{F}_q -points on these schemes.

3.2 Polynomial bounds on Betti numbers of symmetric products

Let X be a topological space. The n -fold cartesian product X^n is endowed with a natural action of the symmetric group S_n , given by permuting the factors. The quotient $\text{Sym}^n X := X^n/S_n$ is called the the n^{th} symmetric product of X .

Proposition 3.2 (Growth of Betti numbers of symmetric products). *Let X be either a space with $\dim(H^*(X; \mathbb{Q})) < \infty$ (e.g. X compact), or a scheme over $\mathbb{Z}[1/N]$ for some N , with geometrically connected fibers of finite type. In the first case let H^i denote singular cohomology with \mathbb{Q} coefficients; in the second, let H^i denote étale cohomology with \mathbb{Q}_ℓ coefficients. In either case let b_i be the associated i^{th} Betti number. Then $b_i(\text{Sym}^n(X))$ is bounded above by a polynomial in i , independent of n .*

⁴[EVW] only needs to deal with the classical, not representation stable, case; that is, the $a = 0$ case.

Proof. We claim that if $m < n$ then $b_i(\text{Sym}^m(X)) \leq b_i(\text{Sym}^n(X))$; further, $b_i(\text{Sym}^n(X)) = b_n(\text{Sym}^n(X))$ for all $i \geq n$. To see the statement for $m < n$, observe that for any graded vector space V (over a field of characteristic 0), a choice of “unit” $1 \in V_0$, determines an injection

$$\begin{aligned}\text{Sym}^m(V)_i &\longrightarrow \text{Sym}^n(V)_i \\ \vec{v} &\mapsto \vec{v} \otimes 1 \otimes \cdots \otimes 1\end{aligned}$$

In particular, $\dim(\text{Sym}^m(V)_i) \leq \dim(\text{Sym}^n(V)_i)$ for all $m < n$ and i . Considering $V = H^*(X; \mathbb{Q})$, Künneth and transfer imply that

$$\text{Sym}^n(H^*(X; \mathbb{Q}))_i \cong H^i(\text{Sym}^n(X); \mathbb{Q})$$

and the first part of the claim follows. For the second, we note that for a graded vector space $V = V_0 \oplus \cdots \oplus V_m$, with $V_0 = \mathbb{Q}$, we have

$$\text{Sym}^n(V) \cong \bigoplus_{a_0 + \cdots + a_m = n} \bigotimes_{j=1}^m \text{Sym}^{a_j}(V_j)$$

and thus

$$\text{Sym}^n(V)_i \cong \bigoplus_{(a_1, \dots, a_m)} \bigotimes_{j=1}^m \text{Sym}^{a_j}(V_j)$$

where the direct sum is over partitions $a_1 + 2a_2 + \cdots + ma_m = i$ such that $a_1 + \cdots + a_m = n$. In particular, the number of pieces in the partition is at most n , and since for any i , the largest number of pieces in any partition is i , we see that for $n \geq i$, the direct sum is independent of n . We conclude the claim by taking $V = H^*(X; \mathbb{Q})$ and invoking Künneth and transfer as above.

We have just shown that for each fixed $i \geq 0$:

$$b_i(\text{Sym}^n(X)) \leq b_i(\text{Sym}^i(X)) = b_i(\text{Sym}^\infty(X)) \quad \text{for all } n \geq 1$$

To prove the proposition, it is therefore enough to bound $b_i(\text{Sym}^\infty(X))$ by a polynomial in i . Well, note that the function $f(z) := \sum_{i=0}^{\infty} b_i(\text{Sym}^\infty(X))z^i$ is just the Poincaré series for the symmetric algebra on the vector space $H^*(X; \mathbb{Q})$. By an elementary argument, this Poincaré series is, writing $B_i := b_i(X)$, the following rational function :

$$f(z) = \frac{(1+z)^{B_1}(1+z)^{B_3} \cdots}{(1-z^2)^{B_2}(1-z^4)^{B_4} \cdots}$$

Since each pole of $f(z)$ lies on the unit circle, it follows (see, e.g. [FS], Theorem IV.9) that the i^{th} coefficient $b_i(\text{Sym}^\infty(X))$ of $f(z)$ is bounded above by a polynomial in i .

□

Consequence: bounding the representation unstable cohomology of products. The following corollary is also a key ingredient in bounding the representation unstable cohomology of configuration spaces.

Corollary 3.3. *Let X be as in Proposition 3.2, and use the notation of that theorem. For each $0 \leq a \leq n$, we have that $\dim(H^i(X^n)^{S_{n-a}})$ is bounded above by a polynomial in i , independent of n .*

Proof. Since the action S_{n-a} leaves invariant the first $n - a$ factors of X^n and acts as the identity on the last a factors, there is, for each $i \geq 0$, an isomorphism:

$$H^i(X^n)^{S_{n-a}} = \bigoplus_{p+q=i} H^p(X^{n-a})^{S_{n-a}} \otimes H^q(X^a). \quad (3.4)$$

Since this sum has $i + 1$ terms, it suffices to bound the dimension of each summand by a polynomial in i . Since $\dim H^*(X; \mathbb{Q}) < \infty$ and since a is fixed, there is a constant C , not depending on q , so that $\dim(H^q(X^a)) \leq C$. It thus suffices to bound each $H^p(X^{n-a})^{S_{n-a}}$ by a polynomial in i . But this follows from transfer together with Proposition 3.2, noting that $p \leq i$. \square

3.3 Bounding the representation unstable cohomology of configuration spaces

We build on the subexponential upper bounds for products in the last section to prove the corresponding result for configuration spaces.

Theorem 3.4 (PConf $_\bullet(X)$ has convergent cohomology). *Let X be either a smooth, orientable manifold with $\dim(H^*(X; \mathbb{Q})) < \infty$ (e.g. X compact), or a scheme, smooth over $\mathbb{Z}[1/N]$ for some N , with geometrically connected fibers of finite type. Then the co-FI manifold (resp. scheme) PConf $_\bullet(X)$ has convergent singular (resp. étale) cohomology.*

Proof. For the case when X is a manifold we let H^i denote singular cohomology with \mathbb{Q} coefficients; in the case when X is a scheme we let H^i denote étale cohomology with \mathbb{Q}_ℓ coefficients.

Fix $a \geq 0$. Denote by S_{n-a} the subgroup $S_{n-a} \times 1 \subset S_n$. We will prove that there is a function $F_a(i)$, subexponential in i , so that:

$$\dim(H^i(\mathrm{PConf}_n(X); \mathbb{Q}))^{S_{n-a}} \leq F_a(i)$$

for all $i \geq 0$. Let m be the real dimension of X , and denote by $A(n, m)$ the graded commutative algebra

$$A(n, m) := \mathbb{Q}[\{G_{ab}\}_{1 \leq a \neq b \leq n}] / I$$

where $|G_{ab}| = 2m - 1$ and I is the ideal generated by the elements

$$\begin{aligned} G_{ab} - G_{ba} \\ G_{ab}G_{ac} + G_{bc}G_{ba} + G_{ca}G_{cb} \end{aligned}$$

for $a < b < c$ distinct. The group S_n acts on $A(n, m)$ via $\sigma \cdot G_{ab} := G_{\sigma(a)\sigma(b)}$. Totaro [To, Theorem 4] has shown that $H^*(\mathrm{PConf}_n(X); \mathbb{Q})$ is isomorphic, as a graded S_n -representation, to a sub-quotient of

$$H^*(X^n; \mathbb{Q}) \otimes A(n, m),$$

with the natural action on each factor. As Totaro indicates, this result holds for both the singular and étale cohomology.

Note that, for any short exact sequence of S_n -representations

$$0 \longrightarrow V_0 \longrightarrow V_1 \longrightarrow V_2 \longrightarrow 0$$

over a field of characteristic 0, there exists an S_n -equivariant splitting

$$V_1 \cong V_0 \oplus V_2.$$

In particular,

$$\dim V_1^{S_n} = \dim V_0^{S_n} + \dim V_2^{S_n},$$

and, more generally, if V is any sub-quotient of an S_n -representation W , we have

$$\dim V^{S_n} \leq \dim W^{S_n}.$$

Let V and W be any two S_n -representations. The identity $\dim(V^{S_n}) = \langle \chi_V, 1 \rangle$ and the Cauchy-Schwarz inequality give:

$$\begin{aligned} \dim(V \otimes W)^{S_n} &= \langle \chi_{V \otimes W}, 1 \rangle \\ &= \frac{1}{n!} \sum_{\sigma \in S_n} \chi_V(\sigma) \chi_W(\sigma) \\ &\leq \frac{1}{n!} \sqrt{\left(\sum_{\sigma \in S_n} \chi_V(\sigma)^2 \right) \left(\sum_{\sigma \in S_n} \chi_W(\sigma)^2 \right)} \\ &= \sqrt{\langle \chi_{V^{\otimes 2}}, 1 \rangle \langle \chi_{W^{\otimes 2}}, 1 \rangle} \\ &= \sqrt{\dim((V^{\otimes 2})^{S_n}) \dim((W^{\otimes 2})^{S_n})}. \end{aligned}$$

Specializing to our setting, we conclude that it suffices to show that

$$(\dim(H^i(X(\mathbb{C})^n; \mathbb{Q})^{\otimes 2}))^{S_{n-a}} \cdot (\dim(A(n, m)^{\otimes 2})^{S_{n-a}}) \leq F_a(i)$$

for some $F_a(i)$ subexponential in i . For the first factor, by Künneth and the definition of the action, we have

$$\begin{aligned} (H^i(X^n; \mathbb{Q})^{\otimes 2})^{S_{n-a}} &\subset H^{2i}(X^n \times X^n; \mathbb{Q})^{S_{n-a}} \\ &\cong H^{2i}((X \times X)^n; \mathbb{Q})^{S_{n-a}} \\ &\cong \bigoplus_{j=0}^{2i} H^j((X \times X)^{n-a}; \mathbb{Q})^{S_{n-a}} \otimes H^{2i-j}((X \times X)^a; \mathbb{Q}). \end{aligned}$$

By transfer, this is isomorphic to

$$\bigoplus_{j=0}^{2i} H^j(\mathrm{Sym}^{n-a}(X \times X); \mathbb{Q}) \otimes H^{2i-j}((X \times X)^a; \mathbb{Q})$$

Let $C = \max_i b_i(X \times X)$, and let $D = C \cdot 2m$. By Künneth, for all $j < 2i$,

$$\dim H^{2i-j}(X^{2a}; \mathbb{Q}) \leq (2i - j)C^a.$$

Combining this with Proposition 3.2, we see that

$$\begin{aligned} \dim(H^i(X^n; \mathbb{Q})^{\otimes 2})^{S_{n-a}} &\leq \sum_{j=0}^{2i} \dim(H^j(\mathrm{Sym}^{n-a}(X \times X); \mathbb{Q})) (2i - j)C^a \\ &\leq \sum_{j=0}^{2i} 2^D D! P(j)(2i - j)C^a \\ &\leq 2^D D! C^a Q(i) \end{aligned}$$

for some polynomials P, Q . It remains to bound $\dim(A(n, m) \otimes A(n, m))_i^{S_{n-a}}$. Well,

$$(A(n, m) \otimes A(n, m))_i = \bigoplus_{p+q=i} (A(n, m)_p \otimes A(n, m)_q) \quad (3.5)$$

Since the right-hand side of (3.5) has at most $2i$ terms, it suffices to bound each $[A(n, m)_p \otimes A(n, m)_q]^{S_{n-a}}$. By the Cauchy-Schwartz inequality, as above, it suffices to bound $[A(n, m)_p \otimes A(n, m)_p]^{S_{n-a}}$ for each $1 \leq p \leq i$. To obtain this bound, first note that the algebra $A(n, m)$ is isomorphic to $A(n, 2)$ via an isomorphism that takes the p^{th} graded piece of $A(n, 2)$ to the $(2m-1)p^{\text{th}}$ graded piece of $A(n, m)$. Since m is fixed and so $2m-1$ is fixed, it suffices to bound $[A(n, 2)_p \otimes A(n, 2)_p]^{S_{n-a}}$ in terms of i , for each $1 \leq p \leq i$.

Lehrer-Solomon [LS] give an explicit description of $A(n, 2)$ as a sum of induced representations

$$A(n, 2)_p = \bigoplus_{\mu} \mathrm{Ind}_{Z(c_{\mu})}^{S_n}(\xi_{\mu})$$

where μ runs over the set of conjugacy classes in S_n of permutations having $n-p$ cycles, c_{μ} is any element of the conjugacy class μ , and ξ_{μ} is a one-dimensional character of the centralizer $Z(c_{\mu})$ of c_{μ} in S_n (we will not need an explicit description of ξ_{μ}). It follows that

$$(A(n, 2)_p \otimes A(n, 2)_p)^{S_{n-a}} = \bigoplus_{\mu, \nu} [\mathrm{Ind}_{Z(c_{\mu})}^{S_n}(\xi_{\mu}) \otimes \mathrm{Ind}_{Z(c_{\nu})}^{S_n}(\xi_{\nu})]^{S_{n-a}} \quad (3.6)$$

where ν is defined similarly to μ . The summands contributing to the first (resp. second) $A(n, 2)_p$ factor in (3.6) correspond to conjugacy classes c_{μ} (resp. c_{ν}) in S_n decomposing into $n-p$ cycles. The number of such conjugacy classes is in bijection with the set of partitions of p , which is less than the number of partitions of i since $p \leq i$. The Hardy-Ramanujan asymptotic for the number $|\{J \vdash i\}|$ of partitions of i gives $C_1, C_2 > 0$ so that

$$|\{J \vdash i\}| \leq C_1 e^{C_2 \sqrt{i}}. \quad (3.7)$$

Thus the number of terms in the sum on the right-hand side of (3.6) is, by (3.7), at most $[C_1 e^{C_2 \sqrt{i}}]^2 = C_1^2 e^{2C_2 \sqrt{i}}$. As this is subexponential in i , it suffices to bound the dimension of $[\mathrm{Ind}_{Z(c_{\mu})}^{S_n}(\xi_{\mu}) \otimes \mathrm{Ind}_{Z(c_{\nu})}^{S_n}(\xi_{\nu})]^{S_{n-a}}$.

Now, a permutation c_μ decomposing into $n - p$ cycles must have at least $n - 2p$ fixed points. This implies that the centralizer $Z(c_\mu)$ contains the subgroup S_{n-2p} , and thus S_{n-2i} since $p \leq i$. It follows that $\text{Ind}_{Z(c_\mu)}^{S_n}(\xi_\mu)$ is a subrepresentation of $\text{Ind}_{S_{n-2i}}^{S_n}(\xi_\mu)$. Thus

$$[\text{Ind}_{Z(c_\mu)}^{S_n}(\xi_\mu) \otimes \text{Ind}_{Z(c_\nu)}^{S_n}(\xi_\nu)]^{S_{n-a}} \subset [\text{Ind}_{S_{n-2i}}^{S_n}(\xi_\mu) \otimes \text{Ind}_{S_{n-2i}}^{S_n}(\xi_\nu)]^{S_{n-a}} \quad (3.8)$$

Let χ_μ and χ_ν denote the characters of ξ_μ and ξ_ν , respectively. The right-hand side of (3.8) consists of the set of bilinear functions $f : S_n \times S_n \rightarrow \mathbb{C}$ satisfying

$$f(\sigma \cdot g, \tau \cdot h) = \chi_\mu(\sigma)\chi_\nu(\tau)f(g, h) \quad \forall \sigma, \tau \in S_{n-2i} \text{ and } \forall g, h \in S_n$$

and

$$f(g \cdot \beta, h \cdot \beta) = f(g, h) \quad \forall \beta \in S_{n-a} \text{ and } \forall g, h \in S_n.$$

It follows that the dimension of this vector space is at most the number of double cosets

$$S_{n-a} \backslash [S_n/S_{n-2i} \times S_n/S_{n-2i}].$$

We claim that this number is polynomial in i . Indeed, it is equal to the number of maps $f : \{1, \dots, a\} \rightarrow \{1, \dots, 2i, \star\} \times \{1, \dots, 2i, \star\}$ such that $|f^{-1}(j, k)| \leq 1$ and $|f^{-1}(j, \star)|, |f^{-1}(\star, k)| \leq (n-2i)^2$. Since a is fixed, this number is bounded by a constant times the number of subsets of $\{1, \dots, 2i, \star\} \times \{1, \dots, 2i, \star\}$ of size $\leq a$, which is $O(i^{2a})$. This completes the proof of Theorem 3.4. \square

4 Stability of arithmetic statistics

Throughout this section we will fix a prime power $q = p^d$ and a prime ℓ not divisible by p .

4.1 Point counting and étale cohomology

Let Y be a scheme of finite type (not necessarily smooth) over $\mathbb{Z}[1/N]$. We can base change to $\text{Spec}(\mathbb{F}_p)$ for any prime $p \nmid N$, and for any positive power $q = p^d$ we can consider both the \mathbb{F}_q -points as well as the $\overline{\mathbb{F}}_q$ -points of Y , where $\overline{\mathbb{F}}_q$ is the algebraic closure of \mathbb{F}_q . The *arithmetic Frobenius* morphism $\text{Frob}_q : Y \rightarrow Y$ acts on $Y(\overline{\mathbb{F}}_q)$ by acting on the coordinates (y_1, \dots, y_d) of any affine chart of y via

$$\text{Frob}_q(y_1, \dots, y_d) := (y_1^q, \dots, y_d^q).$$

A point $y \in Y(\overline{\mathbb{F}}_q)$ will be fixed by Frob_q precisely when $y \in Y(\mathbb{F}_q)$. Thus

$$Y(\mathbb{F}_q) = \text{Fix}(\text{Frob}_q : Y(\overline{\mathbb{F}}_q) \rightarrow Y(\overline{\mathbb{F}}_q)).$$

Fix a prime ℓ not dividing q , and let \mathbb{Q}_ℓ denote the ℓ -adic rationals. Let $H_{et}^*(Y_{/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$ (resp. $H_{et,c}^*(Y_{/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$) denote the étale cohomology groups (resp. compactly supported étale cohomology groups) of the base change $Y_{/\overline{\mathbb{F}}_q}$ of Y to $\overline{\mathbb{F}}_q$ (see, e.g., [De2, Mi]). Denote by $\mathbb{Q}_\ell(-i)$ the rank 1 $\text{Gal}(\overline{\mathbb{F}}_q/\mathbb{F}_q)$ -representation on which Frobenius acts by q^i .

Let \mathcal{V} be a constructible, rational ℓ -adic sheaf on Y (see, e.g., [FK]). If $y \in Y(\overline{\mathbb{F}}_q)$ is a fixed point for the action of Frob_q , then Frob_q acts on the stalk \mathcal{V}_y over y . Attached to this

action is its trace $\text{Tr}(\text{Frob}_q : \mathcal{V}_y \longrightarrow \mathcal{V}_y)$. The *twisted Grothendieck–Lefschetz Trace Formula* ([FK, Theorem II.3.14] and [De2, 6.1.1.1]) gives:

$$\sum_{y \in Y(\mathbb{F}_q)} \text{Tr}(\text{Frob}_q : \mathcal{V}_y \longrightarrow \mathcal{V}_y) = \sum_{i=0}^{2 \dim(Y)} (-1)^i \text{Tr}(\text{Frob}_q : H_{et,c}^i(Y; \mathcal{V}) \longrightarrow H_{et,c}^i(Y; \mathcal{V})) \quad (4.1)$$

When Y is smooth, Poincaré duality for étale cohomology [Mi, Theorem 24.1] gives

$$H_{et,c}^i(Y_{/\overline{\mathbb{F}}_q}; \mathcal{V}) \cong H_{et}^{2 \dim(Y)-i}(Y_{/\overline{\mathbb{F}}_q}; \mathcal{V}(-\dim(Y)))^*. \quad (4.2)$$

Plugging this into Equation (4.1) gives, for smooth Y :

$$\sum_{y \in Y(\mathbb{F}_q)} \text{Tr}(\text{Frob}_q : \mathcal{V}_y \longrightarrow \mathcal{V}_y) = q^{\dim(Y)} \sum_{i=0}^{2 \dim(Y)} (-1)^i \text{Tr}(\text{Frob}_q : H_{et}^i(Y; \mathcal{V})^* \longrightarrow H_{et}^i(Y; \mathcal{V})^*) \quad (4.3)$$

S_n -schemes. Now let Z be smooth and quasi-projective over $\mathbb{Z}[1/N]$. Suppose that the symmetric group S_n acts generically freely on Z by automorphisms, and let $p : Z \longrightarrow Y$ denote the quotient map. By [Mu, Theorem p. 63 and Remark p. 65 (Ch. 2.7)], Y is a scheme. It is typically not smooth even when Z is smooth.

Recall that any finite-dimensional representation of S_n over a field of characteristic 0 is defined over \mathbb{Q} . There is a bijective correspondence between isomorphism classes of finite-dimensional S_n -representations and finite-dimensional constructible sheaves on Y that become isomorphic to $\mathbb{Q}_\ell^{\oplus n}$ when pulled back to Z : Given an S_n -representation V over \mathbb{Q}_ℓ , one can form an S_n -equivariant, locally constant sheaf \mathcal{V} over Z with fiber V . Pushing forward to Y and taking S_n invariants, i.e. $(p_* \mathcal{V})^{S_n}$, we obtain a constructible sheaf of \mathbb{Q}_ℓ vector spaces over Y which is a sheaf-theoretic analogue of the usual topological diagonal quotient “ $Z \times_{S_n} V$ ”.

Suppose that $y \in Y(\overline{\mathbb{F}}_q)$ is fixed by Frob_q . Then Frob_q acts on the fiber $p^{-1}(y)$. Now S_n acts transitively on $p^{-1}(y)$ with some stabilizer H (not depending on $\tilde{y} \in p^{-1}(y)$), and so we can identify $p^{-1}(y)$ with S_n/H . The Frob_q action on $p^{-1}(y)$ commutes with this S_n action, and so it is determined by its action on a single basepoint, which we choose once and for all to be H . Now $\text{Frob}_q(H) = \sigma_y H$ for $\sigma_y \in S_n$. Following Gadish [Ga], for any S_n -representation V and any coset σH of S_n , we set

$$\chi_V(\sigma H) := \frac{1}{|H|} \sum_{h \in H} \chi_V(\sigma h).$$

With this notation we have:

$$\text{Tr}(\text{Frob}_q : \mathcal{V}_y \longrightarrow \mathcal{V}_y) = \chi_V(\sigma_y H) \quad (4.4)$$

which we denote simply by $\chi_V(\text{Frob}_q; \mathcal{V}_y)$. More generally:

Definition 4.1. For any class function P , and any $y \in Y$ fixed by Frob_q , define $P(y)$ by:

$$P(y) := \frac{1}{|H|} \sum_{h \in H} P(\sigma_y h). \quad (4.5)$$

An elementary check shows that the definitions above are independent of the choice of coset H , since the action of S_n is transitive on fibers.

Plugging Equation (4.4) into Equation (4.1) now gives:

$$\sum_{y \in Y(\mathbb{F}_q)} \chi_V(\text{Frob}_q; \mathcal{V}_y) = \sum_{i=0}^{2 \dim(Y)} (-1)^i \text{Tr}(\text{Frob}_q : H_{et,c}^i(Y; \mathcal{V}) \longrightarrow H_{et,c}^i(Y; \mathcal{V})) \quad (4.6)$$

The right-hand side of (4.6) could be computed from the eigenvalues λ_{ij} of Frob_q on each $H_{et,c}^i(Y; \mathbb{Q}_\ell)$. Typically one only has estimates on $|\lambda_{ij}|$. For example, for Y smooth and proper, the Riemann Hypothesis for finite fields (proved by Deligne) gives that $|\lambda_{ij}| = q^{i/2}$. Many natural examples Y , including many of those we study in this paper, are not proper, and finding the λ_{ij} is more difficult.

Given that we only have general bounds on the eigenvalues of Frob_q , to bound the traces of Frob_q we must determine the dimensions of each $H_{et,c}^i(Y; \mathcal{V})$. To do this, we follow the argument in §3.3 of [CEF2]. First note that the pullback $\tilde{\mathcal{V}}$ of \mathcal{V} to Z is trivial. We then compute:

$$\begin{aligned} H_{et,c}^i(Y; \mathcal{V}) &\cong H_{et,c}^i(Z; \tilde{\mathcal{V}})^{S_n} && \text{by transfer} \\ &\cong (H_{et,c}^i(Z; \mathbb{Q}_\ell) \otimes V)^{S_n} && \text{by triviality of } \tilde{\mathcal{V}}|_Z \\ &\cong (H_{et}^{2 \dim(Z)-i}(Z; \mathbb{Q}_\ell(\dim(Z)))^* \otimes V)^{S_n} && \text{by Poincaré duality} \\ &\cong H_{et}^{2 \dim(Z)-i}(Z; \mathbb{Q}_\ell(\dim(Z)))^* \otimes_{\mathbb{Q}_\ell[S_n]} V \end{aligned} \quad (4.7)$$

Because every S_n -representation is self-dual, it follows that

$$\dim_{\mathbb{Q}_\ell} H_{et,c}^i(Y; \mathcal{V}) = \langle V, H_{et}^{2 \dim(Z)-i}(Z; \mathbb{Q}_\ell) \rangle_{S_n} \quad (4.8)$$

where $\langle V, W \rangle_{S_n}$ is the usual inner product of S_n -representations V and W :

$$\langle V, W \rangle = \dim_{\mathbb{Q}_\ell} \text{Hom}_{\mathbb{Q}_\ell[S_n]}(V, W).$$

4.2 Co-FI schemes with convergent étale cohomology

Now that we have discussed schemes, and S_n -schemes, we are ready to discuss sequences of S_n -schemes.

Let Z be a co-FI scheme, smooth and quasi-projective over $\mathbb{Z}[1/N]$. For each $i \geq 0$, the étale cohomology $H_{et}^i(Z_\bullet/\mathbb{F}_q; \mathbb{Q}_\ell)$ is an FI-module over \mathbb{Q}_ℓ . We want to consider the implications of finite generation of this FI-module for point-counting problems over \mathbb{F}_q for the sequence of schemes Z_n/S_n (cf. [Mu, Theorem p. 63 and Remark p. 65]).

As discussed in [CEF1], any partition λ of any $k \geq 1$ determines a finitely-generated FI-module $V(\lambda)$ with $V(\lambda)_n$ being the irreducible representation of S_n corresponding to the partition $(n - |\lambda|) + \lambda$.

Definition 4.2. Let K be a field, and let M_\bullet be a finitely generated $\text{Gal}(\overline{K}/K)$ -FI-module over \mathbb{Q}_ℓ with stable range N . Let λ be a partition of n , let $V(\lambda)$ the associated FI-module, and let $D = \max\{N, \lambda_1\}$. Define the stable λ -isotypic part M_λ of M to be the $\text{Gal}(\overline{\mathbb{F}}_q/\mathbb{F}_q)$ -module

$$M_\lambda := M_D \otimes_{\mathbb{Q}_\ell[S_D]} V(\lambda)_D.$$

More generally, for a character polynomial P , we define the stable P -isotypic part of M to be the \mathbb{Q}_ℓ -virtual Galois module M_P obtained as a linear combination of the M_λ , with the sum taken in the representation ring of $\text{Gal}(\overline{K}/K)$ with \mathbb{Q}_ℓ -coefficients.

Lemma 2.2 shows that for $n \geq D$, there are canonical Galois-equivariant isomorphisms

$$M_n \otimes_{\mathbb{Q}_\ell[S_n]} V(\lambda)_n \xrightarrow{\cong} M_{n+1} \otimes_{\mathbb{Q}_\ell[S_{n+1}]} V(\lambda)_{n+1}$$

and similarly for the stable P -isotypic parts for $n \geq D$.

We can now give the following theorem, which generalizes earlier special cases by Ellenberg-Venkatesh-Westerland [EVW], Ellenberg [E], and Church-Ellenberg-Farb [CEF2]. Its proof is along the exact same lines of the previous proofs. We hope that the generality of the statement here will be useful in future work.

Theorem 4.3 (Convergent Grothendieck-Lefschetz). Let Z be a smooth, quasi-projective co-FI over \mathbb{F}_q , and set $Y_n := Z_n/S_n$ (we do not assume Y_n smooth over $\mathbb{Z}[1/N]$). Assume that for each $i \geq 0$ the FI-module $H_{et}^i(Z_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$ is finitely generated, and, for a character polynomial P , denote by $H_{et}^i(Z)_P^*$ the dual of the stable P -isotypic part. If Z has convergent étale cohomology over $\overline{\mathbb{F}}_q$, then for any character polynomial P :

$$\lim_{n \rightarrow \infty} q^{-\dim Y_n} \sum_{y \in Y_n(\mathbb{F}_q)} P(y) = \sum_{i=0}^{\infty} (-1)^i \text{Tr} (\text{Frob}_q \circ H_{et}^i(Z)_P^*), \quad (4.9)$$

and, taking the absolute value :

$$\lim_{n \rightarrow \infty} q^{-\dim Y_n} \left| \sum_{y \in Y_n(\mathbb{F}_q)} P(y) \right| \leq \sum_{i=0}^{\infty} \frac{\langle P, H_{et}^i(Z) \rangle}{q^{i/2}} < \infty. \quad (4.10)$$

If Z only has weakly convergent cohomology with convergence rate λ , then (4.9) and (4.10) hold for all $q > \lambda$.

Remark 4.4.

1. Specializing the Theorem 4.3 to the case $P = 1$ gives

$$\lim_{n \rightarrow \infty} q^{-\dim Y_n} |Y_n(\mathbb{F}_q)| = \sum_{i=0}^{\infty} (-1)^i \text{Tr} (\text{Frob}_q \circ H_{et}^i(Y)^*), \quad (4.11)$$

where $H_{et}^i(Y)^*$ denotes the stable rational étale cohomology of the sequence Y_1, Y_2, \dots

2. The bound (4.10) is sharp, as is seen by taking $Z_n = (\mathbb{P}^1)^n$, $Y_n = \mathbb{P}^n$, and $P = 1$.

Proof of Theorem 4.3. Because all of the equations in the statement of the theorem are \mathbb{Q}_ℓ -linear in P , it suffices to prove the theorem for $P = P_\lambda$, the character polynomial of the finitely generated FI-module $V(\lambda)$ (cf. Theorem 2.3 above). Let \mathcal{V}_n correspond to the twisted sheaf on Y_n corresponding to the representation $V(\lambda)_n$.

We show that the left side of (4.9) converges by showing that the sequences

$$n \mapsto q^{-\dim Y_n} \sum_{y \in Y_n(\mathbb{F}_q)} P(y) \quad (4.12)$$

is Cauchy. To start, note that

$$\begin{aligned} \sum_{y \in Y_n(\mathbb{F}_q)} P(y) &= \sum_{i=0}^{2 \dim(Y_n)} (-1)^i \text{Tr} (\text{Frob}_q : H_{et,c}^i(Y_n; \mathcal{V}_n) \longrightarrow H_{et,c}^i(Y_n; \mathcal{V}_n)) \\ &= \sum_{i=0}^{2 \dim(Z_n)} (-1)^i \text{Tr} (\text{Frob}_q \circ H_{et}^{2 \dim(Z_n)-i}(Z_n; \mathbb{Q}_\ell(\dim(Z_n)))^* \otimes_{\mathbb{Q}_\ell[S_n]} V(\lambda)_n) \\ &\quad \text{(by Equation (4.7))} \\ &= \sum_{i=0}^{2 \dim(Z_n)} (-1)^i \text{Tr} (\text{Frob}_q \circ H_{et}^i(Z_n; \mathbb{Q}_\ell(\dim(Z_n)))^* \otimes_{\mathbb{Q}_\ell[S_n]} V(\lambda)_n^*) . \end{aligned} \quad (4.13)$$

where the last equation uses the self-duality of S_n -representations.

Denote by $N(n, P)$ the slope of stability of $H_{et}^*(Z_\bullet; \mathbb{Q}_\ell)$ for $V(\lambda)$, i.e. the number such that for all $i \leq N(n, \lambda)$,

$$H_{et}^i(Z_n; \mathbb{Q}_\ell) \otimes_{\mathbb{Q}_\ell[S_n]} V(\lambda)_n \cong H_{et}^i(Z)_P .$$

Let $F_P(i)$ denote the subexponential function in Definition 3.1 guaranteed by the assumption that Z has convergent étale cohomology. Then, for $n > m$:

$$\begin{aligned} &|q^{-\dim Y_n} (\sum_{y \in Y_n(\mathbb{F}_q)} P(y)) - q^{-\dim Y_m} (\sum_{y \in Y_m(\mathbb{F}_q)} P(y))| \\ &= | \sum_{i=0}^{2 \dim(Z_n)} (-1)^i q^{-\dim(Z_n)} \text{Tr} (\text{Frob}_q \circ H_{et}^i(Z_n; \mathbb{Q}_\ell(\dim(Z_n)))^* \otimes_{\mathbb{Q}_\ell[S_n]} V(\lambda)_n^*) \\ &\quad - \sum_{i=0}^{2 \dim(Z_m)} (-1)^i q^{-\dim(Z_m)} \text{Tr} (\text{Frob}_q \circ H_{et}^i(Z_m; \mathbb{Q}_\ell(\dim(Z_m)))^* \otimes_{\mathbb{Q}_\ell[S_m]} V(\lambda)_m^*) | \\ &\quad \text{(by Equations (4.1) and (4.13))} \\ &\leq \sum_{i=0}^{\infty} q^{-i/2} |\langle P, H_{et}^i(Z_n; \mathbb{Q}_\ell) \rangle - \langle P, H_{et}^i(Z_m; \mathbb{Q}_\ell) \rangle| \quad \text{(by Deligne)} \\ &= \sum_{i=N(m, P)}^{\infty} q^{-i/2} |\langle P, H_{et}^i(Z_n; \mathbb{Q}_\ell) \rangle - \langle P, H_{et}^i(Z_m; \mathbb{Q}_\ell) \rangle| \\ &\quad \text{(by étale representation stability)} \\ &\leq \sum_{i=N(m, P)}^{\infty} 2q^{-i/2} F_P(i) . \quad \text{(by convergent cohomology)} \end{aligned}$$

Because $N(m, P)$ tends to ∞ with m , and because $F_P(i)$ is sub-exponential in i , we see that the sequence (4.12) is Cauchy. Similarly, we see that the right side of (4.9)

$$\sum_{i=0}^{\infty} (-1)^i \text{Tr}(\text{Frob}_q \odot H_{et}^i(Z)_P^*)$$

converges as a consequence of the existence of the stable P -isotypic part, Deligne's bounds on the eigenvalues of Frob_q and the existence of the sub-exponential bounds $F_P(i)$.

It remains to show that the two limits agree. For this, we have

$$\begin{aligned} & |q^{-\dim Y_n} \sum_{y \in Y_n(\mathbb{F}_q)} P(y) - \sum_{i=0}^{2 \dim(Z_n)} (-1)^i \text{Tr}(\text{Frob}_q \odot H_{et}^i(Z)_P^*)| \\ &= \left| \sum_{i=0}^{2 \dim(Z_n)} (-1)^i (\text{Tr}(\text{Frob}_q \odot H_{et}^i(Z_n; \mathbb{Q}_\ell(\dim(Z_n)))^* \otimes_{\mathbb{Q}_\ell[S_n]} V(\lambda)_n^*)) \right. \\ &\quad \left. - \text{Tr}(\text{Frob}_q \odot H_{et}^i(Z)_P^*) \right| \quad (\text{by Equation (4.13)}) \\ &= \left| \sum_{i=N(n,P)+1}^{2 \dim(Z_n)} (-1)^i (\text{Tr}(\text{Frob}_q \odot H_{et}^i(Z_n; \mathbb{Q}_\ell(\dim(Z_n)))^* \otimes_{\mathbb{Q}_\ell[S_n]} V(\lambda)_n^*)) \right. \\ &\quad \left. - \text{Tr}(\text{Frob}_q \odot H_{et}^i(Z)_P^*) \right| \quad (\text{by \'etale representation stability}) \\ &\leq \sum_{i=N(n,P)+1}^{2 \dim(Z_n)} 2q^{-i/2} F_P(i). \quad (\text{by Deligne and convergent cohomology}) \end{aligned} \tag{4.14}$$

Because $F_P(i)$ is subexponential in i , we conclude that (4.14) becomes arbitrarily small as n approaches ∞ , which proves the theorem. \square

We can now prove Theorem A from the introduction, as well as the following.

Theorem 4.5 (Statistics for $\text{Sym}^n X$). *Theorem A with $\text{PConf}_n(Y)$ (resp. $\text{UConf}_n(Y)$) replaced by Y^n (resp. $\text{Sym}^n(Y)$) holds.*

Proof of Theorem A and Theorem 4.5. Theorem B gives that, for each $i \geq 0$, the FI-modules $H_{et}^*(X^\bullet; \mathbb{Q}_\ell)$ and $H_{et}^*(\text{PConf}_\bullet(X); \mathbb{Q}_\ell)$ are finitely generated. Corollary 3.3 (resp. Theorem 3.4) gives that the singular cohomology $H^*(X^\bullet; \mathbb{Q})$ (resp. $H^*(\text{PConf}_\bullet(X); \mathbb{Q})$) of the co-FI scheme X^\bullet (resp. $\text{PConf}_\bullet(X)$) is convergent. Now apply Theorem 4.3. \square

In special cases it is possible to compute the right hand side of Equation (4.9) explicitly.

Example 4.6. When $X = \mathbb{A}^r$, we can explicitly compute polynomial statistics on $\text{UConf}_n(\mathbb{A}^r)$, extending the main theorem of [CEF2]. Indeed, the computations of Arnol'd [Ar] and F. Cohen [Coh, §2] combine with results of Björner–Ekedahl [BE, Theorem 4.9] to show that $H_{et}^*(\text{PConf}_n(\mathbb{A}^r)_{\mathbb{F}_q}; \mathbb{Q}_\ell)$ is a graded algebra generated by classes in degree $2r - 1$ with eigenvalues of Frob_q equal to q^r . As a result, for any character polynomial P_λ :

$$\text{Tr}(\text{Frob}_q: H_{et}^i(\text{UConf}_n(\mathbb{A}^r); \mathcal{V})^* \longrightarrow H_{et}^i(\text{UConf}_n(\mathbb{A}^r); \mathcal{V})^*)$$

$$= \begin{cases} 0 & \text{if } i \neq k(2r - 1) \\ q^{-kr} \langle P_\lambda, H_{et}^i(\text{PConf}_n(\mathbb{A}^r); \mathbb{Q}_\ell) \rangle & \text{if } i = k(2r - 1) \end{cases}$$

Here we have, as above, applied Poincarè duality to replace the compactly supported cohomology of the smooth schemes $\mathrm{UConf}_n(\mathbb{A}^r)$ with (the Tate twist of) the dual of ordinary étale cohomology. We thus have, for all P ,

$$\lim_{n \rightarrow \infty} q^{-nr} \sum_{y \in \mathrm{UConf}_n(\mathbb{A}^r)(\mathbb{F}_q)} P(y) = \sum_{i=0}^{\infty} (-1)^{i(2r-1)} q^{-ir} \langle P, H_{et}^{i(2r-1)}(\mathrm{PConf}(\mathbb{A}^r)) \rangle.$$

References

- [Ar] V.I. Arnol'd, The cohomology ring of the colored braid group, *Mathematical Notes* 5, no. 2 (1969), 138–140.
- [BE] A. Björner and T. Ekedahl, Subspace arrangements over finite fields: cohomological and enumerative aspects, *Advances in Math.* 129 (1997), pp. 159–187.
- [CEF1] T. Church, J.S. Ellenberg and B. Farb, FI-modules: a new approach to stability for S_n -representations, *Duke Math Journal*, Vol. 164, No. 9 (2015), pp. 1833–1910.
- [CEF2] T. Church, J.S. Ellenberg and B. Farb, Representation stability in cohomology and asymptotics for families of varieties over finite fields, to appear in *Algebraic Topology: Applications and New Directions*, AMS Contemp. Math. Series., Vol. 620 (2014), pp. 1–54.
- [CEFN] T. Church, J.S. Ellenberg, B. Farb and R. Nagpal, FI-modules over Noetherian rings, *Geometry & Topology*, Vol. 18, No. 5 (2014), pp. 2951–2984.
- [CF] T. Church and B. Farb, Representation theory and homological stability, *Advances in Math.*, Vol. 245 (2013), pp. 250–314.
- [Che] W. Chen, Twisted cohomology of configuration spaces and spaces of maximal tori via point-counting, preprint, March 2016, arXiv:1603.03931v1.
- [Ch] T. Church, Homological stability for configuration spaces of manifolds, *Invent. Math.* 188 (2012) 2, 465–504.
- [Coh] F. Cohen, On configuration spaces, their homology, and Lie algebras, *Jour. of Pure and Appl. Alg.* 100 (1995), 19–42.
- [De1] P. Deligne, La conjecture de Weil, II, *Inst. Hautes Études Sci. Publ. Math.*, No. 52(1980), pp.137–252.
- [De2] P. Deligne, Cohomologie à supports propres, *Theorie des Topos et Cohomologie Étale des Schémas* (SGA 4), Lect. Notes in Math., Vol. 269 (1972).
- [E] J. Ellenberg, Arizona Winter School 2014 notes, <http://swc.math.arizona.edu/aws/2014/2014EllenbergNotes.pdf>.
- [EVW] J. Ellenberg, A. Venkatesh, and C. Westerland, Homological stability and Cohen-Lenstra over function fields, *Annals of Math.*, 183 (2016), no. 3, 729–786.
- [Fa] B. Farb, Representation Stability, *Proc. of the ICM, Seoul, 2014*.
- [FK] E. Freitag and R. Kiehl, *Étale Cohomology and the Weil Conjectures*, Springer-Verlag, 1980.
- [FS] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge Univ. Press, 2009.

- [FM] W. Fulton and R. MacPherson, A compactification of configuration spaces, *Ann. of Math.* (2) **139** (1994), no. 1, 183–225.
- [Ga] N. Gadish, Representation Stability for Families of Linear Subspace Arrangements, preprint, March 2016, arXiv:1603.08547v3.
- [HK] M.L. Hautus and D.A. Klarner, The diagonal of a double power series, *Duke Math. Jour.* Vol. 38, NO. 2 (1971), 229–235.
- [Ho] Q. Ho, Free factorization algebras and homology of configuration spaces in algebraic geometry, preprint, December 2015, arXiv:1512.04490v1.
- [Hor] G. Horel, Motivic homological stability for configurations spaces of the line, preprint, arXiv:1511.09031v3, May 2016.
- [LS] G. Lehrer and L. Solomon, On the action of the symmetric group on the cohomology of the complement of its reflecting hyperplanes, *J. Algebra* 104 (1986), no. 2, 410–424.
- [LT] M. Lipnowski and J. Tsimerman, How large is $A_g(\mathbb{F}_q)$?, arXiv:1511.02212v1, November, 2015.
- [Mac] I.G. Macdonald, The Poincarè polynomial of a symmetric product, *Proc. Cambridge Philos. Soc.* **58** (1962), 563–568.
- [Mi] J. Milne, *Lectures on étale cohomology*, Version 2.21., March 22 2013.
- [Mi2] J. Milne, Values of Zeta Functions of Varieties over Finite Fields, *Amer. J. Math.*, vol. 108 (1986) pp. 297–360.
- [Mu] D. Mumford, *Abelian Varieties*, 2nd ed. (1974).
- [Pe] D. Petersen, A spectral sequence for stratified spaces and configuration spaces of points, preprint, arXiv:1603.01137, March 2016.
- [Se] G. Segal, The Topology of the Space of Rational Functions, *Acta Math.*, Vol. 143 (1979), pp. 39–72.
- [To] B. Totaro, Configuration spaces of algebraic varieties, *Topology* 35 (1996), no. 4, 1057–1067.
- [VW] R. Vakil and M. Matchett Wood, Discriminants in the Grothendieck ring, *Duke Math. Journal*, 164 (2015), no. 6, 1139–1185.
- [We] A. Weber, The Leray spectral sequence for complements of certain arrangements of smooth submanifolds, arXiv:1505.01365v1, May 6, 2015.

Dept. of Mathematics, University of Chicago
E-mail: farb@math.uchicago.edu, wolfson@math.uchicago.edu