

The Rayleigh-Taylor condition for the evolution of irrotational fluid interfaces.

Antonio Cordoba
Universidad Autonoma de Madrid
ctra. de Colmenar Viejo, Km. 15
28049 Madrid
e-mail: antonio.cordoba@uam.es

Diego Cordoba (Corresponding author)
Instituto de Ciencias Matematicas
Consejo Superior de Investigaciones Cientificas
c/Serrano 123
Madrid, 28006
e-mail: dcg@imaff.cfmac.csic.es

Francisco Gancedo
University of Chicago
5734 University Avenue,
Chicago, IL 60637
e-mail: fgancedo@math.uchicago.edu

Classification: Physical Sciences, Mathematics

May 14, 2009

Abstract

For the free boundary dynamics of the two-phase Hele-Shaw and Muskat problems, and also for the irrotational incompressible Euler equation, we prove existence locally in time when the Rayleigh-Taylor condition is initially satisfied for a 2D interface. The result for water waves was first obtained by Wu in a slightly different scenario (vanishing at infinity), but our approach is different because it emphasizes the active scalar character of the system and does not require the presence of gravity.

1 Introduction

There are several interesting problems in Fluid Mechanics regarding the evolution of the interface between two fluids (the Hele-Shaw cell [17, 22] and the Muskat problem [20]) or between a fluid and vacuum or another fluid with zero density, as in models of water waves. In all of them the first important question to be asked is in order to guarantee local existence, usually within the chain of Sobolev spaces. However such a result turns out to be false for general initial data. First Rayleigh [21], Taylor [26] and Saffman-Taylor [22], and later Beale-Hou-Lowengrub [4], Wu [27, 28], Christodoulou-Lindblad [7], Ambrose [1], Lindblad [19], Ambrose-Masmoudi [2], Coutand-Shkoller [12], Córdoba-Gancedo [11], Shatah-Zeng [23] and Zhang-Zhang [29] gave a condition that must be satisfied in order to have a solution locally in time, namely the normal component of the pressure gradient jump at the interface has to have a distinguished sign. This is known as the Rayleigh-Taylor condition.

In references [9] and [10] we have obtained local existence in the 2D case: for the Hele-Shaw and Muskat problems our result is new in the more difficult case when the two fluids have different densities and viscosities; for water waves we give a different proof of the important theorem of Wu [27] where gravity plays a crucial role in the sign of the Rayleigh-Taylor condition. In our proof, however, we consider the two cases, with or without gravity, and with initial data always satisfying the Rayleigh-Taylor condition.

When that condition is not imposed initially there are several cases where ill-posedness has been proved. Let us point out the works of Ebin [14, 15], Caffisch-Orellana [5], Siegel-Caffisch-Howison [24] and Córdoba-Gancedo [11].

We regard these models as transport equations for the density, considered as an active scalar, with a divergence free velocity field given by Darcy's law (Hele-Shaw and Muskat) or Bernoulli's law (irrotational incompressible Euler equation). It follows that the vorticity is then a delta distribution at the interface multiplied by an amplitude. The dynamics of that interface is governed by the Birkhoff-Rott integral of the amplitude from which we may subtract any component in the tangential direction without modifying its evolution (see [18]). We treat the case without surface tension which leads to equality of the pressure on the free boundary, and in both problems it is assumed that the initial interface does not self-intersect. We quantify that property by imposing that the arc-chord quotient be initially strictly positive. It is part of the evolution problem to check carefully that such a positivity prevails for a short time (see [16]), as does the Rayleigh-Taylor condition, although depending, in both cases, upon the initial data.

2 Equations

The free boundary is given by the discontinuity of the densities and the viscosities (in the case of the free boundary for the irrotational incompressible Euler equation the viscosity is zero) of the fluids

$$(\mu, \rho)(x_1, x_2, t) = \begin{cases} (\mu^1, \rho^1), & x \in \Omega^1(t) \\ (\mu^2, \rho^2), & x \in \Omega^2(t) = \mathbb{R}^2 - \Omega^1(t), \end{cases} \quad (1)$$

and $\mu_1, \mu_2, \rho_1, \rho_2$ ($\mu^1 \neq \mu^2$ and $\rho^1 \neq \rho^2$) are constants.

Let the free boundary be parameterized by

$$\partial\Omega^j(t) = \{z(\alpha, t) = (z_1(\alpha, t), z_2(\alpha, t)) : \alpha \in \mathbb{R}\}$$

where

$$(z_1(\alpha + 2k\pi, t), z_2(\alpha + 2k\pi, t)) = (z_1(\alpha, t) + 2k\pi, z_2(\alpha, t)), \quad (2)$$

with the initial data $z(\alpha, 0) = z_0(\alpha)$. We shall study also the case of a closed curve:

$$(z_1(\alpha + 2k\pi, t), z_2(\alpha + 2k\pi, t)) = (z_1(\alpha, t), z_2(\alpha, t)). \quad (3)$$

We consider that the velocity $v = v(x_1, x_2, t)$ is irrotational, i.e. $\omega = \nabla \times v = 0$, in the interior of each domain Ω^j ($j = 1, 2$). Therefore the vorticity ω has its support on the curve $z(\alpha, t)$ and it has the form

$$\omega(x, t) = \varpi(\alpha, t)\delta(x - z(\alpha, t))$$

i.e. ω is a measure defined by

$$\langle \omega, \eta \rangle = \int \varpi(\alpha, t)\eta(z(\alpha, t))d\alpha,$$

with $\eta(x)$ a test function.

Then $z(\alpha, t)$ evolves with a velocity field coming from Biot-Savart law, which can be explicitly computed; it is given by the Birkhoff-Rott integral of the amplitude ϖ along the interface curve:

$$BR(z, \varpi)(\alpha, t) = \frac{1}{2\pi} PV \int \frac{(z(\alpha, t) - z(\beta, t))^\perp}{|z(\alpha, t) - z(\beta, t)|^2} \varpi(\beta, t) d\beta, \quad (4)$$

where PV denotes principal value (see [25]).

We have

$$\begin{aligned} v^2(z(\alpha, t), t) &= BR(z, \varpi)(\alpha, t) + \frac{1}{2} \frac{\varpi(\alpha, t)}{|\partial_\alpha z(\alpha, t)|^2} \partial_\alpha z(\alpha, t), \\ v^1(z(\alpha, t), t) &= BR(z, \varpi)(\alpha, t) - \frac{1}{2} \frac{\varpi(\alpha, t)}{|\partial_\alpha z(\alpha, t)|^2} \partial_\alpha z(\alpha, t), \end{aligned} \quad (5)$$

where $v^j(z(\alpha, t), t)$ denotes the limit velocity field obtained approaching the boundary in the normal direction inside Ω^j and $BR(z, \varpi)(\alpha, t)$ is given by (4). It provides us the velocity field at the interface from which we can subtract any term in the tangential direction without modifying the geometric evolution of the curve

$$z_t(\alpha, t) = BR(z, \varpi)(\alpha, t) + c(\alpha, t)\partial_\alpha z(\alpha, t). \quad (6)$$

A wise choice of $c(\alpha, t)$ namely:

$$\begin{aligned} c(\alpha, t) &= \frac{\alpha + \pi}{2\pi} \int_{\mathbb{T}} \frac{\partial_\alpha z(\alpha, t)}{|\partial_\alpha z(\alpha, t)|^2} \cdot \partial_\alpha BR(z, \varpi)(\alpha, t) d\alpha \\ &\quad - \int_{-\pi}^{\alpha} \frac{\partial_\alpha z(\beta, t)}{|\partial_\alpha z(\beta, t)|^2} \cdot \partial_\beta BR(z, \varpi)(\beta, t) d\beta, \end{aligned} \quad (7)$$

allows us to accomplish the fact that the length of the tangent vector to $z(\alpha, t)$ only depends upon the variable t :

$$A(t) = |\partial_\alpha z(\alpha, t)|^2.$$

Next, in order to close the system we apply Darcy's law, or respectively Bernoulli's law for the case of Euler equations, which leads to an equation relating the parametrization $z(\alpha, t)$ with the amplitude $\varpi(\alpha, t)$.

2.1 Darcy's law

Darcy's law is the following momentum equation for the velocity v

$$\frac{\mu}{\kappa} v = -\nabla p - (0, g\rho), \quad (8)$$

where p is the pressure, μ is the dynamic viscosity, κ is the permeability of the medium, ρ is the liquid density and g is the acceleration due to gravity. Together with the incompressibility condition $\nabla \cdot v = 0$ equation (8) implies the identity

$$p^2(z(\alpha, t), t) = p^1(z(\alpha, t), t),$$

(see also [9]). Let us introduce the following notation:

$$[\mu v](\alpha, t) = (\mu^2 v^2(z(\alpha, t), t) - \mu^1 v^1(z(\alpha, t), t)) \cdot \partial_\alpha z(\alpha, t).$$

Then taking the limit in Darcy's law we obtain

$$\begin{aligned} \frac{[\mu v](\alpha, t)}{\kappa} &= -(\nabla p^2(z(\alpha, t), t) - \nabla p^1(z(\alpha, t), t)) \cdot \partial_\alpha z(\alpha, t) - g(\rho^2 - \rho^1) \partial_\alpha z_2(\alpha, t) \\ &= -\partial_\alpha (p^2(z(\alpha, t), t) - p^1(z(\alpha, t), t)) - g(\rho^2 - \rho^1) \partial_\alpha z_2(\alpha, t) \\ &= -g(\rho^2 - \rho^1) \partial_\alpha z_2(\alpha, t), \end{aligned}$$

which gives us

$$\frac{\mu^2 + \mu^1}{2\kappa} \varpi(\alpha, t) + \frac{\mu^2 - \mu^1}{\kappa} BR(z, \varpi)(\alpha, t) \cdot \partial_\alpha z(\alpha, t) = -g(\rho^2 - \rho^1) \partial_\alpha z_2(\alpha, t),$$

so that

$$\varpi(\alpha, t) = -A_\mu 2BR(z, \varpi)(\alpha, t) \cdot \partial_\alpha z(\alpha, t) - 2\kappa g \frac{\rho^2 - \rho^1}{\mu^2 + \mu^1} \partial_\alpha z_2(\alpha, t). \quad (9)$$

where $A_\mu = \frac{\mu_2 - \mu_1}{\mu_2 + \mu_1}$.

2.2 Bernoulli's law

In this section we deduce the evolution equation for the amplitude of vorticity $\varpi(\alpha, t)$ from Bernoulli's law. Let us consider an irrotational flow satisfying the Euler equations

$$\rho(v_t + v \nabla v) = -\nabla p - (0, g \rho), \quad (10)$$

and the incompressibility condition $\nabla \cdot v = 0$, and let ϕ be such that $v(x, t) = \nabla \phi(x, t)$ for $x \neq z(\alpha, t)$. Then we have the expression

$$\rho(\phi_t(x, t) + \frac{1}{2}|v(x, t)|^2 + gx_2) + p(x, t) = 0.$$

From the Biot-Savart law, for $x \neq z(\alpha, t)$, we get

$$\phi(x, t) = \frac{1}{2\pi} PV \int \arctan\left(\frac{x_2 - z_2(\beta, t)}{x_1 - z_1(\beta, t)}\right) \varpi(\beta, t) d\beta.$$

Let us define

$$\Pi(\alpha, t) = \phi^2(z(\alpha, t), t) - \phi^1(z(\alpha, t), t),$$

where again $\phi^j(z(\alpha, t), t)$ denotes the limit obtained approaching the boundary in the normal direction inside Ω^j . It is clear that

$$\begin{aligned} \partial_\alpha \Pi(\alpha, t) &= (\nabla \phi^2(z(\alpha, t), t) - \nabla \phi^1(z(\alpha, t), t)) \cdot \partial_\alpha z(\alpha, t) \\ &= (v^2(z(\alpha, t), t) - v^1(z(\alpha, t), t)) \cdot \partial_\alpha z(\alpha, t) \\ &= \varpi(\alpha, t), \end{aligned}$$

therefore

$$\int_{-\pi}^{\pi} \varpi(\alpha, t) d\alpha = 0.$$

Now we observe that

$$\begin{aligned} \phi^2(z(\alpha, t), t) &= IT(z, \varpi)(\alpha, t) + \frac{1}{2}\Pi(\alpha, t), \\ \phi^1(z(\alpha, t), t) &= IT(z, \varpi)(\alpha, t) - \frac{1}{2}\Pi(\alpha, t). \end{aligned} \quad (11)$$

where

$$IT(z, \varpi)(\alpha, t) = \frac{1}{2\pi} PV \int \arctan\left(\frac{z_2(\alpha, t) - z_2(\beta, t)}{z_1(\alpha, t) - z_1(\beta, t)}\right) \varpi(\beta, t) d\beta.$$

Then using Bernoulli's law inside each domain and taking limits approaching the common boundary, one finds

$$\rho^j(\phi_t^j(z(\alpha, t), t) + \frac{1}{2}|v^j(z(\alpha, t), t)|^2 + gz_2(\alpha, t)) + p^j(z(\alpha, t), t) = 0,$$

and since

$$p^1(z(\alpha, t), t) = p^2(z(\alpha, t), t)$$

(see also [10]), we get

$$[\rho\phi_t](\alpha, t) + \frac{\rho^2}{2}|v^2(z(\alpha, t), t)|^2 - \frac{\rho^1}{2}|v^1(z(\alpha, t), t)|^2 + (\rho^2 - \rho^1)gz_2(\alpha, t) = 0 \quad (12)$$

where we have introduced the following notation:

$$[\rho\phi_t](\alpha, t) = \rho^2\phi_t^2(z(\alpha, t), t) - \rho^1\phi_t^1(z(\alpha, t), t).$$

Then it is clear that $\phi_t^j(z(\alpha, t), t) = \partial_t(\phi^j(z(\alpha, t), t)) - z_t(\alpha, t) \cdot \nabla\phi^j(z(\alpha, t), t)$, and using (11) we find

$$\begin{aligned} [\rho\phi_t](\alpha, t) &= \frac{\rho^2 + \rho^1}{2}\Pi_t(\alpha, t) + (\rho^2 - \rho^1)\partial_t(IT(z, \varpi)(\alpha, t)) \\ &\quad - z_t(\alpha, t) \cdot (\rho^2v^2(z(\alpha, t), t) - \rho^1v^1(z(\alpha, t), t)). \end{aligned}$$

Equations (5) and (6) in (12) give us

$$\begin{aligned} \Pi_t(\alpha, t) &= -2A_\rho\partial_t(IT(z, \varpi)(\alpha, t)) + c(\alpha, t)\varpi(\alpha, t) + A_\rho|BR(z, \varpi)(\alpha, t)|^2 \\ &\quad + 2A_\rho c(\alpha, t)BR(z, \varpi)(\alpha, t) \cdot \partial_\alpha z(\alpha, t) - A_\rho \frac{|\varpi(\alpha, t)|^2}{4|\partial_\alpha z(\alpha, t)|^2} - 2A_\rho gz_2(\alpha, t). \end{aligned} \quad (13)$$

where $A_\rho = \frac{\rho^2 - \rho^1}{\rho^2 + \rho^1}$.

Easily we find the identity:

$$\begin{aligned} \partial_\alpha\partial_t(IT(z, \varpi)(\alpha, t)) &= \partial_t(BR(z, \varpi)(\alpha, t) \cdot \partial_\alpha z(\alpha, t)) \\ &= \partial_t(BR(z, \varpi)(\alpha, t)) \cdot \partial_\alpha z(\alpha, t) + BR(z, \varpi)(\alpha, t) \cdot \partial_\alpha BR(z, \varpi)(\alpha, t) \\ &\quad + c(\alpha, t)BR(z, \varpi)(\alpha, t) \cdot \partial_\alpha^2 z(\alpha, t) + \partial_\alpha c(\alpha, t)BR(z, \varpi)(\alpha, t) \cdot \partial_\alpha z(\alpha, t) \end{aligned}$$

Then taking a derivative in (13) and using the identity above we get the desired formula for ϖ , which in the case $A_\rho = 1$, i.e. $\rho_1 = 0$, reads as follows

$$\begin{aligned} \varpi_t(\alpha, t) &= -2\partial_t BR(z, \varpi)(\alpha, t) \cdot \partial_\alpha z(\alpha, t) - \partial_\alpha \left(\frac{|\varpi|^2}{4|\partial_\alpha z|^2} \right)(\alpha, t) + \partial_\alpha(c\varpi)(\alpha, t) \\ &\quad + 2c(\alpha, t)\partial_\alpha BR(z, \varpi)(\alpha, t) \cdot \partial_\alpha z(\alpha, t) + 2g\partial_\alpha z_2(\alpha, t). \end{aligned} \quad (14)$$

That is we have obtained the standard water waves model where g is the acceleration due to gravity.

3 Rayleigh-Taylor condition

Our next step is to find the formula for the difference of the gradients of the pressure in the normal direction:

$$\sigma(\alpha, t) = -(\nabla p^2(z(\alpha, t), t) - \nabla p^1(z(\alpha, t), t)) \cdot \partial_\alpha^\perp z(\alpha, t).$$

3.1 Darcy's law

Applying Darcy's law and approaching the boundary, we get

$$\sigma(\alpha, t) = \frac{\mu^2 - \mu^1}{\kappa} BR(z, \varpi)(\alpha, t) \cdot \partial_\alpha^\perp z(\alpha, t) + g(\rho^2 - \rho^1) \partial_\alpha z_1(\alpha, t). \quad (15)$$

3.2 Bernoulli's law

We will consider the case $A_\rho = 1$, which yields $-\nabla p(x, t) = 0$ inside $\Omega^1(t)$ and therefore $\nabla p^1(z(\alpha, t), t) = 0$. Next we define the Lagrangian coordinates for the free boundary with the velocity v^2

$$\begin{aligned} Z_t(\gamma, t) &= v^2(Z(\gamma, t), t) \\ Z(\gamma, 0) &= z_0(\gamma). \end{aligned}$$

We have the same curve with different parameterizations $Z(\gamma, t) = z(\alpha(\gamma, t), t)$ and two equations for the velocity of the curve, namely

$$\begin{aligned} Z_t(\gamma, t) &= z_t(\alpha, t) + \alpha_t(\gamma, t) \partial_\alpha z(\alpha, t) \\ &= BR(z, \varpi)(\alpha, t) + c(\alpha, t) \partial_\alpha z(\alpha, t) + \alpha_t(\gamma, t) \partial_\alpha z(\alpha, t) \end{aligned} \quad (16)$$

and another one given by

$$Z_t(\gamma, t) = BR(z, \varpi)(\alpha, t) + \frac{1}{2} \frac{\varpi(\alpha, t)}{|\partial_\alpha z(\alpha, t)|^2} \partial_\alpha z(\alpha, t). \quad (17)$$

Next we introduce the function (see [4] and [2])

$$\varphi(\alpha, t) = \frac{1}{2} \frac{\varpi(\alpha, t)}{|\partial_\alpha z(\alpha, t)|} - c(\alpha, t) |\partial_\alpha z(\alpha, t)|, \quad (18)$$

and we observe that the dot product of equations (17) and (16) with the tangential vector gives

$$\alpha_t(\gamma, t) = \frac{\varphi(\alpha, t)}{|\partial_\alpha z(\alpha, t)|}. \quad (19)$$

Taking a time derivative in (17) yields

$$\begin{aligned} Z_{tt}(\gamma, t) \cdot \partial_\alpha^\perp z(\alpha, t) &= (\partial_t BR(z, \varpi)(\alpha, t) + \alpha_t(\gamma, t) \partial_\alpha BR(z, \varpi)(\alpha, t)) \cdot \partial_\alpha^\perp z(\alpha, t) \\ &\quad + \frac{1}{2} \frac{\varpi(\alpha, t)}{|\partial_\alpha z(\alpha, t)|^2} (\partial_\alpha z_t(\alpha, t) + \alpha_t(\gamma, t) \partial_\alpha^2 z(\alpha, t)) \cdot \partial_\alpha^\perp z(\alpha, t) \end{aligned}$$

and therefore

$$\begin{aligned} \frac{\sigma(\alpha, t)}{\rho^2} &= (\partial_t BR(z, \varpi)(\alpha, t) + \frac{\varphi(\alpha, t)}{|\partial_\alpha z(\alpha, t)|} \partial_\alpha BR(z, \varpi)(\alpha, t)) \cdot \partial_\alpha^\perp z(\alpha, t) \\ &+ \frac{1}{2} \frac{\varpi(\alpha, t)}{|\partial_\alpha z(\alpha, t)|^2} (\partial_\alpha z_t(\alpha, t) + \frac{\varphi(\alpha, t)}{|\partial_\alpha z(\alpha, t)|} \partial_\alpha^2 z(\alpha, t)) \cdot \partial_\alpha^\perp z(\alpha, t) + g \partial_\alpha z_1(\alpha, t). \end{aligned} \quad (20)$$

4 Local existence

Our main results consist on the existence of a positive time τ (depending upon the initial conditions) for which we have a solution of the periodic Muskat problem (equations (6), (7) and (9)) and of the free boundary of the irrotational incompressible Euler equations in vacuum (equations (6), (7) and (14)) during the time interval $[0, \tau]$ so long as the initial data belong to $H^k(\mathbb{T})$ for k sufficiently large, $\mathcal{F}(z_0)(\alpha, \beta) < \infty$, and

$$\sigma_0(\alpha) = -(\nabla p^2(z_0(\alpha), 0) - \nabla p^1(z_0(\alpha), 0)) \cdot \partial_\alpha^\perp z_0(\alpha) > 0,$$

where p^j denote the pressure in Ω^j and the function $\mathcal{F}(z)$, which measures the arc-chord condition (see [16]), is defined by

$$\mathcal{F}(z)(\alpha, \beta, t) = \frac{|\beta|}{|z(\alpha, t) - z(\alpha - \beta, t)|} \quad \forall \alpha, \beta \in (-\pi, \pi), \quad (21)$$

with

$$\mathcal{F}(z)(\alpha, 0, t) = \frac{1}{|\partial_\alpha z(\alpha, t)|}.$$

Theorem 4.1 *Let $z_0(\alpha) \in H^k(\mathbb{T})$ for $k \geq 3$, $\mathcal{F}(z_0)(\alpha, \beta) < \infty$, and*

$$\sigma_0(\alpha) = -(\nabla p^2(z_0(\alpha), 0) - \nabla p^1(z_0(\alpha), 0)) \cdot \partial_\alpha^\perp z_0(\alpha) > 0.$$

Then there exists a time $\tau > 0$ so that there is a solution to (6), (7) and (9) in $C^1([0, \tau]; H^k(\mathbb{T}))$ with $z(\alpha, 0) = z_0(\alpha)$.

Theorem 4.2 *Let $z_0(\alpha) \in H^k(\mathbb{T})$, $\varpi_0(\alpha) \in H^{k-1}$ and $\varphi(\alpha, 0) = \varphi_0(\alpha) \in H^{k-\frac{1}{2}}$ defined in (18) for $k \geq 4$, $\mathcal{F}(z_0)(\alpha, \beta) < \infty$, $g \geq 0$, and*

$$\sigma_0(\alpha) = -(\nabla p^2(z_0(\alpha), 0) - \nabla p^1(z_0(\alpha), 0)) \cdot \partial_\alpha^\perp z_0(\alpha) > 0.$$

Then there exists a time $\tau > 0$ so that there is a solution to (6), (7) and (14) with $z(\alpha, t) \in C^1([0, \tau]; H^k(\mathbb{T}))$, $\varpi(\alpha, t) \in C^1([0, \tau]; H^{k-1}(\mathbb{T}))$ for $z(\alpha, 0) = z_0(\alpha)$ and $\varpi(\alpha, 0) = \varpi_0(\alpha)$.

Remark 4.3 *Notice that the parametrization is defined by properties (2) or (3). But in the Hele-Shaw and Muskat problems we can show easily that*

$$\int_{-\pi}^{\pi} \sigma(\alpha, t) d\alpha = 0$$

for a closed curve, making impossible the task of prescribing a sign as in the Rayleigh-Taylor condition.

5 Sketch of the proof

First we consider the operator $T(u)(\alpha) = 2BR(z, u)(\alpha) \cdot \partial_\alpha z(\alpha)$ associated to a smooth H^3 curve z satisfying the arc-chord condition. T is a smoothing compact operator in Sobolev space whose adjoint T^* , acting on u , is described in terms of the Cauchy integral of u along the curve z , and whose real eigenvalues have absolute value strictly less than one (see [3]).

In our proof it is crucial to get control of the norm of the inverse operators $(I - \xi T)^{-1}$, $|\xi| \leq 1$. The arguments rely upon the boundedness properties of the Hilbert transforms associated to $C^{1,\alpha}$ curves, for which we need precise estimates obtained with arguments involving conformal mappings, Hopf maximum principle and Harnack inequalities (see [6] and [13]).

We then provide upper bounds for the amplitude of the vorticity, the Birkhoff-Rott integral, the parametrization of the curve, the arc-chord condition and the Rayleigh-Taylor condition, namely:

5.1 A priori estimates for theorem 4.1

$$\|\varpi\|_{H^k} \leq \exp C(\|\mathcal{F}(z)\|_{L^\infty}^2 + \|z\|_{H^{k+1}}^2),$$

$$\|BR(z, \varpi)\|_{H^k} \leq \exp C(\|\mathcal{F}(z)\|_{L^\infty}^2 + \|z\|_{H^{k+1}}^2),$$

$$\begin{aligned} \frac{d}{dt} \|z\|_{H^k}^2(t) &\leq -\frac{\kappa}{2\pi(\mu_1 + \mu_2)} \int_{\mathbb{T}} \frac{\sigma(\alpha, t)}{|\partial_\alpha z(\alpha)|^2} \partial_\alpha^k z(\alpha, t) \cdot \Lambda(\partial_\alpha^k z)(\alpha, t) d\alpha \\ &\quad + \exp C(\|\mathcal{F}(z)\|_{L^\infty}^2(t) + \|z\|_{H^k}^2), \end{aligned}$$

and

$$\frac{d}{dt} \|\mathcal{F}(z)\|_{L^\infty}^2(t) \leq \exp C(\|\mathcal{F}(z)\|_{L^\infty}^2(t) + \|z\|_{H^3}^2(t))$$

where the operator Λ is defined by the Fourier transform $\widehat{\Lambda f}(\xi) = |\xi| \widehat{f}(\xi)$ and $\sigma(\alpha, t)$ is the difference of the gradients of the pressure in the normal direction. The first inequality in our list above is where we use the precise control of the norm of the inverse operator. The third estimate depends crucially upon the positive sign of the Rayleigh-Taylor condition and the chosen well adapted parametrization. The second and the fourth follow from the standard Sobolev embedding.

We then define the quantity $E(t)$ for the Muskat problem given by

$$E(t) = \|z\|_{H^k}^2(t) + \|\mathcal{F}(z)\|_{L^\infty}^2(t)$$

to get

$$\frac{d}{dt} E(t) \leq -\frac{\kappa}{2\pi(\mu_1 + \mu_2)} \int_{\mathbb{T}} \frac{\sigma(\alpha, t)}{|\partial_\alpha z(\alpha)|^2} \partial_\alpha^k z(\alpha, t) \cdot \Lambda(\partial_\alpha^k z)(\alpha, t) d\alpha + \exp C(E(t)). \quad (22)$$

5.2 A priori estimates for theorem 4.2

In the case of Bernoulli's law our previous comments can be applied in the following setting:
For some universal constant q

$$\|BR(z, \varpi)\|_{H^k} \leq C(\|\mathcal{F}(z)\|_{L^\infty}^2 + \|z\|_{H^{k+1}}^2 + \|\varpi\|_{H^k}^2)^q,$$

$$\|z_t\|_{H^k} \leq C(\|\mathcal{F}(z)\|_{L^\infty}^2 + \|z\|_{H^{k+1}}^2 + \|\varpi\|_{H^k}^2)^q,$$

$$\|\varpi_t\|_{H^k} \leq C(\exp C(\|\mathcal{F}(z)\|_{L^\infty}^2 + \|z\|_{H^3}^2))(\|\mathcal{F}(z)\|_{L^\infty}^2 + \|z\|_{H^{k+2}}^2 + \|\varpi\|_{H^{k+1}}^2 + \|\varphi\|_{H^{k+1}}^2)^q,$$

$$\|\varpi\|_{H^k} \leq C(\|\mathcal{F}(z)\|_{L^\infty}^2 + \|z\|_{H^{k+1}}^2 + \|\varpi\|_{H^{k-1}}^2 + \|\varphi\|_{H^k}^2)^q.$$

We define the quantity $E(t)$ in this case by

$$\begin{aligned} E(t) = & \|z\|_{H^{k-1}}^2(t) + \int_{\mathbb{T}} \frac{\sigma(\alpha, t)}{\rho^2 |\partial_\alpha z(\alpha, t)|^2} |\partial_\alpha^k z(\alpha, t)|^2 d\alpha \\ & + \|\mathcal{F}(z)\|_{L^\infty}^2(t) + \|\varpi\|_{H^{k-2}}^2(t) + \|\varphi\|_{H^{k-\frac{1}{2}}}^2(t). \end{aligned}$$

If we take

$$m(t) = \min_{\alpha \in \mathbb{T}} \sigma(\alpha, t)$$

and using that

$$\|\partial_\alpha^4 z\|_{L^2}^2(t) = \int_{\mathbb{T}} \frac{\sigma(\alpha, t)}{\sigma(\alpha, t)} |\partial_\alpha^4 z(\alpha, t)|^2 d\alpha \leq \frac{1}{m(t)} \int_{\mathbb{T}} \sigma(\alpha, t) |\partial_\alpha^4 z(\alpha, t)|^2 d\alpha,$$

we obtain

$$\frac{d}{dt} E(t) \leq \frac{1}{m^p(t)} C \exp(CE(t)), \quad (23)$$

with $p \in \mathbb{N}$.

5.3 The evolution of the Rayleigh-Taylor condition

It is clear that in the evolution of the quantities $E(t)$ in both contour dynamics problems (22) and (23), everything depends on the sign of $\sigma(\alpha, t)$. Therefore, in order to integrate the systems, it is crucial to study the evolution of

$$m(t) = \min_{\alpha \in \mathbb{T}} \sigma(\alpha, t).$$

Let us then introduce the Rayleigh-Taylor condition in a new definition of energy as follows:

$$E_{RT}(t) = E(t) + \frac{1}{m(t)}.$$

Similar arguments (see [8]) allows us to accomplish the fact that $m'(t) = \sigma_t(\alpha_t, t)$ for almost all t . Since we can use the formulas (15) and (20) and the above estimates, it is possible to control $\|\sigma_t(\alpha_t, t)\|_{L^\infty}$ by means of $E_{RT}(t)$. It yields finally

$$\frac{d}{dt}E_{RT}(t) \leq C \exp(CE_{RT}(t)),$$

for C a universal constant.

5.4 Existence

To conclude the existence proof we introduce regularized evolution equations (allowing us to take limits) satisfying uniformly the above a priori estimates and for which the local existence follows by standard arguments. Furthermore, in the case of the Hele-Shaw and Muskat problem, in order to take advantage of the positivity of σ we have to use a pointwise inequality satisfied by the non-local operator Λ (see [8]).

Acknowledgements

The first author was partially supported by the grant MTM2005-04730 of the MEC (Spain). The other two authors were partially supported by the grants MTM2008-03754 of the MEC (Spain) and ERC-2007-StG-203138-CDSIF.

References

- [1] Ambrose D (2004) Well-posedness of Two-phase Hele-Shaw Flow without Surface Tension. *European J Appl Math* 15:597-607.
- [2] Ambrose D, Masmoudi N (2005) The zero surface tension limit of two-dimensional water waves. *Comm Pure Appl Math* 58:1287-1315.
- [3] Baker G, Meiron D, Orszag S (1982) Generalized vortex methods for free-surface flow problems. *J Fluid Mech* 123:477-501.
- [4] Beale T, Hou T, Lowengrub J (1993) Growth rates for the linearized motion of fluid interfaces away from equilibrium. *Comm Pure Appl Math* 46:1269-1301.
- [5] Caffisch R, Orellana O (1989) Singular solutions and ill-posedness for the evolution of vortex sheets. *SIAM J Math Anal* 20:293-307.
- [6] Caffarelli L-A, Córdoba A (2006) Phase transitions: Uniform regularity of the intermediate layers. *J Reine Angew Math* 593:209-235.
- [7] Christodoulou D, Lindblad H (2000) On the motion of the free surface of a liquid. *Comm Pure Appl Math* 53:1536-1602.
- [8] Córdoba A, Córdoba D (2004) A maximum principle applied to Quasi-geostrophic equations. *Comm Math Phys* 249:511-528.

- [9] Córdoba A, Córdoba D, Gancedo F (2009) Interface evolution: the Hele-Shaw and Muskat problems. To appear in *Ann of Math*.
- [10] Córdoba A, Córdoba D, Gancedo F (2008) Interface evolution: the water wave problem in 2D. *ArXiv:0810.5340*.
- [11] Córdoba D, Gancedo F (2007) Contour dynamics of incompressible 3-D fluids in a porous medium with different densities. *Comm Math Phys* 273:445-471.
- [12] Coutand D, Shkoller S (2007) Well-posedness of the free-surface incompressible Euler equations with or without surface tension. *J Amer Math Soc* 20:829-930.
- [13] Dahlberg B-E-J (1979) On the Poisson integral for Lipschitz and C^1 -domains. *Studia Math* 66:13-24.
- [14] Ebin D-G (1987) The equations of motion of a perfect fluid with free boundary are not well posed. *Comm Partial Differential Equations* 12:1175-1201.
- [15] Ebin D-G (1988) Ill-posedness of the Rayleigh-Taylor and Helmholtz problems for incompressible fluids. *Comm Partial Differential Equations* 13:1265-1295.
- [16] Gancedo F (2008) Existence for the α -patch model and the QG sharp front in Sobolev spaces. *Adv Math* 217:2569-2598.
- [17] Hele-Shaw (1898) *Nature* 58 34.
- [18] Hou T, Lowengrub J, Shelley M (1994) Removing the Stiffness from Interfacial Flows with Surface Tension. *J Comput Phys* 114:312-338.
- [19] Lindblad H (2005) Well-posedness for the motion of an incompressible liquid with free surface boundary. *Ann of Math* 162:109-194.
- [20] Muskat M (1937) *The flow of homogeneous fluids through porous media*. New York.
- [21] Lord Rayleigh J-W-S (1879) On the instability of jets. *Proc Lond Math Soc* 10:4-13.
- [22] Saffman P-G, Taylor G (1958) The penetration of a fluid into a porous medium or Hele-Shaw cell containing a more viscous liquid. *Proc R Soc London Ser A* 245:312-329.
- [23] Shatah J, Zeng C (2008) Geometry and a priori estimates for free boundary problems of the Euler equation. *Comm Pure Appl Math* 61:698-744.
- [24] Siegel M, Caffisch R, Howison S (2004) Global Existence, Singular Solutions, and Ill-Posedness for the Muskat Problem. *Comm Pure and Appl Math* 57:1374-1411.
- [25] Stein E (1993) *Harmonic Analysis*. Princeton University Press.
- [26] Taylor G (1950) The instability of liquid surfaces when accelerated in a direction perpendicular to their planes. *Proc Roy Soc London Ser A* 201:192-196.

- [27] Wu S (1997) Well-posedness in Sobolev spaces of the full water wave problem in 2-D. Invent math 130:39-72.
- [28] Wu S (1999) Well-posedness in Sobolev spaces of the full water wave problem in 3-D. J Amer Math Soc 12:445-495.
- [29] Zhang P, Zhang Z (2008) On the free boundary problem of three-dimensional incompressible euler equations. Comm Pure and Appl Math 61:877-940.

Antonio Córdoba

Departamento de Matemáticas
Facultad de Ciencias
Universidad Autónoma de Madrid
Ctra. Colmenar Viejo km. 15, 28049 Madrid, Spain
Email: antonio.cordoba@uam.es

Diego Córdoba

Instituto de Ciencias Matemáticas
Consejo Superior de Investigaciones Científicas
Serrano 123, 28006 Madrid, Spain
Email: dcg@imaff.cfmac.csic.es

Francisco Gancedo

Department of Mathematics
University of Chicago
5734 University Avenue, Chicago, IL 60637
Email: fgancedo@math.uchicago.edu