




Contents lists available at ScienceDirect

Advances in Mathematics

[www.elsevier.com/locate/aim](http://www.elsevier.com/locate/aim)



# A trichotomy theorem for transformation groups of locally symmetric manifolds and topological rigidity



Sylvain Cappell\*, Alexander Lubotzky, Shmuel Weinberger

## ARTICLE INFO

### Article history:

Available online 19 January 2018

To our friend, David Kazhdan, with deep admiration and affection

### Keywords:

Manifolds  
Transformation groups  
Rigidity  
Locally symmetric spaces

## ABSTRACT

Let  $M$  be a locally symmetric irreducible closed manifold of dimension  $\geq 3$ . A result of Borel [6] combined with Mostow rigidity imply that there exists a finite group  $G = G(M)$  such that any finite subgroup of  $\text{Homeo}^+(M)$  is isomorphic to a subgroup of  $G$ . Borel [6] asked if there exist  $M$ 's with  $G(M)$  trivial and if the number of conjugacy classes of finite subgroups of  $\text{Homeo}^+(M)$  is finite. We answer both questions:

- (1) For every finite group  $G$  there exist  $M$ 's with  $G(M) = G$ , and
- (2) the number of maximal subgroups of  $\text{Homeo}^+(M)$  can be either one, countably many or continuum and we determine (at least for  $\dim M \neq 4$ ) when each case occurs.

Our detailed analysis of (2) also gives a complete characterization of the topological local rigidity and topological strong rigidity (for  $\dim M \neq 4$ ) of proper discontinuous actions of uniform lattices in semisimple Lie groups on the associated symmetric spaces.

© 2018 Elsevier Inc. All rights reserved.

\* Corresponding author.

*E-mail address:* [cappell@cims.nyu.edu](mailto:cappell@cims.nyu.edu) (S. Cappell).

## 1. Introduction

A positive dimensional, oriented, closed manifold  $M$  has a very large group of automorphisms (i.e., orientation preserving self-homeomorphisms). In fact this group  $\text{Homeo}^+(M)$  is infinite dimensional. But its finite subgroups are quite restricted. In 1969, Borel showed (in a classic paper [6] but which appeared only in 1983 in his collected works) that if  $M$  is a  $K(\pi, 1)$ -manifold with fundamental group  $\Gamma = \pi_1(M)$ , whose center is trivial, then every finite transformation group  $G$  in  $\text{Homeo}^+(M)$  is mapped injectively into the outer automorphism group  $\text{Out}(\Gamma)$  by the natural map (or more precisely into the subgroup  $\text{Out}^+(\Gamma)$  – see §2 – which has an index at most 2 in  $\text{Out}(\Gamma)$ ).

Let now  $M$  be a locally symmetric manifold of the form  $\Gamma \backslash H/K$  when  $H$  is a connected non-compact semisimple group with trivial center and with no compact factor and  $\Gamma$  a torsion free uniform irreducible lattice in  $H$ . In the situation in which strong rigidity (in the sense of Mostow [39]) holds (i.e., if  $H$  is not locally isomorphic to  $SL_2(\mathbb{R})$ ),  $\text{Out}(\Gamma)$  is a finite group,  $G \leq \text{Out}^+(\Gamma)$ ; in fact,  $\text{Out}^+(\Gamma) \subseteq N_H(\Gamma)/\Gamma$  and it acts on  $M$  as the group of (orientation preserving) self-isometries  $\text{Isom}^+(M)$  of the Riemannian manifold  $M$ . It follows now from Borel’s theorem that every finite subgroup of  $\text{Homeo}^+(M)$  is isomorphic to a subgroup of one finite group,  $G(M) = \text{Isom}^+(M)$ .

Borel ends his paper by remarking: “The author does not know whether the finite subgroups of  $\text{Homeo}^+(M)$  form finitely many conjugacy classes, nor whether one can find a  $\Gamma$  with no outer automorphism.”

The goal of the current paper is to answer these two questions. For an efficient formulation of our results, let us make the following definition(s):

**Definition 1.1.** Let  $G$  be a finite group. An oriented manifold  $M$  will be called  $G$ -dominated (resp.,  $G$ -weakly-dominated) if there is a faithful action of  $G$  on  $M$ , so that  $G$  can be identified with a subgroup of  $\text{Homeo}^+(M)$  and if  $F$  is any finite subgroup of  $\text{Homeo}^+(M)$ , then  $F$  is conjugate (resp., isomorphic) to a subgroup of  $G$ .

Note that Borel’s Theorem combined with strong rigidity implies that unless  $H$  is locally isomorphic to  $SL_2(\mathbb{R})$ ,  $M$  as above is always at least  $\text{Isom}^+(M)$ -weakly-dominated.

We now assert (to be proved in section 2)

**Theorem 1.2.** *For every finite group  $G$  and every  $3 \leq n \in \mathbb{N}$ , there exist infinitely many oriented closed hyperbolic manifolds  $M = M^n(G)$  of dimension  $n$ , with  $G \simeq \text{Isom}^+(M)$  and when  $n \neq 4$  these  $M^n(G)$  are also  $G$ -dominated.*

The very special case  $G = \{e\}$  answers Borel’s second question (where one can also deduce it from [2]). It also answers the question of Schultz [41,42], attributed there to D. Burghelca, who asked whether there exist asymmetric closed manifolds with degree one maps onto hyperbolic manifolds. Our examples are even hyperbolic themselves.

The situation for dimension 2 is very different:

**Theorem 1.3.** *For no group  $G$ , does there exist a  $G$ -weakly-dominated 2-dimensional closed manifold.*

In fact, for every closed, oriented surface  $\Sigma_g$ , of genus  $g$ ,  $\text{Homeo}^+(\Sigma)$  has more than one (but only finitely many) isomorphism classes of maximal finite subgroups, and this number is unbounded as a function of  $g$  – see [Proposition 6.2](#).

As mentioned before, for  $M$  as above with  $\dim M \geq 3$ ,  $M$  is always  $G$ -weakly-dominated. But the  $G$ -dominatedness shown in [Theorem 1.2](#) is not the general phenomenon. We can determine the situation in (almost) all cases. But first we need another definition.

**Definition 1.4.** For an automorphism  $\varphi$  of a manifold  $M$ , denote by  $\text{Fix}(\varphi)$  the fixed point set of  $\varphi$  and for a subgroup  $G \subseteq \text{Homeo}^+(M)$ , denote its *singular set*  $S(G) = \cup\{\text{Fix}(\varphi) \mid \varphi \in G, \varphi \neq \text{id}\}$ . If  $M$  is an oriented Riemannian manifold, then we will call  $S(\text{Isom}^+(M))$  the *singular set* of  $M$  and we denote it  $S_M$ .

We note that  $\dim(M) - \dim(S_M)$  is always even, as we are only considering orientation preserving actions.

Before stating our main theorem, let us recall that in our situation, i.e., when  $M$  is locally symmetric, every finite subgroup of  $\text{Homeo}^+(M)$  is contained in a maximal finite subgroup. We can now give a very detailed answer to Borel’s first question.

**Theorem 1.5** (*Trichotomy theorem*). *Let  $M = \Gamma \backslash H/K$  a locally symmetric manifold as above, and assume  $\dim M \neq 2$  or 4. Let  $G = \text{Isom}^+(M)$ , so  $G \cong N/\Gamma$  where  $N = N_H(\Gamma)$ . Then one of the following holds:*

- (a)  $\text{Homeo}^+(M)$  has a unique conjugacy class of maximal finite subgroups, all of whose members are conjugate to  $\text{Isom}^+(M)$ .
- (b)  $\text{Homeo}^+(M)$  has countably infinite many maximal finite subgroups, up to conjugacy or
- (c)  $\text{Homeo}^+(M)$  has a continuum of such subgroups (up to conjugacy).

*These cases happen, if and only if the following hold, respectively:*

- (a) (i)  $S_M = \emptyset$ , i.e.,  $\text{Isom}^+(M)$  acts freely on  $M$ , or  
(ii) the singular set  $S_M$  is 0-dimensional and either  $\dim(M)$  is divisible by 4 or all elements of order 2 act freely.
- (b)  $M$  is of dimension equal 2 (mod 4), the singular set is 0-dimensional and some element of order 2 has a non-empty fixed point set, or
- (c) the singular set  $S_M$  is positive dimensional, i.e.,  $M$  has some non-trivial isometry with a positive dimensional fixed point set.

The cases treated in [Theorem 1.2](#) are with  $N = N_H(\Gamma)$  torsion free (see §2), i.e.,  $S(G) = \emptyset$  where  $G = \text{Isom}^+(M)$ , so we are in case (a)(i) and these manifolds  $M$  are  $\text{Isom}^+(M)$ -dominated also by [Theorem 1.5](#).

An interesting corollary of the theorem is that if  $\text{Homeo}^+(M)$  has only finitely many conjugacy classes of maximal finite subgroups then it has a unique one, the class of  $\text{Isom}^+(M)$ , in contrast to [Theorem 1.3](#).

In dimension 4 when the action has positive dimensional singular set, we do construct uncountably many actions. If the singular set is finite, then we have countability, but we do not know whether/when this countable set of actions consists of a unique possibility. As a consequence, the following dichotomy holds in all dimensions:

**Corollary 1.6.** Let  $M$  be as above with arbitrary dimension. Then  $\text{Homeo}^+(M)$  has an uncountable number of conjugacy classes of finite subgroups if and only if the singular set of  $\text{Isom}^+(M)$  acting on  $M$  is positive dimensional.

The uniqueness in [Theorem 1.5](#) fails in the smooth case (i.e., for  $\text{Diff}^+(M)$ ). In that case, the number of conjugacy classes is always countable. (This is a straightforward consequence of the equivariant tubular neighborhood [8] theorem and countability of the number of compact manifolds in any dimension, which in turn follows, in the smooth case from handlebody theory.)

Finiteness without uniqueness is also possible. The simplest example occurs in dimension 7 where one can take any exotic differential structure on  $S^7$  and connect sum it with  $T^7$  28 times to get the standard  $T^7$ . This can easily be used to construct free exotic  $Z_{28}$  actions on  $T^7$ . (That the quotient manifold is not  $T^7$  is proved exactly the same way as the proof by Milnor that the original exotic sphere is not standard.)

The boundary between finite and infinite number of conjugacy classes of finite subgroups of  $\text{Diff}^+(M)$  can be largely analyzed by the methods of this paper, but works out somewhat differently (e.g., one has finiteness in some cases of one dimensional singular set) and is especially more involved when the singular set is 2-dimensional. We shall not discuss this here.

Finally, let us present our result from an additional point of view: Given  $H$  as above and  $\Gamma$  a uniform lattice in it. It acts via the standard action  $\rho_0$  by translation on the symmetric space  $H/K$  which topologically is  $\mathbb{R}^d$ . The Farrell–Jones topological rigidity result implies that if  $\Gamma$  is torsion free, every proper discontinuous (orientation preserving) action  $\rho$  of  $\Gamma$  on  $H/K$  is conjugate within  $\text{Homeo}^+(H/K)$  to  $\rho_0$ . It has been known for a long time (cf. [46] for discussion and references) that this is not necessarily the case if  $\Gamma$  has torsion. Our discussion above (with some additional ingredient based on [15,46] – see §7) gives the essentially complete picture. But first a definition:

**Definition.** For  $\Gamma, H, K$  and  $\rho_0$  as above, say

- (1) The lattice  $\Gamma$  has topological strong rigidity if every proper discontinuous action  $\rho$  of  $\Gamma$  on  $H/K$ , is conjugate to  $\rho_0$  by an element of  $\text{Homeo}^+(H/K)$ .
- (2)  $\Gamma$  has local topological rigidity if for every proper discontinuous action  $\rho$  of  $\Gamma$  on  $H/K$ , there exists a small neighborhood  $U$  of  $\rho$  in  $\text{Hom}(\Gamma, \text{Homeo}^+(H/K))$  such that any  $\rho' \in U$  is conjugate to  $\rho$  by an element of  $\text{Homeo}^+(H/K)$ .

The following two results follow from [Corollary 1.6](#) and [Theorem 1.5](#) (see §7):

**Theorem 1.7.** *Let  $H$  be a semisimple group,  $K$  a maximal compact subgroup and  $\Gamma$  an irreducible uniform lattice in  $H$ . Then  $\Gamma$  satisfies the topological local rigidity if and only if for every non-trivial element of  $\Gamma$  of finite order, the fixed point set of its action on  $H/K$  is zero dimensional.*

**Theorem 1.8.** *For  $H, K$  and  $\Gamma$  as in [Theorem 1.7](#) but assuming  $\dim(H/K) \neq 2, 4$ . Then one of the following holds:*

- (a)  $\Gamma$  has topological strong rigidity, i.e., it has a unique (up to conjugation) proper discontinuous action on  $H/K \simeq \mathbb{R}^n$ .
- (b)  $\Gamma$  has an infinite countable number of such actions, yet all are locally rigid.
- (c)  $\Gamma$  has uncountably many (conjugacy classes) of such actions.

*These cases happen if and only if the following hold, respectively:*

- (a) (i)  $\Gamma$  acts freely on  $H/K$  or  
(ii) every torsion (i.e., non-trivial of finite order) element of  $\Gamma$  has 0-dimensional fixed point set in  $H/K$  and either  $\dim(M) \equiv 0 \pmod{4}$  or there are no elements of order 2.
- (b)  $\dim(H/K) \equiv 2 \pmod{4}$ , the fixed point set of every torsion element is 0-dimensional and there is some element of order 2, or
- (c) there exist a torsion element in  $\Gamma$  with a positive dimensional fixed point set.

The paper is organized as follows. In §2, we prove [Theorem 1.2](#). In §3, we give preliminaries for the proof of [Theorem 1.5](#), which will be given in §4. In this proof we depend crucially on the deep works of Farrell and Jones [18,19] and Bartels and Lueck [2] related to the (famous) Borel conjecture as well as recent work of [15]. In §5, we analyze manifolds of dimension 4, while in §6 we prove [Theorem 1.3](#). Section 7 discusses topological rigidity of lattices and proves [Theorem 1.7](#) and [1.8](#).

**Remark.** If one allows orientation reversing actions then if  $\dim M \geq 7$  there is a trichotomy theorem; rigidity holds if the action is free or  $\dim M \equiv 1 \pmod{4}$  or if

$\dim M \equiv 3(4)$  and the elements of order 2 act freely. The proof of this is similar to the one we give below for [Theorem 1.5](#). We believe that the remaining cases, at least when  $\dim M \neq 4$ , work out similarly to that theorem.

**Acknowledgments:** This work was partially done while the authors visited Yale University. They are grateful to Dan Mostow for a useful conversation. We are also grateful to NYU, the Hebrew University, ETH-ITS for their hospitality as well as to the NSF, ERC, ISF, Dr. Max Rössler, the Walter Haefner Foundation and the ETH Foundation, for their support.

This paper is dedicated to David Kazhdan on his 70th birthday, recognizing the profound impact he has had on our and our community's mathematics, and wishing him many more years of leadership.

## 2. Proof of [Theorem 1.2](#)

The proof of the theorem is in four steps:

**Step I:** In [\[4\]](#), M. Belolipetsky and the second named author showed that for every  $n \geq 3$  and for every finite group  $G$ , there exist infinitely many closed, oriented, hyperbolic manifolds  $M = M^n(G)$  with  $\text{Isom}^+(M) \simeq G$ . More precisely, it is shown there that if  $\Gamma_0$  is the non-arithmetic cocompact lattice in  $H = PO^+(n, 1) = \text{Isom}^+(\mathbb{H}^n)$  constructed in [\[26\]](#), then it has infinitely many finite index subgroups  $\Gamma$  with  $N_H(\Gamma)/\Gamma \simeq G$ . The proof shows that  $\Gamma$  can be chosen so that  $N_H(\Gamma)$  is torsion free and moreover  $N_H(\Gamma)/\Gamma \simeq \text{Isom}^+(M) = \text{Isom}(M)$  for  $M = \Gamma \backslash \mathbb{H}^n$ . This implies that  $G = N_H(\Gamma)/\Gamma$  acts on  $M$  freely, (as a fixed point for an element of  $G$  could be lifted, by covering space theory, to an isometry of the universal cover, fixing a point, which would necessarily give us an element of finite order in  $N_H(\Gamma)$ ), a fact we will use in Step IV below.

Let  $M = M^n(G)$  be one of these manifolds,  $\Gamma = \pi_1(M)$ . So  $\Gamma$  can be considered as a cocompact lattice in  $\text{Isom}^+(\mathbb{H}^n) = PO^+(n, 1)$ , the group of orientation preserving isometries of the  $n$ -dimensional hyperbolic space  $\mathbb{H}^n$ .

**Step II:** The Mostow Strong Rigidity Theorem [\[39\]](#) for compact hyperbolic manifolds asserts that if  $\Gamma_1$  and  $\Gamma_2$  are torsion free cocompact lattices in  $\text{Isom}(\mathbb{H}^n)$ , then every group theoretical isomorphism from  $\Gamma_1$  to  $\Gamma_2$  is realized by a conjugation within  $\text{Isom}(\mathbb{H}^n)$  (or in a geometric language, homotopical equivalence of hyperbolic manifolds implies an isomorphism as Riemannian manifolds). Indeed, as  $\Gamma$  has no center, this conjugating element is unique.

Applying Mostow's theorem for the automorphisms of  $\Gamma = \pi_1(M)$  implies that  $\text{Aut}(\Gamma)$  can be identified with  $N_{\text{Isom}(\mathbb{H}^n)}(\Gamma)$ , the normalizer of  $\Gamma$  in  $\text{Isom}(\mathbb{H}^n)$ . Hence the outer automorphism group  $\text{Out}(\Gamma) = \text{Aut}(\Gamma)/\text{Inn}(\Gamma)$  of  $\Gamma$  is isomorphic to  $N_{\text{Isom}(\mathbb{H}^n)}(\Gamma)/\Gamma$  and hence also to  $\text{Isom}(M)$ , which in our case is equal to  $\text{Isom}^+(M)$  by step I.

**Step III:** In [\[6\]](#), Borel showed that if  $\Gamma$  is a torsion free cocompact lattice in any simple non-compact Lie group  $H$ , including our  $H = PO^+(n, 1)$ , with a maximal compact

subgroup  $K$  and associated symmetric space  $X = H/K$ , then every finite subgroup  $F$  of  $\text{Homeo}(M)$  where  $M = \Gamma \backslash X$ , is mapped injectively into  $\text{Out}(\pi_1(M)) = \text{Out}(\Gamma)$  by the natural map. If  $F \leq \text{Homeo}^+(M)$ , then its image is in  $\text{Out}^+(\Gamma)$  which is the kernel of the action of  $\text{Out}(\Gamma)$  on  $H^n(\Gamma, \mathbb{Z}) \simeq \mathbb{Z}$ , so  $[\text{Out}(\Gamma) : \text{Out}^+(\Gamma)] \leq 2$ . Borel's result is actually much more general; the reader is referred to that short paper for the general result and the proof which uses Smith theory and cohomological methods.

Anyway, applying Borel's result for our  $M = M^n(G)$  finishes the proof of the first part of [Theorem 1.2](#). In particular, one sees that in all these examples,  $F$  acts freely on  $M$  (and is abstractly a subgroup of  $G$ ) since the isometry group, in this case, is the group of covering transformations, which acts freely on  $M$ .

**Step IV:** We have shown so far that whenever a finite group  $F$  acts on  $M$  as above, there is a natural injective homomorphism  $F \hookrightarrow \text{Out}^+(\pi_1(M)) \cong \text{Isom}^+(M)$ . Denote the image of  $F$  in  $\text{Isom}^+(M)$  by  $L$ . Our next goal is to show that  $F$  is conjugate to  $L$  within  $\text{Homeo}^+(M)$ . For ease of reading we will call  $M$  with the action of  $F$ ,  $M'$ , to avoid confusion.

There is actually an equivariant map  $M' \rightarrow M$  that is a homotopy equivalence. To see this, recall that  $N_H(\Gamma)$  is torsion free and hence so is  $\bar{\Gamma}$ , the preimage of  $L$  in  $N_H(\Gamma)$  with respect to the natural projection  $N_H(\Gamma) \rightarrow \text{Out}(\Gamma) = \text{Out}(\pi_1(M))$ . Similarly, let us consider all of the possible lifts of all of the elements of  $F$  to the universal cover, which form a group  $\bar{\Gamma}'$  (the orbifold fundamental group of  $M'/F$ , which we presently show is the genuine fundamental group) that fits in an exact sequence:

$$1 \rightarrow \Gamma (= \pi_1(M)) \rightarrow \bar{\Gamma}' \rightarrow F \rightarrow 1$$

As  $\Gamma$  is centerless and  $F$  and  $L$  induce the same outer automorphism group, it follows that  $\bar{\Gamma}'$  is also torsion free and as a corollary  $F = \bar{\Gamma}'/\Gamma$  acts freely on  $M'$ . Hence  $M'/F$  is homotopy equivalent to  $M/L$  as both have  $\bar{\Gamma}' \simeq \bar{\Gamma}$  as their fundamental group. [Note that two groups containing the same centerless group as a normal subgroup, with an isomorphism between quotient groups that preserves the action on the normal subgroup are canonically isomorphic.] By the Borel conjecture for hyperbolic closed manifolds (which is a Theorem of Farrell and Jones [\[18\]](#) for  $n \geq 5$  and of Gabai–Meyerhoff–Thurston [\[23\]](#) for  $n = 3$ ) the map  $M'/F \rightarrow M/L$  is homotopic to a homeomorphism which preserves  $\pi_1(M') = \pi_1(M)$ , as did the original homotopy equivalence. Since liftability in a covering space is a homotopy condition, this homeomorphism can be lifted to the cover  $M' \rightarrow M$ , producing a conjugating homeomorphism between the actions. [Theorem 1.2](#) is now proved.

In summary, the above proof is analogous to (and relies on) the fact that Mostow rigidity gives a uniqueness of the isometric action (or in different terminology, the uniqueness of the Nielsen realization of a subgroup of  $\text{Out}(\Gamma)$ ). At the same time, the Farrell–Jones/Gabai–Meyerhoff–Thurston rigidity gives the uniqueness of the topological realization in the case of free actions. We will see later that this freeness condition is essential.

### 3. Some ingredients for the proof of Theorem 1.5

The proof of Theorem 1.5 is based on results, sometimes deep theorems, some of which are well-known and others which might be folklore (or new). We present them in this section and use them in the next one.

**Ingredient 3.1.** *For  $v \geq 3$ , there exist infinitely many non-simply connected homology spheres  $\Sigma^v$ , each bounding a contractible manifold  $X^{v+1}$  such that the different fundamental groups  $\pi_1(\Sigma)$  are all freely indecomposable and are nonisomorphic to each other. Moreover,  $X \times [0, 1]$  is a ball.*

**Proof.** For  $v > 4$  this is very straightforward. According to Kervaire [32], a group  $\pi$  is the fundamental group of a (PL) homology sphere iff it is finitely presented and superperfect (i.e.,  $H_1(\pi) = H_2(\pi) = 0$ ). It is quite simple to produce an infinite family of freely indecomposable groups that satisfy these conditions. Among finite groups, one can take the universal central extensions of an infinite family of simple groups. Moreover, as Kervaire shows in that paper, every PL homology sphere bounds a PL contractible manifold (this is true for  $v \geq 4$ , and for  $v = 3$  in the topological category [21]). The product of a contractible manifold with  $[0, 1]$  is a ball as an immediate application of the  $h$ -cobordism theorem (see [37]).

For  $v = 3$ , we could rely on the work of Mazur in the PL category, but would then need to use subsequent work on the structure of manifolds obtained by surgery on knots. Instead, as we will be working in the topological category, we rely on [21] which shows that the analogue of all of the above holds topologically for  $v = 3$ , aside from the characterization of fundamental groups: however, using the uniqueness of the JSJ ([28,29]) decomposition of Haken 3-manifolds, homology spheres obtained by gluing together nontrivial knot complements are trivially distinguished from one another. These fundamental groups are also freely indecomposable, because the constructed 3-manifolds are aspherical. (If they were decomposable, the manifolds would be connected sums, and the universal cover would have nontrivial  $\pi_2$ ).

For  $v = 4$ , note that all the fundamental groups of the  $v = 3$  case arise here as well: if  $\Sigma^3$  is a homology sphere then  $\partial(\Sigma^o \times D^2)$  is a homology 4-sphere with the same fundamental group (where  $\Sigma^o$  denotes, as usual, the punctured manifold).  $\square$

We also need:

**Ingredient 3.2.** *For  $m - 1 > c_0 \geq 3$  and every orientation preserving linear free action  $\rho$  of  $G = \mathbb{Z}_p$  on  $S^{c_0}$  (in particular,  $c_0$  is odd), there exist an infinite number of homology spheres  $\Sigma^{c_0}$  with non-isomorphic fundamental groups and with a  $G = \mathbb{Z}_p$ -free action satisfying: For each such  $\Sigma$  there exists an action of  $G$  on  $B^m$  fixing  $0 \in B^m$  such that*

- (1) *The action of  $G$  on  $S^{m-1}$  is isomorphic to the linear action  $\rho \oplus \text{Identity}$ , and*



- (2) the local fundamental group  $\pi_1^{local}(B^m \setminus F, 0)$  is isomorphic to  $\pi_1(\Sigma)$ , when  $F$  is the fixed point set. Moreover, this action is topologically conjugate to a PL action on a polyhedron.

Let us recall what is meant by the local fundamental group: This is the inverse limit  $\varprojlim_{\alpha} \pi_1(U_{\alpha}, x_{\alpha})$  where the  $\{U_{\alpha}\}$  is a sequence of connected open neighborhoods converging down to 0, and  $x_{\alpha} \in U_{\alpha} \setminus F$  is a sequence of base points. Note that by the Jordan Curve Theorem,  $U_{\alpha} \setminus F$  is connected as  $\text{codim } F \geq 2$ . Also the induced maps are well defined up to conjugacy, so the limit is well defined.

**Proof.** For every homology sphere  $\Sigma'$  of odd dimension  $c_0$ , let  $\Sigma = p\Sigma' = \Sigma' \# \Sigma' \# \cdots \# \Sigma'$   $p$  times. We now give  $\Sigma$  a free  $\mathbb{Z}_p$  action, by taking connected sum along an orbit of the free linear action on  $S^{c_0}$  with the permutation action on  $p\Sigma'$ . The action on  $S^{c_0}$  bounds a linear disk  $D^{c_0+1}$ . One can take the (equivariant) boundary connect sum of this disk with  $pX$ ,  $X$  the contractible manifold that  $\Sigma'$  bounds to get a contractible manifold  $Z$  with  $\mathbb{Z}_p$  action fixing just one point which is locally smooth at that point and has the given local representation  $\rho$  there. (Let  $Z$  be the locally linear contractible  $\mathbb{Z}_p$ -manifold we just constructed.)

For motivation, consider now  $c(\Sigma) \times B^{m-c_0-1}$  where  $c(\Sigma)$  is the cone of  $\Sigma$ . It is a ball, by Edwards's theorem [11], [34] and [16, p. 3]. (The contractible manifold  $\Sigma$  bounds maps to  $c\Sigma$  in a cell like way, and its product with a ball is a ball by the  $h$ -cobordism theorem), and has an obvious  $\mathbb{Z}_p$  action as desired except that the action on the boundary is not linear: the fixed set is  $S^{m-c_0-2}$  but it is not locally flat.

We will now make use of  $Z$  to solve this problem. The manifold  $Z \cup (\Sigma \times [0, 1]) \cup Z$  is a sphere (by the Poincaré conjecture). If one maps this to  $[0, 1]$  by the projection on  $\Sigma \times [0, 1]$  and extending by constant maps on the two copies of  $Z$ , then the mapping cylinder of this map  $\varphi : Z \cup (\Sigma \times [0, 1]) \cup Z \rightarrow [0, 1]$  is a manifold, again by Edwards's theorem. It has an obvious  $\mathbb{Z}_p$  action with fixed set an interval. The action on the boundary sphere is locally smooth with two fixed points, so that an old argument of Stallings [44] shows that it is topologically linear with  $\rho$  as above, see [27] for the details. Note that the nonlocally flat points of the fixed point set correspond to the points where the local structure is  $c(\Sigma) \times [0, 1]$ ; hence the local fundamental group is  $\pi_1(\Sigma)$ , as required. This proves the result for the case  $m = c_0 + 2$ .

For  $m > c_0 + 2$ , one can spin this picture:  $\text{Map}(S^{m-c_0-1} \times Z^{c_0+1}) \cup (B^{m-c_0-1} \times \Sigma)$  to  $B^{m-c_0}$  in the obvious way and again the mapping cylinder produces a ball with locally linear boundaries and desired fixed set. This time the linearity of the boundary action follows from the deeper results of Illman [27].  $\square$

We will also need the following group theoretical result:

**Proposition 3.3.** *If  $\{\pi_i\}_{i=1}^\infty$  and  $\{\pi'_i\}_{i=1}^\infty$  are two infinite countable families of non-isomorphic freely indecomposable groups such that  $\prod_{i=1}^\infty \pi_i$  is isomorphic to  $\prod_{i=1}^\infty \pi'_i$ , then after reordering for every  $i$ ,  $\pi_i$  is isomorphic to  $\pi'_i$ .*

**Proof.** Recall that by the Bass–Serre theory, a group  $\Gamma$  is a free product  $\prod_{i=1}^\infty \pi_i$  if and only if  $\Gamma$  acts on a tree  $T$  with trivial edge stabilizers and a contractible quotient and with one to one correspondence between the vertices of  $T$  and the conjugates of  $\pi_i (i \in \mathbb{N})$  in  $\Gamma$ , where each vertex corresponds to its stabilizer. Now assume  $\Gamma \simeq \prod_{i=1}^\infty \pi_i$  and also  $\Gamma \simeq \prod_{j=1}^\infty \pi'_j$  with the corresponding trees  $T$  and  $T'$ . Fix  $i \in \mathbb{N}$ , as  $\Gamma$  acts on  $T'$  with trivial edge stabilizers and  $\pi_i$  is freely indecomposable,  $\pi_i$  fixes a vertex of  $T'$ . Hence there exists  $j \in \mathbb{N}$  s.t.  $\pi_i \subseteq \pi'_j{}^\tau$ . In the same way  $\pi'_j{}^\tau$  is a subgroup of some  $\pi_k^\delta$  for some  $\delta \in \Gamma$ . This means that  $\pi_i \subseteq \pi_k^\delta$ . But in a free product a free factor cannot have a non-trivial intersection with another factor or with a conjugate of it. Moreover, if  $\pi_i \cap \pi_i^\delta \neq \{e\}$ , then  $\pi_i^\delta = \pi_i$ . Indeed, if  $g$  is in this intersection, it fixes the fixed vertex of  $\pi_i$  as well as that of  $\pi_i^\delta$ , hence also the geodesic between them, in contradiction to the fact that  $\Gamma$  acts with trivial edge stabilizers.

We deduce that  $\pi_i \subseteq \pi'_j{}^\tau \subseteq \pi_i$  and hence  $\pi_i = \pi'_j{}^\tau$ .

This shows by symmetry that the collections  $\{\pi_i\}$  and  $\{\pi'_j\}$  are identical.  $\square$

Finally, let us recall the famous Borel conjecture which asserts that two aspherical manifolds which are homotopy equivalent are homeomorphic. Moreover, the original homotopy equivalence is homotopic to a homeomorphism. This conjecture of Borel was proved by Farrell and Jones [19] for the locally symmetric manifolds  $M$  discussed in this paper, if  $\dim(M) \geq 5$  and by Gabai–Meyerhoff–Thurston [23] for the case of  $\dim(M) = 3$ . This, in particular, says that there is a unique cocompact proper topological action of  $\Gamma = \pi_1(M)$  on the symmetric space  $H/K$  for any uniform torsion-free lattice.

But if  $\bar{\Gamma} \triangleright \Gamma$  is a finite extension with torsion, then the situation is more delicate. In fact, as we will see, there is no rigidity anymore in the topological setting and  $\bar{\Gamma}$  may have many inequivalent actions on  $H/K$ . In other words, the “equivariant Borel conjecture” is not true. It is of interest (though not really relevant to the goals of this paper) to compare this with the analogous situation in the setting of  $C^*$ -algebras, where the analogue of the Borel conjecture is the Baum–Connes conjecture. This latter conjecture is known to be true in many situations, even in its equivariant form, i.e., for groups with torsion, while the equivariant Borel conjecture fails in some of those cases ([14,40,45,46]). The failure is due to the non-vanishing of the Nil and the first author’s UNil groups (see [3] and [12]).<sup>1</sup> The latter is the source of non-rigidity in the case of isolated singularities.

The specific outcome relevant for our needs is the following:

<sup>1</sup> More accurately, these algebraic reasons explain the failures of the equivariant Borel conjecture relevant here. [46] gives other sorts of examples when the singular set is high dimensional.

**Ingredient 3.4.** *If  $\Delta$  is a lattice containing  $\Gamma$  a torsion free uniform lattice in  $H$  as a normal subgroup of finite index, so that the normalizer of any finite subgroup of  $\Delta$  in  $\Delta$  is finite, then the proper discontinuous actions of  $\Delta$  on (the topological manifold)  $H/K$  are in a one to one correspondence with the action of  $\Delta/\Gamma$  on  $\Gamma \backslash H/K$ , inducing the given outer automorphism of  $\Gamma$ . If  $\dim(H/K) \neq 4$  then this action is unique unless  $\dim(H/K)$  is  $2 \pmod 4$  and  $\Delta$  contains an infinite dihedral subgroup. In that case, the number of conjugacy classes is infinite and countable.*

**Proof.** Any proper action of  $\Delta$  is automatically free when restricted to  $\Gamma$  (by torsion freeness). The action is cocompact, because if it were not, this quotient space would show that the cohomological dimension of  $\Gamma$  is less than  $\dim(H/K)$  which cannot happen, since there is a cocompact action. As a result, the Borel conjecture, proved for uniform lattices by [19], shows that all of these actions are standard for the  $\Gamma$  subgroup, i.e., equivalent to the original action of  $\Gamma$  on  $H/K$ .

Note that this argument did not use the fact that the manifold on which  $\Delta$  acts is  $H/K$ ; it would apply automatically to any contractible manifold. This shows that such a manifold is automatically Euclidean space, as follows quite directly from [43].

With this preparation, the result now follows from [15] together with [2]: The condition on normalizers is equivalent to the discreteness of the fixed point set for the isometric action. The exotic action now has the same property by Smith theory (see e.g. [7]), as one can see that each element of the isometry group must have discrete fixed set from the fact that all the elements of order  $p$  do (which is the Smith theoretic statement, noting that because for each finite group  $G$ ,  $N_H(G)/G$  acts properly on the fixed set of  $G$  on  $H/K$ ,  $\pmod p$  acyclicity of a component of fixed set implies that component is a point.) Assuming the Farrell–Jones conjecture for  $\Delta$ , which is a theorem of [2,15] gives a description of the set of actions in terms of UNil groups and maximal infinite dihedral subgroups  $\Delta$ .<sup>2</sup> A lattice that contains an infinite dihedral subgroup, contains a maximal one (by discreteness: the  $\mathbf{Z}$  subgroups cannot keep growing in a nested sequence, since they correspond to shorter and shorter closed geodesics and a compact manifold has a positive injectivity radius).  $\square$

#### 4. Proof of Theorem 1.5

For Theorem 1.2, we have depended on the fact that the construction of [4] produced free actions. The constructions we presently describe show that whenever a manifold  $M$  of dimension  $\geq 5$  has an (orientation preserving) action whose singular set (i.e., the union

<sup>2</sup> Essentially the argument shows that, unlike what is done in the next section in the situation where the singular set is positive dimensional, the action of  $\Delta/\Gamma$  is equivariantly homotopy equivalent to the linear one. Since the singular set is very low dimensional, one can promote this to an isovariant homotopy equivalence. At that point, surgery theory can be used to reduce this problem to issues in  $K$ -theory and  $L$ -theory that are handled by the Farrell–Jones conjecture. It turns out that the  $K$ -groups of  $\Delta$  are the limit of those of the finite subgroups of  $\Delta$ ; however, because of UNil, the analogous statement is not true for  $L(\Delta)$  and the calculation, *in this case*, reduces to the infinite dihedral subgroups.

of the fixed sets of all nontrivial subgroups) is positive dimensional, there are actually continuously many actions on  $M$  that induce the same outer automorphisms of their fundamental group but are not topologically conjugate.

To prove part (c) of the Theorem it suffices to prove it for cyclic group, i.e., that if a cyclic group  $C = \mathbb{Z}_p$ ,  $p$  prime, has a positive dimensional fixed point set, then there is a continuum of such non-equivalent actions. The case of general  $G$  then easily follows.

Let  $V$  be a component of the fixed set of the action of  $C$ . Let  $v = \dim(V)$  and let  $\rho$  be the normal representation of  $\mathbb{Z}_p$  on  $\mathbb{R}^c$  ( $c = m - v$ , the codimension of  $V$  in  $M$ ). If  $v > 2$ , take  $\Sigma^v$  and  $X^{v+1}$  as in Ingredient 3.1. The product  $X \times D(\rho)$  is a ball with an action of  $\mathbb{Z}_p$  whose fixed set is  $X$ . The action of  $C$  on the boundary  $\partial(X \times D^c(\rho)) = (X \times S^{c-1}(\rho)) \cup_{(\Sigma \times S^{c-1}(\rho))} (\Sigma \times D(\rho))$  has  $\Sigma$  as its fixed set (as  $C$  has no fixed points on  $S^{c-1}(\rho)$  and a unique fixed point – the origin – in the disk  $D(\rho)$ ). The normal representation to this fixed set is still  $\rho$ . We can take equivariant connected sum of  $M$  with this  $C$ -sphere to obtain a new  $C$ -action on  $M \# S^{c+v} \simeq M$  whose fixed set is  $V \# \Sigma$ . Since the fixed set of the new action does not have the same fundamental group as  $V$  (e.g., by Grushko’s theorem as  $\pi_1(\Sigma)$  was assumed nontrivial), it is not conjugate to the original action. Of course it induces the same outer automorphism on  $\pi_1$ . Notice that we can think of this procedure as being a local equivariant insertion; near a point  $x \in V$  we modify the action of  $C$  only in a small specified ball. This procedure can be done any finite number of times to get countably many non-conjugate actions.

In fact, we can even get a continuum of actions of  $C$  on  $M$ . Let us first make a definition: For a finite group  $G$  acting topologically on a manifold  $M$ , we say that  $x$  in  $M$  is a *decent fixed point*, if the action of  $G$  in some open neighborhood of  $x$  is topologically conjugate to a simplicial action on a polyhedron. Now, apply the process above with smaller and smaller disjoint balls in  $M$  converging to some point  $x_0$  using any choice of  $\pi_1(\Sigma)$ ’s provided by Ingredient 3.1. The outcome is a copy of  $M$  with an action of  $C$  on it with a fixed point set  $W$  containing  $x_0$ , which is the unique non-decent point on  $M$ . This set  $W$  is not a manifold, but  $W \setminus \{x_0\}$  is. Moreover, the fundamental group of  $W \setminus \{x_0\}$  is isomorphic to the free product of the fundamental group of the original set  $V$  with the free product of the infinitely many different  $\pi_1(\Sigma)$ ’s which have been used. Now, if two such constructions lead to equivalent actions of  $C$  on  $M$ , then this unique non-decent point of the fixed point set should be preserved. Proposition 3.3 would imply that the two collections of  $\pi_1(\Sigma)$ ’s are equal. An infinite countable set has a continuum number of subsets and we can therefore get a continuum number of non-conjugate actions of  $C$ . This proves case (c) for  $v \geq 3$ .

In the case of  $v \geq 3$ , we replaced balls  $B^m$  (with isometric local  $C$ -action) of  $M$ , by copies of the same ball with new  $C$ -actions. The new action preserved the original normal action  $\rho$  but changed dramatically the fixed point set. For  $v = 1$  or  $2$ , we are not able to argue like that (for lack of suitable  $\Sigma$ ’s as above). Instead we will keep the fixed point set in  $B^m$  but deform the normal action  $\rho$ . In fact, this second method works whenever  $c = m - v > 2$ , so altogether the two methods cover all cases if  $\dim M \geq 5$ .

One now imitates the procedure described before to modify the original isometric action of  $C$  on  $M$  at a ball around a fixed point by replacing it with some  $B^m$  as in Ingredient 3.2. The resulting action is not equivalent to the original one as  $\pi_1(\Sigma)$  can be recovered from it as the local fundamental group at a fixed point. Doing this procedure any finite number of times with different  $\Sigma$ 's each time, gives us an infinite countable collection of nonconjugate actions. To get a continuum, we argue as before. In fact, this version is easier: The family of  $\pi_1(\Sigma)$ 's used can be recovered from the action of  $G$  on  $M$  as being exactly the collection of non-trivial local fundamental groups at decent fixed points of  $G$ .

This finishes the proof of part c of Theorem 1.5.

Now, the proof of part (a) is exactly the same as Step IV in the proof of Theorem 1.2.

Part (b) can be deduced from Ingredient 3.4. The existence of a dihedral subgroup in  $\bar{\Gamma}$  is equivalent to the existence of an involution in  $G$  fixing a point in  $M$ . This is indeed the case: If  $\bar{\Gamma}$  contains a dihedral group, it contains an element of order 2. This element has a fixed point on  $H/K$  and hence also on  $M$ . In the other direction: assume  $\tau \in G$  is an involution fixing a point  $p$  of  $M$ . Then by [13],  $\tau$  has at least a second fixed point  $q$ . Let  $\alpha$  be a geodesic from  $p$  to  $q$ , then  $\tau(\alpha)$  is another such geodesic and indeed,  $\alpha \cup \tau(\alpha)$  is a closed geodesic  $\gamma \in \pi_1(M, p) = \Gamma$ . The group generated by  $\gamma$  and  $\tau$  is a dihedral group. When we have a dihedral subgroup and the dimension is  $2 \bmod 4$ , ingredient 3.4 gives infinitely many conjugacy classes of  $\Delta$  actions. By the Farrell–Jones theorem, these action are all conjugate as  $\Gamma$  actions, so the actions of  $\Delta/\Gamma$  are nonconjugate actions on  $M = \Gamma \backslash G/K$ .

This finishes the proof of Theorem 1.5 for  $\dim M \geq 5$ .

To prove Theorem 1.5 for dimension 3 observe that case (b) does not occur; since the fixed point set is of even codimension. Now, case (a) is exactly as before, with this time the work of Gabai–Meyerhoff–Thurston [23] replacing the work of Farrell and Jones.

For part (c), we use the same procedure of replacing a ball around one fixed point by an exotic action. This time the work of Bing [5], provides us with uncountably many actions of  $G = \mathbb{Z}_p$  on  $B^3$ . (Bing discusses  $\mathbb{R}^3$ , but his construction clearly works on  $B^3$  and produces actions with given linear  $G$  action on  $\partial B^3$ .) As the actions are distinguished by the structures of this non-locally flat set as a subset of the line, they remain inequivalent in any manifold.  $\square$

## 5. The case of $\dim M = 4$

Theorem 1.5 may hold true in dimension 4 as well, but we can only prove part (c) of it, namely:

**Theorem 5.1.** *Let  $M$  be a locally symmetric irreducible manifold of dimension 4, with a  $G = \mathbb{Z}_p$  faithful isometric action whose fixed point set  $V$  is positive dimensional. Then  $G$  has a continuum number of inequivalent topological actions on  $M$ .*

**Proof.** First note that  $\dim V = 2$  as the codimension must be even. The Smith conjecture asserts that if  $\mathbb{Z}_p$  acts topologically on  $S^n$ ,  $n \geq 3$  with a smooth  $S^{n-2}$  as its fixed point set then  $S^{n-2}$  is isotopic to the unknot. While this conjecture is true for  $n = 3$ , it turns out to be false when  $n \geq 4$  as we now discuss.

**Theorem 5.2.** *There exists an infinite number of  $G = \mathbb{Z}_p$  actions  $\rho_i$ ,  $i \in \mathbb{N}$  on  $S^4$  satisfying:*

- (1) *The fixed point set is a nontrivial knotted  $S^2 \subset S^4$ .*
- (2) *The complement of the knot is a 4-dimensional manifold  $N_i$ , which fibers over the circle, with fibres  $F_i^0$ , each a 3-dimensional manifold with boundary  $S^2$ . Once this  $S^2$  is filled by a ball, the resulting closed manifold  $F_i$  is irreducible.*
- (3) *For every  $i \neq j$ ,  $\pi_1(F_i)$  is not isomorphic to a subgroup of  $\pi_1(F_j)$ .*

Note that  $\pi_1(F_i) = \pi_1(F_i^0)$  and this is a normal subgroup of  $\pi_1(N_i)$ , equal to  $[\pi_1(N_i), \pi_1(N_i)]$  with the quotient isomorphic to  $\mathbb{Z}$ . Also, as  $F_i$  is irreducible,  $\pi_1(F_i)$  is freely indecomposable. We postponed the proof of [Theorem 5.2](#), using it first to prove [5.1](#).

Given  $x \in V \subset M$ , we will replace a punctured sphere  $S^4$  around  $x \in M$  (with its  $G$ -action) by a punctured  $S^4$  around a fixed point provided by [Theorem 5.2](#). (We can adjust the action on the sphere to have the same normal representation as that of  $V$  just by changing the generator of  $G = \mathbb{Z}_p$ .) More precisely, as in the proof of [Theorem 1.5](#), we replace  $M$  by  $M \# S^4$ , but the latter is homeomorphic to  $M$ . The new fixed point set is  $V \# S^2$ .

This time, neither the fundamental group of the fixed set nor the local fundamental group distinguishes the two actions. Thus we will argue slightly differently: Look at the universal cover  $\tilde{M}$  of  $M$ . Take  $\tilde{V}$  to be one of the components of the lift of  $V$  in  $\tilde{X}$  containing a lift of  $x$  (actually  $\tilde{V}$  is the universal cover of  $V$  since  $\pi_1(V)$  injects into  $\pi_1(M)$ ). The complement of it has fundamental group isomorphic to  $\mathbb{Z}$  (as by Hadamard's Theorem this is diffeomorphic to the linear inclusion  $\tilde{V} \simeq \mathbb{R}^2 \subset \mathbb{R}^4 \simeq \tilde{M}$ , whose complement is homotopy equivalent to a circle). On the other hand, if  $\tilde{W}$  is a lift of  $W$  (= the fixed points of the modified action) then the fundamental group of its complement is an infinite amalgamated free product of  $\pi_1(N_i)$  with itself amalgamated along  $\mathbb{Z}$  (generated by the meridian). This group is certainly not isomorphic to  $\mathbb{Z}$  and hence the two actions are not equivalent. Repeating this with the different  $N_i$ 's gives us countably many inequivalent actions, as the different  $\pi_1(N_i)$ 's are not isomorphic to each other. (These groups have  $\pi_1 F_i * \pi_1 F_i * \dots$  as their commutator subgroups.)

To get a continuum number, we argue as follows: Let  $\Omega$  be an infinite subset of  $\mathbb{N}$ . For every fixed  $i \in \Omega$ , apply the procedure above infinitely many times around disjoint balls in  $M$  converging to a point  $x_i \in M$ . Do it in such a way that the set  $\{x_i\}_{i \in \Omega}$  has a unique limit point  $x_0$ .

We got therefore a new action on  $G$  on  $M$  depending on  $\Omega$ . We want to show that different  $\Omega$ 's lead to non-equivalent actions. Indeed, the indecent points are  $\{x_i\}_{i \in \Omega}$  as well as  $x_0$  (which is a unique limit point of the indecent points). Now if two families  $\Omega$  and  $\Omega'$  lead to equivalent actions then the conjugating homeomorphism takes, after reordering,  $x_i, i \in \Omega$  to  $x'_{j'}, j' \in \Omega'$ . Looking, as before, at the neighborhood of a lift of  $x_i$  (and  $x_{j'}$ ), in the universal cover, we deduce that  $\pi_1(F_{j'})$  is a subgroup of the infinite amalgamated product of infinitely many copies of  $\pi_1(N_i)$ . In fact it is in the (unique) kernel of the map from this group onto  $\mathbb{Z}$ . Thus  $\pi_1(F_{j'})$  is a subgroup of this kernel which is just a free product of infinitely many copies of  $\pi_1(F_i)$ . By the Kurosh Subgroup Theorem every subgroup of this kernel is a free product of subgroups of  $\pi_1(F_i)$  and of a free group. As  $\pi_1(F_{j'})$  is freely indecomposable (since  $F_{j'}$  is irreducible) we deduce that  $\pi_1(F_{j'})$  is isomorphic to a subgroup of  $\pi_1(F_i)$ . By part (3) of [Theorem 5.2](#) this implies  $i = j'$  and hence  $\Omega = \Omega'$  and [Theorem 5.1](#) is now proven.  $\square$

Let us now prove [Theorem 5.2](#).

**Proof.** There are many ways of doing this construction. We follow the work of [\[24\]](#) and [\[47\]](#) and use work of [\[38\]](#) to choose an explicit set of examples. In [\[47\]](#), Zeeman modified Artin's spinning construction of knots in  $S^4$  to twist spinning. Artin took a knot and imagined a child holding it in two hands and spinning it through the 4-th dimension, so that the trajectory of this rotated interval (the part beyond the hands) forms an  $S^2$  in  $S^4$ . He observed that the new knot complement has the fundamental group as the original knot complement. Zeeman [\[47\]](#) suggested having the knot rotate in the normal direction  $q$  times as it makes a rotation. He then saw that this knot in  $S^4$  has a Seifert surface which is the  $q$ -fold branched cyclic cover of  $S^3$  over the original knot, and that the natural generator of this covering is the monodromy of the action. If  $p$  is prime to  $q$ , then monodromy has a  $q$ -th root. Using the 4-dimensional Poincaré conjecture [\[21\]](#), Giffen observes<sup>3</sup> that this action can be extended to one on the sphere with the twist spun knot as fixed set.

If one starts with  $(r, s)$  torus knots, one obtains the Brieskorn manifold associated to  $(q, r, s)$  as the closed fibers. These groups, as observed in [\[38\]](#) are central extensions of the  $(q, r, s)$ -Delta groups. Their quotients by their centers have as torsion exactly the cyclic groups of order  $\{q, r, s\}$ ; then by, for example, letting  $q, r, s$  run through primes these groups do not embed in one another.  $\square$

## 6. The case of $\dim M = 2$

The phenomena in dimension 2 genuinely differ from those in higher dimension. Let us note that Step I and Step III in the proof of [Theorem 1.2](#) work equally well in

<sup>3</sup> Giffen's paper also shows that for  $p$  odd, one can avoid using the Poincaré conjecture (which was not known in any category at the time that paper was written).

dimension 2. In Borel’s result there is no assumption on the dimension. The paper [4] assumes  $n \geq 3$ , but the result is true also for  $n = 2$ . In fact, it was proved earlier by Greenberg [25] with the following elegant argument: He showed that in the Teichmüller space classifying the hyperbolic structures on a given surface  $S_g$  of genus  $g$ , or better yet, classifying conjugacy classes of cocompact lattices of  $PSL_2(\mathbb{R})$  isomorphic to  $\pi_1(S_g)$ , almost every such lattice is maximal and non-arithmetic. Hence by Margulis’ criterion for arithmeticity, it is equal to its commensurability group. Now given a finite group  $G$ , choose  $g$  large enough so that  $\pi_1(S_g)$  is mapped onto  $G$  (this is possible since  $\pi_1(S_g)$  is mapped onto  $F_g$ , the free group on  $g$  generators, so taking  $g$  greater or equal to the number of generators of  $G$  will do). Now, let  $\Gamma$  be a non-arithmetic maximal lattice with epimorphism  $\pi : \Gamma \twoheadrightarrow G$  with kernel  $\Lambda$ . Then  $N_{PSL_2(\mathbb{R})}(\Lambda) = \Gamma$  and so  $\text{Isom}^+(\Sigma) = G$  for  $\Sigma = \Lambda \backslash \mathbb{H}^2$ , as needed.

Moreover, it is also true that given a surface  $S$  of genus  $\sigma \geq 2$ , there are only finitely many conjugacy classes of finite subgroups in  $\text{Homeo}^+(S)$ . (These are, by the Nielsen realization theorem [31], in one to one correspondence with the conjugacy classes of finite subgroups of the mapping class group  $MCG(S)$ .) The analysis below shows that this number is more than 1 for every genus, in contrast to parts (a) and (b) of Theorem 1.5, in spite of the fact that the singular set is always 0-dimensional. To see the finiteness note first that for any finite group  $G$  there are only finitely many conjugacy classes of subgroups of  $G$ . That all topological actions of finite groups on surfaces can be smoothed is classical [30]. Smooth actions can be made isometric on some hyperbolic structure either by direct construction (cut paste PL methods) or by using the uniformization theorem: there is a unique hyperbolic structure conformal to any invariant Riemannian metric, and that hyperbolic metric has an isometric action of  $G$ .

The finiteness of the number of actions is either obvious by thinking of the data required to reconstruct  $\Sigma \rightarrow \Sigma/G$  in terms of the quotient manifold, ramification points, and group homomorphisms from  $\pi_1$  (Nonsingular part of  $\Sigma/G$ )  $\rightarrow G$  or by using the uniformization theorem to see that the actions then correspond to the strata of moduli space – and there are of course only finitely many strata in any variety. (See our discussion of the Riemann–Hurwitz formula below.) As one varies over all finite groups, one has only a finite amount of data for any fixed genus.

Despite all this, let us show that Theorem 1.2 fails in dimension 2 in the strongest possible way, namely:

**Theorem 6.1.** *For no finite group  $G$  does there exist a  $G$ -weakly-dominated 2-dimensional closed manifold. In fact, for every genus  $\sigma > 1$ , the set of isomorphism classes of finite maximal subgroups of  $\text{Homeo}^+(S_\sigma)$  is finite with more than one element, while for  $\sigma = 0$  or 1, there are no maximal finite subgroups.*

Recall first that closed, oriented surfaces are classified by their genus  $0 \leq \sigma \in \mathbb{Z}$ . Clearly the surfaces of genus 0 (the sphere) and genus 1 (the torus) cannot be  $G$ -weakly-dominated for any  $G$  since each of them has self-automorphisms of unbounded finite



order. So from now on assume  $\sigma \geq 2$ . Now, if  $S = S_\sigma$  and  $G$  is a finite group of automorphisms, then by [30] and the Hurwitz upper bound  $|G| \leq 84(\sigma - 1)$ . The Riemann–Hurwitz formula [9] asserts that in this case, letting  $\bar{S} = S/G$ ,  $\pi : S \rightarrow \bar{S}$  the quotient map which is a ramified covering, ramified at  $0 \leq r \in \mathbb{Z}$  points with ramification indices  $m_1, \dots, m_r$  and if  $\bar{S}$  is of genus  $\rho$  then the following holds

$$2\sigma - 2 = |G|(2\rho - 2 + \sum_{i=1}^r (1 - \frac{1}{m_i})). \tag{1}$$

What is even more important for us is the converse. Namely if  $G$  is a finite group generated by elements  $a_1, \dots, a_\rho, b_1, \dots, b_\rho, c_1, \dots, c_r$  where

$$\prod_{j=1}^\rho [a_j, b_j] \prod_{j=1}^r c_j = 1 \text{ and} \tag{2}$$

$$\text{for } i = 1, \dots, r, \ c_i \text{ is of order } m_i \tag{3}$$

and if (1) holds, then  $G$  is a quotient of the fundamental group of a surface of genus  $\rho$  with  $r$  ramification points with ramification indices  $m_1, \dots, m_r$ . The kernel is a surface group of genus  $\sigma$ . Hence  $G$  acts faithfully on  $S = S_\sigma$  with quotient  $\bar{S} = S/G$  of genus  $\rho$  and ramification indices  $m_1, \dots, m_r$ .

The Hurwitz upper bound actually follows from equation (1): To get the largest  $G$ , for a fixed  $\sigma$ , one wants the term in brackets on the right-hand side to be minimal but positive. A careful elementary analysis shows that the smallest value is  $\frac{1}{42}$  and it is obtained only if  $\rho = 0, r = 3$  and  $\{m_1, m_2, m_3\} = \{2, 3, 7\}$ . One also observes that if this value is not attained then the next one is  $\frac{1}{24}$  with  $\rho = 0, r = 3$  and  $\{m_1, m_2, m_3\} = \{1, 2, 8\}$ . It is known that for infinitely many  $g$ 's, the upper bound of  $84(\sigma - 1)$  is attained but for infinitely many others it is not (cf. [33]). In the second case it follows that  $|G| \leq 48(\sigma - 1)$ .

The converse result allows one to prove that various groups act faithfully on  $S_\sigma$ . For example, by taking  $\rho = 2$  and  $r = 0$  we see that

- (a) The cyclic group  $c_{\sigma-1}$  of order  $\sigma - 1$  acts faithfully on  $S_\sigma$ .

Similarly, by taking  $\rho = 1, r = 2$  and  $m_1 = m_2 = \sigma$

- (b) The cyclic group  $C_\sigma$  of order  $\sigma$  acts faithfully on  $S_\sigma$ .

Finally, the following group

$$H_\sigma = \langle x, y | x^4 = y^{2(\sigma+1)} = (xy)^2 = x^{-1}y)^2 = 1 \rangle$$

is shown in [1] and [36] to be of order  $8(\sigma + 1)$ . (Note that by the two last relations every element of  $H_\sigma$  can be written as  $x^a y^b$  with  $0 \leq a < 4$  and  $0 \leq b < 2(\sigma + 1)$ ). By taking as 3 generators  $c_1 = x, c_2 = y$  and  $c_3 = (xy)^{-1}$  which are of orders 4,  $2(\sigma + 1)$  and 2,

respectively, one sees that equation (1) is satisfied with  $\rho = 0$ ,  $r = 3$  and  $(m_1, m_2, m_3) = (4, 2(\sigma + 1), 2)$  and hence:

(c) The group  $H_\sigma$  of order  $8(\sigma + 1)$  acts faithfully on  $S_\sigma$ .

Let us mention in passing that Accola [1] and Macmillan [36] used (c) to prove a lower bound (as an analogue to the upper bound of Hurwitz) and they showed that for infinitely many  $\sigma$ 's, this lower bound of  $8(\sigma + 1)$  is best possible.

Back to our goal: We want to show that  $S = S_\sigma$  cannot be  $G$ -weakly-dominated for any finite group  $G$ . Assume it is, then (a), (b) and (c) imply that  $C_{\sigma-1}, C_\sigma$  and  $H_\sigma$  are subgroups of  $G$  and hence:

$$\ell.c.m(\sigma - 1, \sigma, 8(\sigma + 1)) \mid |G|. \quad (4)$$

Now clearly  $\ell.c.m(\sigma - 1, \sigma, 8(\sigma + 1)) \geq \frac{(\sigma-1)\sigma(\sigma+1)}{2}$  and by the Hurwitz Theorem  $|G| \leq 84(\sigma - 1)$ . This implies  $\frac{1}{2}\sigma(\sigma + 1) \leq 84$ , i.e.,  $\sigma \leq 12$ .

Now checking case by case for  $\sigma = 6, 7, 9, 10, 11, 12$ , one sees that  $\ell.c.m(\sigma - 1, \sigma, 8(\sigma + 1)) > 84(\sigma - 1)$  in all these cases, which leads to a contradiction. We are left with  $\sigma = 8$  and  $2 \leq \sigma \leq 5$ .

For  $\sigma = 8$ , we observe that if  $S_8$  is  $G$ -weakly-dominated then by (b),  $C_8$  is a subgroup of  $G$  and so is  $H_8$  of (c). But  $C_8$  is cyclic, while the 2-sylow subgroup of  $H_8$  contains the *non-cyclic* subgroup of order 4, generated by  $xy$  and  $x^{-1}y$ . Thus the 2-sylow subgroup of  $G$  is non-cyclic and hence of order greater than 8, i.e., at least 16. This implies that  $G$  is of order at least  $7 \cdot 16 \cdot 9 > 84 \cdot 7$ . This contradicts the Hurwitz upper bound and hence  $S_8$  cannot be  $G$ -weakly-dominated.

To handle the case  $\sigma = 5$ , let us observe that the Hurwitz upper bound of  $84(\sigma - 1) = 336$  is obtained in this case. Indeed, look at  $G = SL_2(7)$ , a group of order 336 with the generators

$$c_1 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad c_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

and  $c_3$  where

$$c_3^{-1} = c_1 c_2 = \begin{pmatrix} -1 & 1 \\ -1 & 0 \end{pmatrix}$$

of orders 7, 2 and 3, respectively.

Thus if  $S_5$  is  $G$ -weakly-dominated,  $G$  must be  $SL_2(7)$  since Hurwitz bound is attained for this group. But  $SL_2(7)$  does not contain the group  $H_5$  of (c) of order 48, since 48 does not divide 336.

Consider now the case  $\sigma = 4$ . We claim that  $Sym(5)$  of order 120 acts on it. Indeed, taking  $\rho = 0, r = 3$  and

$$c_1 = (1, 2, 3, 4, 5), \quad c_2 = (1, 2) \quad \text{and} \quad c_3 = (c_1 c_2)^{-1} = (5, 4, 3, 1)$$

of orders 5, 2 and 4 respectively. We get a solution to (1) and hence  $Sym(5)$  acts on  $S_4$ . Assume now that  $S_4$  is  $G$ -weakly-dominated. The Hurwitz bound  $84.3 = 254$  cannot be obtained in such a case, since 120 does not divide 254. Thus  $G$  is of order at most  $48 \cdot 3 = 154$ , but we know that its order should be divisible by 120. Hence  $|G| = 120$  and  $G = Sym(5)$ . But (c) above shows that  $G$  should also contain  $H_4$  which is of order 40. As  $Sym(5)$  has no subgroup of order 40, we get a contradiction. Hence  $S_4$  is not  $G$ -weakly-dominated.

For  $\sigma = 2$  and 3, a full classification of the finite groups acting on  $S_\sigma$  is given in [10]. From the list there it is clear that  $S_\sigma$  is not  $G$ -weakly-dominated also in these last two cases. The Theorem is now fully proved.  $\square$

The Theorem says in particular that for every  $g$ ,  $Homeo^+(\Sigma_g)$  has at least two conjugacy classes of maximal finite subgroups (even isomorphism classes). In fact, the number of those is unbounded as a function of  $g$ :

**Proposition 6.2.** *The number of isomorphism classes of maximal finite subgroups of  $Homeo^+(\Sigma_g)$  (or equivalently of  $MCG(\Sigma_g)$ ) is unbounded as a function of  $g$ .*

**Proof.** Let  $\Gamma$  be a fixed cocompact lattice in  $PSL(2, R)$  which is the fundamental group of a surface of genus 2. This group is mapped onto  $F_2$  – the free group on 2 generators and  $F_2$  is mapped onto every 2-generated finite group. This holds therefore also for  $\Gamma$ . Every normal subgroup  $\Delta$  of  $\Gamma$  of index  $n$  defines a covering surface  $\Sigma_r$  when  $r = n + 1$ , for which  $\Gamma/\Delta$  serves as a group of (orientation preserving) isometries and hence define a finite subgroup of  $Homeo^+(\Sigma_r)$ . Now, the number of isomorphism classes of finite groups of order at most  $n$  which are generated by 2 elements is superpolynomial (in fact, this number of groups grows like  $n^{O(\log n)}$  – see [35]). Thus, there is an infinite set of  $r$ 's for which there is an unbounded number of nonisomorphic finite subgroups of  $Homeo^+(\Sigma_r)$  of order  $r - 1$ . Even if these subgroups are not maximal, there are also unboundedly many isomorphism classes of maximal subgroups. Indeed, each one of the above is of index at most 84 (by Hurwitz upper bound) in a maximal subgroup. Now, every maximal subgroup containing one of these subgroups is generated by at most  $2 + \log_2(84) < 9$  elements. The number of bounded index subgroups in any group with a bounded number of generators is uniformly bounded. This finishes the proof of the Proposition.  $\square$

## 7. Topological rigidity

In this section we prove [Theorem 1.7](#) and [1.8](#).

[Theorem 1.8](#) is proved essentially the same way as [Theorem 1.5](#), but two modifications are needed: [Theorem 1.5](#) was proved by restricting to actions of the cyclic group  $G = \mathbb{Z}/p$ , while now we need similar results for actions of general finite groups  $G$ . Let us indicate how the method of proof for  $\mathbb{Z}/p$  generalizes to general  $G$ .

- (1) Note that if  $H$  is any group acting semifreely (i.e., with only two kinds of orbits, fixed points and free orbits) the literally same proof as for  $\mathbb{Z}/p$  works.
- (2) Now if  $G$  is a finite group, consider the least singular of the singular points, i.e., the non-singular points of the singular set. Each of these will be fixed by some group  $H$ . The  $H$ -fixed set consists of points fixed by  $H$ , and maybe also some more singular points fixed by a larger group. We will do our modification near points that are fixed only by  $H$  or by a conjugate of it. These points are the  $G$  orbit of points fixed just by  $H$ .
- (3) The modification will be done as follows. Start with a semi-free  $H$ -sphere  $S$  with fixed set  $\Sigma$  and normal representation – the  $H$  representation that occurs at a fixed point of  $H$  (that is on the top stratum of the singular set, as in # 2). We can consider the product space  $(G \times S)/H$ , where  $H$  acts on the left on  $G$  and the right on  $S$ . So,  $G$  acts on this product. The underlying topological space is  $G/H \times S$ , but the action is more interesting. It is called the induction of the  $H$ -action on  $S$  to  $G$ .
- (4) Now take connected sum along an orbit of  $M$  with  $(G \times S)/H$ . It is homeomorphic to  $M$ , but the singular set is modified by connect sums of copies of  $\Sigma$  in various places.
- (5) Similar tricks work when we do Edwards modifications.

With the above modification all the results proved in Sections 2–5 can be modified to work with general finite group  $G$ .

The second modification is easier: We should think of proper discontinuous actions of  $\Gamma$  on  $H/K$  as follows. Let  $\Delta$  be a normal finite index torsion free subgroup of  $\Gamma$ . Then  $M = \Delta \backslash H/K$  is a compact manifold upon which  $G = \Gamma/\Delta$  acts. Note that  $M$  is indeed compact whatever the (proper discontinuous) action of  $\Gamma$  on  $H/K$  is, since the cohomological dimension of  $\Delta$  is  $\dim(H/K)$  as deduced from its original isometric action. Now, taking the above mentioned modification (from  $\mathbb{Z}/p$  to  $G$ ), [Theorem 1.8](#) is deduced from [Theorem 1.5](#) by standard covering space theory, changing from  $M$  to its universal cover  $H/K$ . Note, however, that the formulation of the two theorems is slightly different due to the fact that  $\gamma \in \Gamma$  has a fixed point in  $H/K$  if and only if it is of finite order. Also, an automorphism of  $M = \Delta \backslash H/K$  with a fixed point can be lifted to an element of finite order in  $\Gamma = N_H(\Delta)$ .

[Theorem 1.7](#) is essentially equivalent to [Corollary 1.6](#), but one needs to ensure that when there are countably many proper discontinuous actions of  $\Gamma$  on  $H/K$ , these actions are isolated, i.e., a small perturbation of each such action is conjugate to it. This is indeed the case (in contrast with the theory of deformations into Lie groups in which case, if there are infinitely many cocompact discrete representations, then local rigidity fails and there are continuously many such actions.) The point is that when the fixed points form a discrete set one does have a local topological rigidity (and even in dimension 4). This follows from Edmonds' Theorem [[17](#), [Theorem 2.8](#)] in high dimensions. The work of [[22](#)] (see also [[20](#)]) shows it is true also in dimension 4. The case of dimension 3 is always covered by either [Theorem 1.5](#) (a) or (c), and case (b) does not happen.  $\square$

## References

- [1] R. Accola, On the number of automorphisms of a closed Riemann surface, *Trans. Amer. Math. Soc.* 131 (1968) 398–407.
- [2] A. Bartels, W. Lueck, The Borel conjecture for hyperbolic and CAT(0)-groups, *Ann. of Math.* 2 (175) (2012) 631–689.
- [3] H. Bass, A. Heller, R.G. Swan, The Whitehead group of a polynomial extension, *Publ. Math. Inst. Hautes Études Sci.* 22 (1964) 61–79.
- [4] M. Belolipetsky, A. Lubotzky, Finite groups and hyperbolic manifolds, *Invent. Math.* 162 (2005) 459–472.
- [5] R.H. Bing, Inequivalent families of periodic homeomorphisms of  $E^3$ , *Ann. of Math.* 80 (2) (1964) 78–93.
- [6] A. Borel, On period maps of certain  $K(\pi, 1)$ , in: *Borel Collected Papers III*, Springer, 1983, pp. 57–60.
- [7] A. Borel, *Seminar on Transformation Groups*, *Annals of Math. Studies*, vol. 46, Princeton University Press, 1960.
- [8] G. Bredon, *Introduction to Compact Transformation Groups*, Academic Press, 1972.
- [9] T. Breuer, *Characters and Automorphism Groups of Compact Riemann Surfaces*, Cambridge University Press, 2001.
- [10] S. Broughton, Classifying finite group actions on surfaces of low genus, *J. Pure Appl. Algebra* 69 (1990) 233–270.
- [11] J. Cannon, Shrinking cell-like decompositions of manifolds. Codimension three, *Ann. of Math.* (2) 110 (1) (1979) 83–112.
- [12] S.E. Cappell, Splitting obstructions for Hermitian forms and manifolds with  $Z^2 \subset \Pi_1$ , *Bull. Amer. Math. Soc.* 79 (1973) 909–913.
- [13] P.E. Conner, E.E. Floyd, *Differentiable Periodic Maps*, *Ergebnisse der Mathematik und Ihrer Grenzgebiete, N. F.*, vol. 33, Academic Press Inc./Springer-Verlag, New York/Berlin–Göttingen–Heidelberg, 1964, vii+148 pp.
- [14] F. Connolly, T. Koźniewski, Examples of lack of rigidity in crystallographic groups, in: *Algebraic Topology Poznań, 1989*, in: *Lecture Notes in Math.*, vol. 1474, Springer, Berlin, 1991, pp. 139–145.
- [15] F. Connolly, J.F. Davis, Q. Khan, Topological rigidity of actions on contractible manifolds with discret singular set, arXiv:1402.0280.
- [16] R.J. Daverman, *Decompositions of Manifolds*, *Pure and Applied Mathematics*, vol. 124, Acad. Press, Orlando, FL, 1986, p. xii+317.
- [17] A. Edmonds, Local connectivity of spaces of group actions, *Quad. Mat.* 27 (1976) 71–84.
- [18] F.T. Farrell, L. Jones, A topological analogue of Mostow’s rigidity theorem, *J. Amer. Math. Soc.* 2 (2) (1989) 257–370.
- [19] F.T. Farrell, L. Jones, *Topological Rigidity for Compact Non-positively Curved Manifolds in Differential Geometry: Riemannian Geometry*, Amer. Math. Society, Providence, RI, 1993, pp. 229–274.
- [20] S. Ferry, S. Weinberger, Curvature, tangentiality, and controlled topology, *Invent. Math.* 105 (1991) 401–414.
- [21] M.H. Freedman, The topology of four-dimensional manifolds, *J. Differential Geom.* 17 (3) (1982) 357–453.
- [22] M. Freedman, F. Quinn, *Topology of 4-Manifolds*, Princeton University Press, 1990.
- [23] D.Gabai.G.R. Meyerhoff, N. Thurston, Homotopy hyperbolic 3-manifolds are hyperbolic, *Ann. of Math.* (2) 157 (2003) 335–431.
- [24] C. Giffen, The generalized Smith conjecture, *Amer. J. Math.* 88 (1966) 187–198.
- [25] L. Greenberg, Maximal Fuchsian groups, *Bull. Amer. Math. Soc.* 69 (1963) 569–573.
- [26] M. Gromov, I. Piatetski-Shapiro, Nonarithmetic groups in Lobachevsky spaces, *Publ. Math. Inst. Hautes Études Sci.* 66 (1988) 93–103.
- [27] S. Illman, Equivariant absorptions and recognition of linear actions on spheres, (Russian). *Topology, Moscow, 1979 Tr. Mat. Inst. Steklova* 154 (1983) 124–126.
- [28] W. Jaco, P. Shalen, Seifert fibered spaces in 3-manifolds, *Mem. Amer. Math. Soc.* 21 (1979), viii+192 pp.
- [29] K. Johansson, *Homotopy Equivalences of 3-Manifolds with Boundaries*, LNM, vol. 761, Springer-Verlag, 1979.
- [30] B. Kerékjártó, Über die periodischen transformationen der Kreisscheibe und der Kugelfläsche, *Math. Ann.* 80 (1919) 36–38.

- [31] S. Kerckhoff, The Nielsen realization problem, *Ann. of Math. (2)* 117 (1983) 235–265.
- [32] M. Kervaire, Smooth homology spheres and their fundamental groups, *Trans. Amer. Math. Soc.* 144 (1969) 67–72.
- [33] M. Larsen, How often is  $84(g - 1)$  achieved?, *Israel J. Math.* 126 (2001) 1–16.
- [34] F. Latour, Double suspension d’une sphère d’homologie [d’après R. Edwards], in: *Séminaire Bourbaki vol. 1977/78 Exposés 507–524*, in: *Lecture Notes in Math.*, vol. 710, Springer-Verlag, Berlin, New York, 1979, pp. 169–186, in French.
- [35] A. Lubotzky, Enumerating boundedly generated finite groups, *J. Algebra* 238 (1) (2001) 194–199.
- [36] C. Maclachlan, A bound for the number of automorphisms of a compact Riemann surface, *J. Lond. Math. Soc. (2)* 44 (1969) 265–276.
- [37] J. Milnor, Lectures on the  $h$ -cobordism theorem, in: *Notes by L. Siebenmann and J. Sondow*, Princeton University Press, Princeton, N.J., 1965, v+116 pp.
- [38] J. Milnor, On the 3-Dimensional Brieskorn Manifolds  $M(p, q, r)$ . Knots, Groups, and 3-Manifolds, Papers dedicated to the memory of R. H. Fox *Ann. of Math. Studies*, vol. 84, Princeton Univ. Press, Princeton, N.J., 1975, pp. 175–225.
- [39] G.D. Mostow, Strong Rigidity of Locally Symmetric Spaces, *Annals of Mathematics Studies*, vol. 78, Princeton Univ. Press/University of Tokyo Press, Princeton, N.J./Tokyo, 1973, v+195 pp.
- [40] F. Quinn, Applications of topology with control, in: *Proceedings of the International Congress of Mathematicians, Vol. 1, 2*, Berkeley, CA, 1986, Amer. Math. Soc., Providence, RI, 1987, pp. 598–606.
- [41] R. Schultz, Group actions on hypertoral manifolds. I, in: *Topology Symposium, Siegen 1979, Proc. Sympos.*, Univ. Siegen, Siegen, 1979, in: *Lecture Notes in Math.*, vol. 788, Springer, Berlin, 1980, pp. 364–377.
- [42] R. Schultz, Group actions on hypertoral manifolds. II, *J. Reine Angew. Math.* 325 (1981) 75–86.
- [43] J. Stallings, The piecewise-linear structure of Euclidean space, *Math. Proc. Cambridge Philos. Soc.* 58 (1962) 481–488.
- [44] J. Stallings, On infinite processes leading to differentiability in the complement of a point, in: *Differential and Combinatorial Topology (A Symposium in Honor of Marston Morse)*, Princeton Univ. Press, Princeton, N.J., 1965, pp. 245–254.
- [45] S. Weinberger, *The Topological Classification of Stratified Spaces*, University of Chicago Press, 1994.
- [46] S. Weinberger, Variations on a theme of Borel, preprint, 2015.
- [47] E.C. Zeeman, Twisting spun knots, *Trans. Amer. Math. Soc.* 115 (1965) 471–495.