

# An Etale Approach to the Novikov Conjecture

A.N. DRANISHNIKOV

*University of Florida, Gainesville, FL 32611-8105, USA*

S. FERRY

*Department of Mathematics, Rutgers University, Piscataway, NJ 08854-8019*

AND

S. WEINBERGER

*Department of Mathematics, The University of Chicago, Chicago, IL 60637*

## Abstract

We show that the rational Novikov conjecture for a group  $\Gamma$  of finite homological type follows from the mod 2 acyclicity of the Higson compactification of an  $E\Gamma$ . We then show that for groups of finite asymptotic dimension the Higson compactification is mod  $p$  acyclic for all  $p$ , and deduce the integral Novikov conjecture for these groups. © 2000 Wiley Periodicals, Inc.

## 1 Introduction

Ten years ago, the most popular approach to the Novikov conjecture went via compactifications. If a compact aspherical manifold, say, had a universal cover which suitably equivariantly compactified, already Farrell and Hsiang [20] proved that the Novikov conjecture follows. Subsequent work by many authors weakened the various hypotheses and extended the idea to other settings. (See e.g. [9, 22, 37].)

In recent years, other coarse methods (most notably the embedding method of [41] in the  $C^*$ -algebra setting) have supplanted the compactification method. The reason for this is that the embedding method seemed to have a better chance of applying generally, since compactifications effective for the Novikov conjecture seem to require some special geometry for their construction. For a brief period, it seemed that a general construction might solve the Novikov conjecture. Higson introduced a general compactification of metric spaces (somewhat reminiscent of the Stone-Ćech compactification) that automatically has most of the needed properties for application to the Novikov conjecture. The Novikov conjecture would follow if it could be proven that the Higson compactification of the universal cover of an aspherical polyhedron is acyclic. (See [37].)

Unfortunately, it was soon realized by Keesling and others (see [29, 15]) that the Higson compactification, even for manifolds as small as  $\mathbb{R}$ , has nontrivial rational cohomology. Thus, it was felt that such universal compactifications were not suitable for Novikov. Recently, however, Gromov has harpooned the embedding approach by constructing finitely generated groups that do not uniformly embed in Hilbert space [23] and a number of authors (e.g. [25, 32]) have shown that Gromov's groups can be used to construct counterexamples to general forms of the Baum-Connes conjecture. Currently, however, there are no candidates for counterexamples to the Novikov conjecture nor is there a potential method for proving the conjecture.

Moreover, for reasons that are not entirely clear, the embedding method has never been translated into pure topology; the results on integral Novikov conjecture so obtained, from the L-theoretic viewpoint never account for the prime 2, and there do not seem to be many results in pure algebraic K-theory (or A-theory) provable by this method.

This paper seeks to rehabilitate the approach to the Novikov conjecture via the Higson compactification. We do this by examining the Higson compactification with finite coefficients.

**Theorem 1.** *If the Higson compactification of  $E\pi$  is mod 2 acyclic, and  $B\pi$  is a finite complex, then the integral Novikov conjecture for  $\pi$  holds at the prime 2.*

In other words, the assembly map localized at the prime two is an injection on homotopy groups. At odd primes we do not know how to prove any corresponding statement, in general, for reasons related to the examples in [16]: The L-spectrum away from 2 is periodic K-theory, and cohomologically acyclic spaces may not be acyclic for periodic K-theory.

This difficulty is an infinite dimensional phenomenon, so for groups of *finite asymptotic dimension* this issue does not arise, and it is possible to prove integral results.

**Theorem 2.** *If  $B\pi$  is a finite complex and  $\pi$  has finite asymptotic dimension, then if  $\mathbf{R}$  is a ring such that  $\mathbf{L}(\mathbf{R})$  is finitely generated, then the  $\mathbf{L}(\mathbf{R})$  assembly map for  $\pi$  is injective.*

In this paper, by  $\mathbf{L}(\mathbf{R})$  we will always mean  $\mathbf{L}^{-\infty}(\mathbf{R})$ .

This is not a new theorem. For  $K(C^*\pi)$ , this is due to [44]. In algebraic K-theory there are papers by [1, 8], which apply to L-theory as well (without the restrictions on  $\mathbf{R}$ ); moreover [11] has yet another approach to the stronger L-theory result. The

novelty here is the method of using finite primes to regain the acyclicity of the Higson compactification and then using the etale idea to get our Novikov result.<sup>1</sup>

Theorem 2 follows from the general discussion above, together with our third theorem.

**Theorem 3.** *If  $B\pi$  is a finite complex and  $\pi$  has finite asymptotic dimension, then the Higson compactification of  $E\pi$  is mod  $p$  acyclic for every prime  $p$ .*

To motivate and clarify our results, we close with a brief discussion of the Higson compactification and of the Keesling elements. If  $X$  is a metric space, and  $f : X \rightarrow \mathbb{R}$  is a bounded continuous function, we say that  $f$  has *decaying variation* if

$$\lim_{r \rightarrow \infty} \text{diam}(f(B_r(x))) = 0$$

for any fixed  $r$  and any sequence of points  $x$  going to infinity in  $X$ . Higson's compactification is the smallest one containing  $X$  densely so that all bounded continuous functions with decaying variation extend. Let  $C_h$  be the set of such functions. We obtain the Higson compactification of  $X$  by embedding  $X$  into  $\prod_{f \in C_h} \mathbb{R}$  and taking the closure of the image. This closure is compact by the Tychonoff theorem, since the maps  $f$  are bounded.

Let us now consider the first cohomology of the compactification of a uniformly simply connected  $X$ . We represent an element of  $\check{H}^1(\bar{X})$  by a map  $f : \bar{X} \rightarrow S^1$ . On  $X$ , we can lift  $f|_X$  and view our class as being given by  $\exp(2\pi i \tilde{f})$ , where  $\tilde{f}$  has decaying variation but is not necessarily bounded. Maps  $f, g : \bar{X} \rightarrow S^1$  represent the same cohomology class iff  $\tilde{f} - \tilde{g}$  is bounded.

Note that  $H^1(\bar{X}; \mathbb{Z})$  is now an  $\mathbb{R}$ -vector space – one can multiply the function  $\tilde{f}$  by a real number, project back to  $S^1$ , and extend to  $\bar{X}$ . In particular, this cohomology vanishes with mod  $p$  coefficients. In this regard, the cohomology of the Higson compactification resembles the uniformly finite homology of a uniformly contractible  $n$  manifold, [4], which is known to be an  $\mathbb{R}$  vector space, aside from the class coming from the fundamental class of  $X$ .

Our paper is organized as follows. Section 2 is devoted to *descent*, i.e. deducing Novikov conjectures from metric results on the universal cover. The last section is devoted to verifying the mod  $p$  acyclicity result for the finite asymptotic dimension case. This is done using ideas of *quantitative algebraic topology*. The key idea is that when homotopy groups are finite, it is sometimes possible to get extra a priori Lipschitz conditions on maps.

---

<sup>1</sup>Note that the celebrated paper of [5] on the algebraic K-theory analogue of the Novikov conjecture is a very nice illustration of a completely different etale idea. We thank Alain Connes for suggesting this evocative description of our method.

## 2 L-theory and assembly map with $\mathbf{Z}_p$ coefficients

For every ring with involution  $R$  (even for any additive category with involution) Ranicki defines a 4-periodic spectrum  $\mathbf{L}_*(R)$  with  $\pi_i(\mathbf{L}_*(R)) = L_i(R)$  [34, 35]. We use the notation  $\mathbf{L} = \mathbf{L}_*(\mathbf{Z})$ . Strictly speaking,  $\mathbf{L}$ -spectra are indexed by  $K$ -groups. Our  $\mathbf{L}(\mathbf{R})$  equals  $\mathbf{L}^{-\infty}(\mathbf{R})$ , the limit of the  $\mathbf{L}$ -spectra associated to the negative  $K$ -groups. This applies to all  $\mathbf{L}$ -spectra occurring in this paper.

For a metric space  $(X, d)$  we will denote by  $C_X(R)$  the boundedly controlled Peder- sen-Weibel category whose objects are locally finite direct sums  $A = \bigoplus_{x \in X} A(x)$  of finite dimensional free  $R$ -modules and whose morphisms are given by matrices with bounded propagations (see [21]). For a subset  $V \subset X$  we denote by  $A(V)$  the sum  $\bigoplus_{x \in V} A(x)$ . Ranicki [35] defined  $X$ -bounded quadratic  $L$ -groups  $L_*(C_X(R))$  and the corresponding spectrum  $\mathbf{L}_*(C_X(R))$ . We will use the notation  $\mathbf{L}^{bdd}(X) = \mathbf{L}_*(C_X(\mathbf{Z}))$ .

Suppose that  $\bar{X}$  is a compactification of  $X$  with corona  $Y = \bar{X} \setminus X$ . Then one can define the *continuously controlled category*  $B_{X,Y}(R)$  by taking the same objects as above with morphisms  $f : A \rightarrow B$  in  $B_{X,Y}(R)$  taken to be homomorphisms such that for every  $y \in Y$  and every neighborhood  $y \in U \subset \bar{X}$  there is a smaller neighborhood  $y \in V \subset U$  such that  $f(A(V)) \subset A(U)$ . This category is additive and hence also admits an  $L$ -theory. The corresponding spectrum for  $R = \mathbf{Z}$  we denote by  $\mathbf{L}^{cc}(X) = \mathbf{L}_*(B_{X,Y}(\mathbf{Z}))$ .

If  $\mathbf{E} = \{E_k \mid k \in \mathbf{Z}\}$  is a spectrum, we will denote the  $\mathbf{E}$ -homology of  $X$  by  $H_i(X; \mathbf{E})$ . If  $X$  is a complex, then  $H_i(X; \mathbf{E}) = \pi_i(X_+ \wedge \mathbf{E}) = \lim_{k \rightarrow \infty} \pi_{i+k}(X_+ \wedge E_k)$ . If  $X$  is a compact metric space, we denote the Steenrod  $\mathbf{E}$ -homology of  $X$  by  $H_i(X; \mathbf{E})$ , i.e.  $H_i(X; \mathbf{E}) = \pi_i(\text{holim}\{N_+^i \wedge \mathbf{E}\})$  where  $X = \varprojlim N^i$  is the inverse limit of polyhedra [CP2], [EH].

The following theorem was proven in [33, 9, 10].

**Theorem 2.1.**  $L_i(B_{X,Y}(R)) = \bar{H}_{i-1}(Y; \mathbf{L}(R))$  for all  $i$  where the homology on the right is reduced Steenrod  $\mathbf{L}_*(R)$ -homology.

A metric space  $X$  is called *proper* if every ball  $B_r(x)$  in  $X$  is compact. Subsets  $A, B \subset X$  of a proper metric space  $(X, d)$  are called *diverging* if

$$\lim_{r \rightarrow \infty} d(A \setminus B_r(x_0), B \setminus B_r(x_0)) = \infty$$

where  $x_0 \in X$  is any basepoint,  $B_r(x_0)$  is the  $r$ -ball centered at  $x_0$ , and the distance  $d(A', B')$  between sets is the infimum of distances between  $d(a, b)$ ,  $a \in A'$ ,  $b \in B'$ . We note that the minimal compactification of a proper metric space  $X$  with respect to the property that the closures of every pair of diverging subsets in  $X$  have empty intersection in the corona is the Higson compactification.

A compactification  $\tilde{X}$  of  $X$  is called *Higson dominated* if the identity map  $1_X : X \rightarrow X$  admits a continuous extension  $\tilde{X} \rightarrow \tilde{X}$ , where  $\tilde{X}$  is the Higson compactification. We note that for every Higson dominated compactification  $\tilde{X}$  of  $X$  with corona  $Y = \tilde{X} - X$ , there is a forgetful functor  $C_X(R) \rightarrow B_{X,Y}(R)$ . This functor defines a map of spectra  $\phi : \mathbf{L}^{bdd}(X) \rightarrow \mathbf{L}^{cc}(X)$ .

If  $X$  is the universal covering of a finite complex with fundamental group  $\Gamma$ , then  $\Gamma$  acts on  $C_X(R)$  and hence on  $\mathbf{L}_*(C_X(R))$  with fixed set  $\mathbf{L}_*(C_X(R))^\Gamma = \mathbf{L}_*(R\Gamma)$  [9]. If a compactification of  $X$  with corona  $Y$  is equivariant, then  $\Gamma$  acts on  $B_{X,Y}(R)$  and hence on  $\mathbf{L}_*(B_{X,Y}(R))$ .

We recall that the homotopy fixed set  $X^{h\Gamma}$  of a pointed space  $X$  with a  $\Gamma$  action on it is defined as the space of equivariant maps  $\text{Map}_\Gamma(\mathbf{E}\Gamma_+, X)$ .

For a general spectrum  $\mathbf{E}$  and a torsion free group  $\Gamma$  it was proven [6, 9] that

$$\mathbf{H}_*(B\Gamma; \mathbf{E}) \cong \mathbf{H}_*^{lf}(\mathbf{E}\Gamma; \mathbf{E})^\Gamma \cong \mathbf{H}_*^{lf}(\mathbf{E}\Gamma; \mathbf{E})^{h\Gamma}.$$

For a locally compact space  $X$ ,  $H_i^{lf}(X; \mathbf{E})$  is defined to be the Steenrod  $\mathbf{E}$ -homology rel infinity of the one point compactification  $\alpha X$ . We note that for a complex  $N$ ,  $\mathbf{H}_*(N; \mathbf{E}) = N_+ \wedge \mathbf{E}$  and for a compact space  $Y$ ,  $\mathbf{H}_*(Y; \mathbf{E}) = \text{holim}\{N_+^\alpha \wedge \mathbf{E}\}$  where  $N^\alpha$  runs over nerves of all finite open covers of  $Y$ . If  $\Gamma$  acts on a compact space  $Y$ , then it acts on the set of all finite open covers of  $Y$  and hence on the spectrum  $\mathbf{H}_*(Y; \mathbf{E})$ .

The following theorem is due to Carlsson-Pedersen [9, 10]. See also Ranicki [36]. The first part of it is discussed in a more general setting in [39], Theorem 3.3.

**Theorem 2.2.** *For every group  $\Gamma$  with finite classifying complex  $B\Gamma$  there is a morphism of spectra called the bounded control assembly map*

$$A^{bdd} : \mathbf{H}_*^{lf}(\mathbf{E}\Gamma; \mathbf{L}) \rightarrow \mathbf{L}^{bdd}(\mathbf{E}\Gamma)$$

which fits into a homotopy commutative diagram

$$\begin{array}{ccc} \mathbf{H}_*(B\Gamma; \mathbf{L}) & \xrightarrow{A} & \mathbf{L}_*(Z\Gamma) \\ \simeq \downarrow & & \text{trf} \downarrow \\ \mathbf{H}_*^{lf}(\mathbf{E}\Gamma; \mathbf{L})^{h\Gamma} & \xrightarrow{A^{bdd, h\Gamma}} & \mathbf{L}^{bdd}(\mathbf{E}\Gamma)^{h\Gamma} \end{array}$$

where  $A$  is the standard assembly map, and the vertical arrows are the natural maps from fixed sets to homotopy fixed sets.

If the universal covering space  $E\Gamma$  admits a Higson dominated equivariant compactification  $X$  with corona  $Y$ , then the diagram can be extended

$$\begin{array}{ccc}
\mathbf{H}_*(\mathbf{B}\Gamma; \mathbf{L}) & \xrightarrow{A} & \mathbf{L}_*(\mathbf{Z}\Gamma) \\
\cong \downarrow & & \text{tr}f \downarrow \\
\mathbf{H}_*^{lf}(\mathbf{E}\Gamma; \mathbf{L})^{h\Gamma} & \xrightarrow{A^{bdd, h\Gamma}} & \mathbf{L}^{bdd}(\mathbf{E}\Gamma)^{h\Gamma} \\
= \downarrow & & \phi^{h\Gamma} \downarrow \\
\mathbf{H}_*^{lf}(\mathbf{E}\Gamma; \mathbf{L})^{h\Gamma} & \xrightarrow{A^{cc, h\Gamma}} & \mathbf{L}^{cc}(\mathbf{E}\Gamma)^{h\Gamma}
\end{array}$$

where  $A^{cc, h\Gamma} = \phi^{h\Gamma} \circ A^{bdd, h\Gamma}$ . Moreover,  $A^{cc}$  coincides with the boundary map in the Steenrod  $\mathbf{L}$ -homology exact sequence of pair  $(X, Y)$ :

$$A_p^{cc} = \partial : \mathbf{H}_*^{lf}(\mathbf{E}\Gamma; \mathbf{L}) \rightarrow \mathbf{H}_{*-1}(Y; \mathbf{L}).$$

Let  $M(p)$  denote the Moore spectrum for the group  $\mathbf{Z}_p = \mathbf{Z}/p\mathbf{Z}$ ,  $p$  prime.

**Lemma 2.3.** *Let  $X$  be a finite-dimensional compact metric space. Suppose that  $X$  is  $\mathbf{Z}_p$ -acyclic, i.e.,  $\tilde{H}^*(X; \mathbf{Z}_p) = 0$  for reduced Čech cohomology. Then  $X$  is acyclic for reduced Steenrod  $\mathbf{L} \wedge M(p^k)$ -homology for all  $k$ .*

*Proof.* In view of the universal coefficient formula [30], we note that  $\tilde{H}_*(X; \mathbf{Z}_p) = 0$  for reduced Steenrod homology. The coefficient exact sequence shows that  $\tilde{H}_*(X; \mathbf{Z}_{p^k}) = 0$  for all  $k$ . For every spectrum  $\mathbf{S}$  there is a Steenrod homology Atiyah-Hirzebruch spectral sequence with

$$E_{i,j}^2 = \tilde{H}_i(X; H_j(S^0, \mathbf{S}))$$

which converges to  $\tilde{H}_*(X; \mathbf{S})$  provided that  $X$  is a finite dimensional compact metric space [19, 28]. Since all of the groups  $H_q(S^0, \mathbf{L} \wedge M(p^k))$  are  $p$ -primary, we have  $E_{i,j}^2 = 0$  for all  $i$  and  $j$ .  $\square$

*Remark 2.4.* The finite dimensionality condition is essential here. There are acyclic compacta that have nontrivial mod  $p$  complex  $K$ -theory [42]. Namely, by results of Adams and Toda there is a map  $f : \Sigma^k M(\mathbf{Z}_p, m) \rightarrow M(\mathbf{Z}_p, m)$  which induces an isomorphism in  $K$ -theory. The inverse limit  $X$  of suspensions of  $f$  is an acyclic compactum, since all bonding maps are trivial in cohomology, yet it has nontrivial  $K$ -theory and nontrivial mod  $p$   $K$ -theory. Hence, for  $p$  odd,  $X$  has nontrivial mod  $p$   $L$ -theory.

*Remark 2.5.* The spectrum  $\mathbf{L}$  localized at 2 is equivalent (see [31]) to the Eilenberg-MacLane 4-periodic spectrum generated by the space

$$\prod_{i=1}^{\infty} K(\mathbf{Z}_{(2)}, 4i) \times K(\mathbf{Z}_2, 4i - 2).$$

Thus,  $\mathbf{L} \wedge M(2) = \mathbf{L}_{(2)} \wedge M(2)$  is the Eilenberg-MacLane spectrum generated by the space

$$\prod_{i=1}^{\infty} K(\mathbf{Z}_2, 4i) \times K(\mathbf{Z}_2, 4i - 1) \times K(\mathbf{Z}_2, 4i - 2).$$

Indeed, for any  $R$ ,  $\mathbf{L}(R)$  is Eilenberg-MacLane at 2 [43]. Thus, if a compactum  $X$  is  $\mathbf{Z}_2$ -acyclic, then it is  $\mathbf{L} \wedge M(2)$ -acyclic without any finite dimensionality assumption on  $X$ .

**Definition 2.6.** The mod  $p$  assembly map  $A_p$  is

$$A \wedge 1_{M(p)} : \mathbf{H}_*(B\Gamma; \mathbf{L}) \wedge M(p) \rightarrow \mathbf{L}_*(\mathbf{Z}\Gamma) \wedge M(p).$$

**Theorem 2.7.** *Suppose that the Higson compactification of the universal cover  $X$  of a finite aspherical complex  $B$  is finite-dimensional and acyclic in Steenrod mod  $p$  homology. Then the mod  $p^k$  assembly map for  $\Gamma = \pi_1(B)$  is a split monomorphism for every  $k$ .*

*Proof.* Let let  $m$  be the dimension of the Higson corona  $\dim vX$ . Using Schepin's spectral theorem [12] one can obtain a metrizable  $\mathbf{Z}_p$ -acyclic  $\Gamma$ -equivariant compactification  $\bar{X}$  of  $X$  with corona  $Y$  such that  $\dim Y = m$  (see the proof of Lemma 8.3. in [12]). We introduce coefficients to the second diagram of Theorem 2.2 by forming the smash product with  $M(p^k)$ .

$$\begin{array}{ccc} \mathbf{H}_*(B; \mathbf{L}) \wedge M(p^k) & \xrightarrow{A_p} & \mathbf{L}_*(\mathbf{Z}\Gamma) \wedge M(p^k) \\ \simeq \downarrow & & \downarrow \phi^{h\Gamma} \text{tr } f \wedge 1 \\ \mathbf{H}_*^{lf}(X; \mathbf{L})^{h\Gamma} \wedge M(p^k) & \xrightarrow{A^{cc, h\Gamma} \wedge 1} & \mathbf{L}^{cc}(X)^{h\Gamma} \wedge M(p^k) \end{array}$$

Note also that  $\mathbf{L} \wedge p^k = \text{cof}(\times p^k : \mathbf{L} \rightarrow \mathbf{L})$  and  $\times p^k$  commutes with all of the structure maps defining the spectral structure and the  $\Gamma$ -action. We therefore have the commutative diagram

$$\begin{array}{ccc} \mathbf{H}_*(B; \mathbf{L}) \wedge M(p^k) & \xrightarrow{A_p} & \mathbf{L}_*(\mathbf{Z}\Gamma) \wedge M(p^k) \\ \simeq \downarrow & & \downarrow \phi^{h\Gamma} \text{tr } f \wedge 1 \\ (\mathbf{H}_*^{lf}(X; \mathbf{L}) \wedge M(p^k))^{h\Gamma} & \xrightarrow{(A^{cc} \wedge 1)^{h\Gamma}} & (\mathbf{L}^{cc}(X) \wedge M(p^k))^{h\Gamma}. \end{array}$$

It suffices to show that

$$A^{cc} \wedge 1 : \mathbf{H}_*^{lf}(X; \mathbf{L}) \wedge M(p^k) \rightarrow \mathbf{L}^{cc}(X) \wedge M(p^k)$$

is an isomorphism because this implies that  $(A^{cc} \wedge 1)^{h\Gamma}$  is an isomorphism and hence that  $A_{p^k}$  is a split monomorphism.

In view of Theorems 2.1 and 2.2 we need to show that

$$\partial \wedge 1 : \mathbf{H}_*^{lf}(X; \mathbf{L}) \wedge M(p^k) \rightarrow \mathbf{H}_{*-1}(Y; \mathbf{L}) \wedge M(p^k)$$

is isomorphism. Note that  $\partial \wedge 1_{M(p^k)}$  is equivalent to the boundary homomorphism for the pair  $(\bar{X}, Y)$  in  $\mathbf{L} \wedge M(p^k)$ -homology. By Lemma 2.3,  $\mathbf{H}_*(\bar{X}; \mathbf{L} \wedge M(p^k)) = \mathbf{H}_*(pt; \mathbf{L} \wedge M(p^k))$  and hence  $\partial \wedge 1_{M(p^k)}$  is indeed an isomorphism.  $\square$

It is known that the universal coefficient formula with  $\mathbf{Z}_p$  coefficients (UCF) holds true for every generalized homology theory and is natural with respect to morphisms of spectra:

**Proposition 2.8.** *For every morphism of spectra  $A : \mathbf{E}_1 \rightarrow \mathbf{E}_2$  and every  $p$  and  $i$  there is a commutative diagram*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \pi_i(\mathbf{E}_1) \otimes \mathbf{Z}_p & \longrightarrow & \pi_i(\mathbf{E}_1 \wedge M(p)) & \longrightarrow & \pi_{i-1}(\mathbf{E}_1) * \mathbf{Z}_p \longrightarrow 0 \\ & & \downarrow A_* \otimes 1 & & \downarrow (A \wedge 1)_* & & \downarrow (A_{*-1})_* \otimes 1 \\ 0 & \longrightarrow & \pi_i(\mathbf{E}_2) \otimes \mathbf{Z}_p & \longrightarrow & \pi_i(\mathbf{E}_2 \wedge M(p)) & \longrightarrow & \pi_{i-1}(\mathbf{E}_2) * \mathbf{Z}_p \longrightarrow 0. \end{array}$$

*Proof.* We apply the smash product with  $\mathbf{E}_i$ ,  $i = 1, 2$  to the cofibration of spectra  $\mathbf{S} \rightarrow \mathbf{S} \rightarrow M(p)$ , where  $\mathbf{S}$  is the sphere spectrum. Then the result follows from the homotopy exact sequence of the resulting cofibrations of spectra and the induced morphism between them

$$\begin{array}{ccccc} E_1 & \longrightarrow & E_1 & \longrightarrow & E_1 \wedge M(p) \\ \downarrow & & \downarrow & & \downarrow \\ E_2 & \longrightarrow & E_2 & \longrightarrow & E_2 \wedge M(p). \end{array}$$

$\square$

**Theorem 2.9.** *Suppose that the the universal cover of a finite complex  $B\Gamma$  has finite dimensional Higson compactification which is mod  $p$  acyclic for every prime  $p$ . Then the integral assembly map  $A$  is a monomorphism.*

*Proof.* In view of compactness of  $B\Gamma$  we have  $v\Gamma = vE\Gamma$ . By Theorem 2.7  $A \otimes 1_G$  is a split monomorphism for every finite abelian group  $G$ . Since  $B\Gamma$  is a finite complex, the standard induction argument on the number of cells show that the group  $H_i(B\Gamma; \mathbf{L})$  is finitely generated for every  $i$ . Hence for every  $\alpha \in H_i(B\Gamma; \mathbf{L})$  there is  $p$  such that  $\alpha \otimes 1 \in H_i(B\Gamma; \mathbf{L}) \otimes \mathbf{Z}_{p^k}$  is not zero. By the UCF there is a



monomorphism  $H_i(\mathbf{B}\Gamma; \mathbf{L}) \otimes \mathbf{Z}_{p^k} \rightarrow H_i(\mathbf{B}\Gamma; \mathbf{L}(\mathbf{Z}) \wedge M(p^k))$  which, together with the assembly map, produces a commutative diagram

$$\begin{array}{ccc} H_i(\mathbf{B}\Gamma; \mathbf{L}) \otimes \mathbf{Z}_{p^k} & \longrightarrow & H_i(\mathbf{B}\Gamma; \mathbf{L} \wedge M(p^k)) \\ A \otimes 1 \downarrow & & A_p \downarrow \\ L_i(\mathbf{Z}\Gamma) \otimes \mathbf{Z}_{p^k} & \longrightarrow & \pi_i(\mathbf{L}(\mathbf{Z}\Gamma) \wedge M(p^k)). \end{array}$$

This diagram implies that  $A(\alpha) \neq 0$ .  $\square$

**Theorem 2.10.** *Suppose that  $\Gamma$  is a group with  $\mathbf{B}\Gamma$  a finite complex. If  $\mathbf{E}\Gamma$  has mod 2 acyclic Higson compactification, then the rational Novikov conjecture holds for  $\Gamma$ .*

*Proof.* In view of Remark 2.5 the argument of Theorem 2.7 for  $p = 2$  works without the assumption  $\dim \nu \mathbf{E}\Gamma < \infty$ . Hence the mod  $2^k$  assembly map

$$A \wedge 1_{M(2^k)} : \mathbf{H}_*(\mathbf{B}\Gamma; \mathbf{L}) \wedge M(2^k) \rightarrow \mathbf{L}_*(\mathbf{Z}\Gamma) \wedge M(2^k)$$

is a split monomorphism. On the group level by the Universal Coefficient Formula we have the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_i(\mathbf{B}\Gamma; \mathbf{L}) \otimes \mathbf{Z}_{2^k} & \longrightarrow & H_i(\mathbf{B}\Gamma; \mathbf{L} \wedge M(2^k)) & \longrightarrow & H_{i-1}(\mathbf{B}\Gamma; \mathbf{L}) * \mathbf{Z}_{2^k} \longrightarrow 0 \\ & & A_* \otimes 1 \downarrow & & (A \wedge 1)_* \downarrow & & (A_{*-1})_* \downarrow \\ 0 & \longrightarrow & \pi_i(\mathbf{L}_*(\mathbf{Z}\Gamma)) \otimes \mathbf{Z}_{2^k} & \longrightarrow & \pi_i(\mathbf{L}_*(\mathbf{Z}\Gamma) \wedge M(2^k)) & \longrightarrow & \pi_{i-1}(\mathbf{L}_*(\mathbf{Z}\Gamma)) * \mathbf{Z}_{2^k} \longrightarrow 0. \end{array}$$

Since the group  $H_i(\mathbf{B}\Gamma; \mathbf{L})$  is finitely generated, it can be presented as  $\bigoplus_{F_i} \mathbf{Z} \oplus \text{Tor}_i$ . Since  $A_* \otimes 1_{\mathbf{Z}_{2^k}}$  is a monomorphism, the kernel  $\ker(A_*)$  consists of  $2^k$  divisible elements. Since  $k$  is arbitrary,  $\ker(A_*|_{\bigoplus \mathbf{Z}}) = 0$ . Therefore,  $A_* \otimes 1_{\mathbf{Q}}$  is a monomorphism.  $\square$

This proves Theorem 1 of the introduction.

### 3 Mod $p$ acyclicity of Higson compactifications of asymptotically finite dimensional spaces

We recall that a map  $f : X \rightarrow Y$  between metric spaces is  $\lambda$ -Lipschitz if  $d_Y(f(x), f(x')) \leq \lambda d_X(x, x')$  for all  $x, x' \in X$ . Denote by

$$\text{Lip}(f) = \sup_{x \neq x'} \left\{ \frac{d_Y(f(x), f(x'))}{d_X(x, x')} \right\}$$

the minimal Lipschitz constant of  $f$ .

Every simplicial complex  $K$  carries a metric where all simplexes are isometric to the standard euclidean simplex. We will call the maximal such metric on  $K$  *uniform*

and usually we will denote the corresponding metric space by  $K_U$ .<sup>2</sup> Note that the metric space  $K_U$  is geodesic. If no metric is specified, we will assume that a finite complex is supplied with this uniform metric. In particular, the complexes in the two lemmas below are assumed to have the uniform metric.

The following lemma is a special case of Theorem A from [40].

**Lemma 3.1.** *Let  $Y$  be a finite simplicial complex with  $\pi_n(Y)$  finite. Then for every  $\lambda > 0$  there is a  $\mu > 0$  such that every map  $f : B^n \rightarrow Y$  with  $\text{Lip}(f|_{S^{n-1}}) \leq \lambda$  can be deformed to a  $\mu$ -Lipschitz map  $g : B^n \rightarrow Y$  by means of a homotopy  $h_t : B^n \rightarrow Y$  with  $h_t|_{S^{n-1}} = f|_{S^{n-1}}$ .*

**Lemma 3.2.** *Let  $L$  be a finite dimensional complex and let  $K$  be a finite complex with finite homotopy groups  $\pi_i(K)$  for  $i \leq \dim L + 1$ . Let  $f, g : L \rightarrow K$  be homotopic Lipschitz maps. Then every homotopy between  $f$  and  $g$  can be deformed to a Lipschitz homotopy  $H : L \times [0, 1] \rightarrow K$ .*

*Proof.* Let  $F : L \times I \rightarrow K$  be a homotopy between  $f$  and  $g$ . By induction on  $n$  using Lemma 3.1, we construct a  $\mu_n$ -Lipschitz map  $H_n : L^{(n)} \times I \cup L \times \{0, 1\} \rightarrow K$  which is a deformation of  $F$  restricted to the  $n$ -skeleton  $L^{(n)}$  such that  $H_n$  extends  $H_{n-1}$ . Here the fact that  $L$  (and hence  $L \times I$ ) is geodesic is essential for the argument because this condition guarantees that the union of  $\lambda$ -Lipschitz maps on  $\Delta^n \times I$  is  $\lambda$ -Lipschitz on  $L^{(n)} \times I$  (see [13] or [14] for details). Then  $H = H_m$  for  $m = \dim L$ .  $\square$

Let  $x_0 \in X$  be a basepoint in a metric space  $X$ . For  $x \in X$  and  $A \subset X$  we denote by  $\|x\| = \text{dist}(x, x_0)$  and  $\|A\| = \max\{\|z\| \mid z \in A\}$ . By  $B_r(x)$  we denote the closed  $r$ -ball in  $X$  centered at  $x$ . We use notation  $B_r = B_r(x_0)$ .

Let  $\sigma$  be an  $n$ -dimensional simplex spanned in a Euclidean space. By  $s_\sigma : \sigma \rightarrow \Delta^n$  we denote a simplicial homeomorphism onto the standard  $n$ -simplex  $\Delta^n = \{(x_i) \in \mathbb{R}^{n+1} \mid \sum x_i = 1, x_i \geq 0\}$ . A locally finite simplicial complex  $L$  with a geodesic metric on it is called an *asymptotic polyhedron* if every simplex in  $L$  is isometric to a simplex  $\sigma$  spanned in a Euclidean space and  $\lim_{\|\sigma\| \rightarrow \infty} \text{Lip}(s_\sigma) = 0$ . Asymptotic polyhedra usually do not have the uniform metric.

**Proposition 3.3.** *Let  $(K, d)$  be an asymptotic polyhedron and let  $d_U$  denote the uniform geodesic metric on  $K$ . Then the identity map  $u : (K, d) \rightarrow (K, d_U)$  satisfies the condition  $\lim_{x \rightarrow \infty} \text{diam}(u(B_r(x))) = 0$ .*

*Proof.* Let  $r$  and  $\varepsilon > 0$  be given. There is a  $t$  so that  $\text{Lip}(s_\sigma) < \varepsilon/(2r)$  for every  $\sigma$  with  $\|\sigma\| \geq t$ . Choose  $x$  so that  $\|x\| \geq t + 2r$ . For every two points  $z, z' \in B_r(x)$ , a geodesic segment  $J = [z, z']$  joining them does not intersect  $B_r$ . There is a partition of  $J$ :  $z = z_0 < z_1 \cdots < z_m = z'$  such that every segment  $J_i = [z_i, z_{i+1}]$

<sup>2</sup>An important exception to this convention occurs with *asymptotic polyhedra*, which are defined below.

lies in a simplex  $\sigma_i$ . By our choice of  $t$ ,  $\text{Lip}(s_{\sigma_i}) < \varepsilon/(2r)$  for each  $i$ . Therefore  $d_U(z_i, z_{i+1}) \leq \varepsilon/(2r)d(z_i, z_{i+1})$  and hence  $d_U(z, z') \leq \sum \varepsilon/(2r)d(z_i, z_{i+1}) = \varepsilon/(2r)d(z, z') \leq \varepsilon$ .  $\square$

A family of subsets  $\mathcal{A}$  is said to be  $d$ -disjoint if for every pair of sets  $A$  and  $A'$  in  $\mathcal{A}$  we have  $d(x, x') > d$  for all  $x \in A$  and  $x' \in A'$ . A family of subsets  $\mathcal{U}$  of a metric space  $X$  is *uniformly bounded* if there is a  $B > 0$  so that the diameter of each element of  $\mathcal{U}$  is less than  $B$ . Gromov, [23], defined the *asymptotic dimension*  $\text{asdim} X$  of a metric space  $X$  as follows:  $\text{asdim} X \leq n$  if for every  $d$  there are  $n + 1$   $d$ -disjoint uniformly bounded families  $\mathcal{U}_i$ ,  $i = 0, \dots, n$  of subsets of  $X$  such that the union  $\mathcal{U} = \cup \mathcal{U}_i$  is a cover of  $X$ .

Gromov's definition can be equivalently reformulated as follows:  $\text{asdim} X \leq n$  if for any arbitrary large number  $d$  there is a uniformly bounded open cover  $\mathcal{U}$  of  $X$  with multiplicity  $\leq n + 1$  and with Lebesgue number  $\geq d$  (see Assertion 1 in [BD2] for a proof).

We note that for every  $n$ -dimensional asymptotic polyhedron  $L$ ,  $\text{asdim} L \leq n$ .

Let  $\mathcal{U}$  be a locally finite open cover of a metric space  $X$ . The *canonical projection* to the nerve  $p : X \rightarrow \text{Nerve}(\mathcal{U})$  is defined using the partition of unity  $\{\phi_U : X \rightarrow \mathbb{R}\}_{U \in \mathcal{U}}$  defined by  $\phi_U(x) = d(x, X \setminus U) / \sum_{V \in \mathcal{U}} d(x, X \setminus V)$ . The family  $\{\phi_U : X \rightarrow \mathbb{R}\}_{U \in \mathcal{U}}$  defines a map  $p$  to the Hilbert space  $l_2(\mathcal{U})$  of square summable functions on  $\mathcal{U}$  with the Dirac functions  $\delta_U$ ,  $U \in \mathcal{U}$  as the basis. The nerve  $N(\mathcal{U})$  of the cover  $\mathcal{U}$  is realized in  $l_2(\mathcal{U})$  by taking every vertex  $U$  to  $\delta_U$ . Clearly, the image of  $p$  lies in the nerve. Given a family of positive numbers  $\bar{\lambda} = \{\lambda_U\}_{U \in \mathcal{U}}$  we can change the above imbedding of the nerve  $N(\mathcal{U})$  into  $l_2(\mathcal{U})$  by taking each vertex  $U$  to  $\lambda_U \delta_U$ . Then the projection  $p_{\bar{\lambda}}^{\mathcal{U}} : X \rightarrow N(\mathcal{U})$  to modified realization is given by the formula  $p_{\bar{\lambda}}^{\mathcal{U}}(x) = \sum_{U \in \mathcal{U}} \lambda_U \phi_U(x) \delta_U$ .

For a subset  $A \subset X$  we denote by  $L(\mathcal{U})|_A$  the Lebesgue number of  $\mathcal{U}$  restricted to  $A$ . More precisely,  $L(\mathcal{U})|_A = \inf_{y \in A} \max_{V \in \mathcal{U}} d(y, X \setminus V)$ .

**Proposition 3.4.** *Let  $\mathcal{U}$  be a locally finite cover of a geodesic metric space  $X$  with the multiplicity  $\leq m$ . Then the above projection to the nerve  $p_{\bar{\lambda}}^{\mathcal{U}} : X \rightarrow N = N(\mathcal{U})$  for  $\bar{\lambda} = \{\lambda_U\}$  with  $\lambda_U = L(\mathcal{U})|_U / (2m + 1)^2$  is 1-Lipschitz where the nerve is taken with the intrinsic metric induced from  $l_2(\mathcal{U})$ .*

*Proof.* We will show that the map  $\bar{p} = p_{\bar{\lambda}}^{\mathcal{U}}$  is 1-Lipschitz as a map to  $l_2(\mathcal{U})$ . Then the partition of  $N$  into simplices defines a locally finite partition on  $X$  such that  $\bar{p}$  is 1-Lipschitz on every piece, considered as the map to  $N$  with the induced path metric. Since  $X$  is a geodesic metric space, this will imply that  $\bar{p}$  is 1-Lipschitz.

Let  $x, y \in X$  and  $U \in \mathcal{U}$ . The triangle inequality implies

$$|d(x, X \setminus U) - d(y, X \setminus U)| \leq d(x, y).$$

It is easy to see that  $\sum_{V \in \mathcal{U}} d(x, X \setminus V) \geq L(\mathcal{U})|U| = (2m+1)^2 \lambda_U$  for  $x \in U$ . Then

$$\begin{aligned}
|\phi_U(x) - \phi_U(y)| &= \left| \frac{d(x, X \setminus U)}{\sum_{V \in \mathcal{U}} d(x, X \setminus V)} - \frac{d(y, X \setminus U)}{\sum_{V \in \mathcal{U}} d(x, X \setminus V)} \right. \\
&\quad \left. + \frac{d(y, X \setminus U)}{\sum_{V \in \mathcal{U}} d(x, X \setminus V)} - \frac{d(y, X \setminus U)}{\sum_{V \in \mathcal{U}} d(y, X \setminus V)} \right| \\
&\leq \frac{1}{\sum_{V \in \mathcal{U}} d(x, X \setminus V)} d(x, y) \\
&\quad + d(y, X \setminus U) \left| \frac{1}{\sum_{V \in \mathcal{U}} d(x, X \setminus V)} - \frac{1}{\sum_{V \in \mathcal{U}} d(y, X \setminus V)} \right| \\
&\leq \frac{1}{(2m+1)^2 \lambda_U} d(x, y) \\
&\quad + \frac{d(y, X \setminus U)}{\sum_{V \in \mathcal{U}} d(x, X \setminus V) \sum_{V \in \mathcal{U}} d(y, X \setminus V)} \sum_{V \in \mathcal{U}} |d(x, X \setminus V) - d(y, X \setminus V)| \\
&\leq \frac{1}{(2m+1)^2 \lambda_U} d(x, y) \\
&\quad + \frac{1}{(2m+1)^2 \lambda_U} \left( \sum_{V \in \mathcal{U}} |d(x, X \setminus V) - d(y, X \setminus V)| \right) \\
&\leq \frac{1}{(2m+1) \lambda_U} d(x, y) \leq \frac{1}{\sqrt{2m+1} \lambda_U} d(x, y)
\end{aligned}$$

Then

$$\begin{aligned}
\|p(x) - p(y)\| &= \left( \sum_{U \in \mathcal{U}} \lambda_U^2 (\phi_U(x) - \phi_U(y))^2 \right)^{\frac{1}{2}} \\
&\leq ((2m) \frac{1}{2m+1} d(x, y)^2)^{\frac{1}{2}} \leq d(x, y).
\end{aligned}$$

□

**Lemma 3.5.** *Suppose that  $X$  is a geodesic metric space with  $\text{asdim} X \leq n$  and let  $f : X \rightarrow \mathbb{R}_+$  be a proper function. Then there exist a compact set  $C \subset X$  and a map  $\phi : X \rightarrow N$  to a  $2n+1$ -dimensional asymptotic polyhedron with  $\text{diam}(\phi^{-1}(\Delta)) \leq f(z)$  for all  $z \in \phi^{-1}(\Delta) \setminus C$  and for all simplices  $\Delta$  of  $N$ . Moreover, every vertex of  $N$  lies in the image of  $\phi$ .*

*Proof.* We construct  $\phi$  as the projection  $p_{\mathcal{U}}^{\bar{\lambda}}$  from Proposition 3.4 to the nerve of a cover  $\mathcal{U}$  of  $X$ .

Fix a monotone sequence  $\{l_i\}$  tending to infinity. Since  $\text{asdim} X \leq n$ , for every  $i$  there is a uniformly bounded cover  $\mathcal{U}_i$  of multiplicity  $n+1$  with the Lebesgue number  $L(\mathcal{U}_i) > l_i$ . Let  $m_i$  be an upper bound for the diameter of elements of  $\mathcal{U}_i$ . Choose a sequence  $\{r_i\}$  such that  $f(X \setminus B_{r_i - m_i}) \geq 2m_{i+1}$  and take  $\mathcal{U} = \cup_i \mathcal{U}'_i$  where

$$\mathcal{U}'_i = \{U \setminus B_{r_{i-1}} \mid U \in \mathcal{U}_i, U \cap B_{r_i} \neq \emptyset\}.$$

We take  $C = B_{r_1}$  and check that the conclusion of our Lemma holds. By definition, the preimage  $\phi^{-1}(\Delta)$  lies in a union  $\cup_{j \in J} U'_j$  of sets from  $\mathcal{U}$  with nonempty intersection. Let  $x \in \cap_{j \in J} U'_j$ . Each  $U'_j$  belongs to the cover  $\mathcal{U}'_i$  for some  $i$ . Let  $k$  be maximal among those  $i$ 's for all  $j \in J$ . Then  $x \notin B_{r_{k-1}}$ . Therefore  $U'_j \subset X \setminus B_{r_{k-1} - m_{k-1}}$  for all  $j$ . Hence  $f(z) \geq 2m_k \geq \text{diam} \cup_j U'_j$  for  $z \in X \setminus B_{r_{k-1} - m_{k-1}}$  and  $k > 1$ .

Since  $l_i \rightarrow \infty$  the nerve  $N$  realized in  $l_2(\mathcal{U})$  as above with the intrinsic metric is an asymptotic polyhedron. Since the covers  $\mathcal{U}'_i$  and  $\mathcal{U}'_j$  do not intersect for  $|i-j| > 1$ , the multiplicity of  $\mathcal{U}$  is  $\leq 2n+2$ .

We can ensure that  $\phi$  is onto the vertices by the following trick: omit from the cover any elements where the Lebesgue number  $L(x)$  is not actually attained. Then choose an element  $x_U \in U$  for each  $U \in \mathcal{U}$  where the Lebesgue number is attained. Subtract a ball of radius  $L(x_U)/4$  around  $x_U$  from each of the other elements of the cover. The result is an irreducible cover with Lebesgue number going to infinity and  $\phi$  is onto.  $\square$

*Remark 3.6.* Using a standard dimension theoretic trick the polyhedron  $N$  can be chosen to be  $n$ -dimensional.

A metric space  $X$  is called *uniformly contractible* if there is a function  $S: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  such that every ball  $B_r(x)$  is contractible to a point in the ball  $B_{S(r)}(x)$ . The function  $S$  is called a *contractibility function* for  $X$ . We will always take our contractibility functions to be strictly monotone.

**Proposition 3.7.** *Let  $X$  be a uniformly contractible proper metric space with  $\text{asdim} X \leq n$ . Then given a proper function  $g: X \rightarrow \mathbf{R}_+$  there exist a  $2n+1$ -dimensional asymptotic polyhedron  $N$ , a proper 1-Lipschitz map  $\phi: X \rightarrow N$ , and a proper homotopy inverse map  $\gamma: N \rightarrow X$  with  $d(x, \gamma\phi(x)) < g(x)$  for all  $x \in X$ . Moreover, there is a compact set  $C \subset X$  such that  $\text{diam}(\phi^{-1}(\Delta)) \leq g(x)$  for all  $x \in \Delta \setminus C$ .*

*Proof.* Let  $S$  be a contractibility function. We define  $\rho(t) = S^{-1}(t/2)$  where  $S^{-1}$  is the inverse function for  $S$ . Then we take  $f = \rho^{2n+1} \circ g$ , the composition of  $g$  and the  $2n+1$ -fold iteration of  $\rho$ . Clearly  $g \leq f$ . We assume here that  $g(x) \leq \|x\|/2$ . We define by induction on  $i$  a lift  $\gamma$  on the  $i$ -skeleton  $N^{(i)}$  of the nerve of a cover of  $X$  given by Lemma 3.5 for this choice of  $f$ . We take  $\gamma(v) \in \phi^{-1}(v)$  for every vertex  $v$ . Then using the uniform contractibility of  $X$  we can extend  $\gamma$  with control over the 1-skeleton  $N^{(1)}$  and so on. Without loss of generality (and for simplicity

of exposition) we may assume that  $X$  is a polyhedron of the dimension  $n$  supplied with a triangulation of mesh  $\leq 1$ . By induction on  $i$  we define a homotopy  $H : X^{(i)} \times I \rightarrow X$  joining the identity map with  $\gamma \circ \phi$ . We consider a function  $\psi(x) = \|x\| - \max\{d(x, y) \mid y \in H(x \times I)\}$ . If  $\psi$  tends to infinity, then the map  $H$  is proper, but it is easy to verify that  $\psi(x) \geq \|x\|/2$ .  $\square$

We recall that the Higson compactification  $\bar{X}$  of a proper metric space  $X$  can be defined as the maximal ideal space of the completion of the ring of bounded functions with the gradient tending to zero at infinity [37]. The defining property of the Higson corona is the following:

(\*A continuous map  $f : X \rightarrow Z$  to a compact metric space is extendable to the Higson corona  $\nu X$  if and only if it satisfies the condition: For arbitrary large  $R$

$$\lim_{\|x\| \rightarrow \infty} \text{diam}(f(B_R(x))) = 0.$$

Note that a proper Lipschitz map  $f : X \rightarrow Y$  induces a continuous mapping between the Higson coronas  $\bar{f} : \nu X \rightarrow \nu Y$ .

**Theorem 3.8.** *Let  $X$  be a uniformly contractible geodesic proper metric space with finite asymptotic dimension and let  $\bar{X}$  be the Higson compactification. Then  $\check{H}^n(\bar{X}; \mathbf{Z}_p) = 0$  for all  $n > 0$  and all  $p$ .*

*Proof.* We show that every map  $\alpha : \bar{X} \rightarrow K(\mathbf{Z}_p, n)$  is null homotopic. Since  $\bar{X}$  is compact, the image  $\alpha(\bar{X})$  is contained in the  $k$ -skeleton  $K = K(\mathbf{Z}_p, n)^{(k)}$  for some  $k$ . We will assume that  $K$  is a finite complex of dimension at least  $2n + 2$ . For convenience, we replace  $K$  by a Riemannian manifold with Riemannianly collared boundary. Let  $4\varepsilon_K$  be a convexity radius for  $K$ , i.e. a positive constant such that every two points in an  $4\varepsilon_K$ -ball can be joined by a unique geodesic in that  $4\varepsilon_K$ -ball.

Since the map  $\alpha|_X \rightarrow K$  is extendable over the Higson corona, the constructions above allow us to replace  $\alpha|_X$  by a function, also called  $\alpha$  by abuse of notation, such that the function  $R_\alpha(t) = \text{Lip}(\alpha|_{X \setminus B_t(x_0)})$  tends to zero at infinity. We apply Lemma 3.7 with

$$g(x) \leq \min\{\varepsilon_K/R_\alpha(\|x\|/2), \|x\|/4\}$$

to obtain an asymptotic polyhedron  $N$  and maps  $\phi : X \rightarrow N$  and  $\gamma : N \rightarrow X$ .

We may assume that  $\phi$  is surjective on vertices. Let  $[u, v]$  be an edge in  $N$ , then

$$d_K(\alpha\gamma(u), \alpha\gamma(v)) \leq R_\alpha(t_0)d_X(\gamma(u), \gamma(v))$$

where  $t_0 = \min\{\|\gamma(u)\|, \|\gamma(v)\|\}$ . We may assume that there are  $x, y \in X$  such that  $\phi(x) = u$  and  $\phi(y) = v$ . Then

$$\begin{aligned} d_X(\gamma(u), \gamma(v)) &\leq d_X(\gamma(\phi(x)), \gamma(\phi(y))) \\ &\leq d_X(\gamma(\phi(x)), x) + d_X(x, y) + d_X(y, \gamma(\phi(y))) \\ &\leq g(x) + \text{diam } \phi^{-1}[x, y] + g(y) \leq 2g(x) + g(y) \leq 3g(x) \\ &\leq 3\varepsilon_K/R_\alpha(\|x\|/2) \end{aligned}$$

provided that  $\|x\| \geq \|y\|$ .

As a result, we obtain the inequality  $d_K(\alpha\gamma(u), \alpha\gamma(v)) \leq 3\varepsilon_K$ . This means that there is a map  $\beta : N \rightarrow K$  which coincides on vertices with the map  $\alpha \circ \gamma$  and which is  $3\varepsilon_K$ -Lipschitz for  $N_U = N$  taken with the uniform metric:  $\beta$  is obtained by linear extension of the restriction of  $\alpha \circ \gamma$  to vertices. Note that  $\beta$  is  $\varepsilon_K$ -close to  $\alpha \circ \gamma$  and that these maps are therefore homotopic.

Since  $X$  is contractible, the map  $\alpha \circ \gamma$  is null homotopic. Therefore, so is  $\beta$ . Note that the homotopy groups  $\pi_i(K)$  are finite for  $i \leq \dim N + 1$ . We apply Lemma 3.2 to obtain a  $\lambda$ -Lipschitz homotopy  $H : N_U \times I \rightarrow K$  of  $\beta$  to a constant map. This homotopy defines a Lipschitz map  $\tilde{H} : N_U \rightarrow K_\lambda^I$  to the space of  $\lambda$ -Lipschitz mappings of the unit interval  $I$  to  $K$ . We note that the space  $K_\lambda^I$  is compact. Then, by Proposition 3.3,  $\tilde{H} \circ u : N \rightarrow K_\lambda^I$  satisfies the Higson extendibility condition (\*). Let  $\tilde{h} : \bar{N} \rightarrow K_\lambda^I$  be the extension over the Higson corona. This extension defines a map  $\bar{H} : \bar{N} \times I \rightarrow K$ . The map  $\bar{H}$  is a homotopy between the extension  $\bar{\beta}$  and a constant map. To complete the proof, we show that  $\alpha$  is homotopic to  $\bar{\beta} \circ \bar{\phi}$  where  $\bar{\phi}$  is the extension of the Lipschitz map  $\phi$  to the Higson compactifications. Note that

$$d_K(\alpha(x), \alpha\gamma\phi(x)) \leq R_\alpha(t_0)d(x, \gamma\phi(x)) \leq R_\alpha(t_0)\varepsilon_K/R_\alpha(\|x\|/2) \leq \varepsilon_K$$

where  $t_0 = \min\{\|x\|, \|\gamma\phi(x)\|\} \geq \|x\|/2$ . Therefore,  $d_K(\alpha(x), \beta \circ \phi(x)) \leq 4\varepsilon_K$ . Then for every  $x \in X$  we join the points  $\alpha(x)$  and  $\beta\phi(x)$  by the unique geodesic  $\psi_x : I \rightarrow K$ . This defines a map  $\tilde{\psi} : X \rightarrow K_\mu^I$ . Since both  $\alpha$  and  $\beta \circ \phi$  satisfy the condition (\*), the map  $\tilde{\psi}$  has the property (\*). Let  $\bar{\psi} : \bar{X} \rightarrow K_\mu^I$  be the extension of  $\tilde{\psi}$  to the Higson corona. The map  $\bar{\psi}$  defines a homotopy  $\Psi : \bar{X} \times I \rightarrow K$  between  $\alpha$  and  $\bar{\beta} \circ \bar{\phi}$ .  $\square$

We have now proven Theorem 3. The following asserts Theorem 2:

**Corollary 3.9.** *Suppose that a group  $\Gamma$  has finite asymptotic dimension and that  $B\Gamma$  is a finite complex. Then the integral assembly map  $A : H_*(B\Gamma; \mathbf{L}) \rightarrow L_*(\mathbf{Z}\Gamma)$  is a monomorphism.*

*Proof.* In view of the inequality  $\dim v\Gamma \leq \text{asdim}\Gamma$  [18] we can apply Theorem 2.9.  $\square$

**Corollary 3.10.** *If  $\Gamma$  and  $R$  satisfy the finiteness conditions of Theorem 2, then  $H_i(B\Gamma; \mathbf{L}(R)) \rightarrow L_i(R\Gamma)$  is injective for all  $i$ .*

### Acknowledgment.

This paper was partially supported by NSF grants. The first author would like to thank the Max-Planck Institute für Mathematik in Bonn for hospitality and for presenting the preliminary version of this paper as an MPI preprint. The second author would like to thank the University of Chicago for support during several visits. The third author would like to thank the Courant Insitute for its support during the time that this paper was being written.

### Bibliography

- [1] Bartels, A. Squeezing and higher algebraic  $K$ -theory *K-Theory* **28** (2003), no. 1, 19–37.
- [2] Bell, G. and Dranishnikov, A. On asymptotic dimension of groups *Algebr. Geom. Topol.* **1** (2001), 57–71.
- [3] Bell, G. and Dranishnikov, A. On asymptotic dimension of groups acting on trees *Geom. Dedicata* **103** (2004), 89–101.
- [4] Block, J. and Weinberger, S. Aperiodic tilings, positive scalar curvature and amenability of spaces *J. Amer. Math. Soc.* **5** (1992), no. 4, 907–918.
- [5] Bökstedt, M., Hsiang, W. C., and Madsen, I. The cyclotomic trace and the  $K$ -theoretic analogue of Novikov’s conjecture *Proc. Nat. Acad. Sci. U.S.A.* **86** (1989), no. 22, 8607–8609.
- [6] Carlsson, G. Bounded  $K$ -theory and the assembly map in algebraic  $K$ -theory. *Novikov conjectures, index theorems and rigidity, Vol. 2 (Oberwolfach, 1993)*, 5–127. London Math. Soc. Lecture Note Ser., 227, Cambridge Univ. Press, Cambridge, 1995.
- [7] Carlsson, G. and Goldfarb, B. On homological coherence of discrete groups *J. Algebra* **276** (2004), no. 2, 502–514.
- [8] Carlsson, G., and Goldfarb, B. The integral  $K$ -theoretic Novikov conjecture for groups with finite asymptotic dimension *Invent. Math.* **157** (2004), no. 2, 405–418.
- [9] Carlsson, G., and Pedersen, E. Controlled algebra and the Novikov conjectures for  $K$ - and  $L$ -theory *Topology* **34** (1995), no. 3, 731–758.
- [10] Carlsson, G., and Pedersen, E. Čech homology and the Novikov conjectures for  $K$ - and  $L$ -theory. *Math. Scand.* **82** (1998), no. 1, 5–47.
- [11] Chang, S., Ferry, S., and Yu, G. Bounded rigidity of manifolds and asymptotic dimension growth *K-Theory (to appear)*.
- [12] Dranishnikov, A. Asymptotic topology (*Russian*) *Uspekhi Mat. Nauk* **55** (2000), no. 6(336), 71–116; *translation in Russian Math. Surveys* **55** (2000), no. 6, 1085–1129.
- [13] Dranishnikov, A. On hypersphericity of manifolds with finite asymptotic dimension *Trans. Amer. Math. Soc.* **355** (2003), no. 1, 155–167.
- [14] Dranishnikov, A. *Geom. Dedicata* **117** (2006), 215–231.
- [15] Dranishnikov, A. N. and Ferry, S. On the Higson-Roe corona (*Russian*) *Uspekhi Mat. Nauk* **52** (1997), no. 5(317), 133–146 *translation in Russian Math. Surveys* **52** (1997), no. 5, 1017–1028
- [16] Large Riemannian manifolds which are flexible Dranishnikov, A., Ferry, S. and Weinberger S., *Ann. of Math. (2)* **157** (2003), no. 3, 919–938.



- [17] Dranishnikov, A. and Januszkiewicz, T., Every Coxeter group acts amenably on a compact space *Proceedings of the 1999 Topology and Dynamics Conference (Salt Lake City, UT)*, *Topology Proc.* **24**, (1999), 135–141.
- [18] Dranishnikov, A. N. , Keesling, J. , and Uspenskij, V. V., On the Higson corona of uniformly contractible spaces *Topology* **37** (1998), no. 4, 791–803.
- [19] Edwards, David A. ; Hastings, Harold M. *Čech and Steenrod homotopy theories with applications to geometric topology* Lecture Notes in Mathematics, Vol. 542. Springer-Verlag, Berlin-New York, 1976.
- [20] Farrell, F. T. and Hsiang, W. C. On Novikov’s conjecture for nonpositively curved manifolds. I. *Ann. of Math. (2)* **113** (1981), no. 1, 199–209.
- [21] Ferry, S., Ranicki, A., and Rosenberg, J, A history and survey of the Novikov conjecture, *Novikov conjectures, index theorems and rigidity, Vol. 1 (Oberwolfach, 1993)*, London Math. Soc. Lecture Note Ser., **226**, 7–66, Cambridge Univ. Press, Cambridge, 1995.
- [22] Ferry, S., and Weinberger, S., A coarse approach to the Novikov conjecture *Novikov conjectures, index theorems and rigidity, Vol. 1 (Oberwolfach, 1993)*, 147–163, London Math. Soc. Lecture Note Ser., **226**, Cambridge Univ. Press, Cambridge, 1995.
- [23] Gromov, M. *Asymptotic invariants of infinite groups* Geometric group theory, Vol. 2 (Sussex, 1991), 1–295, London Math. Soc. Lecture Note Ser., 182, Cambridge Univ. Press, Cambridge, 1993.
- [24] Gromov, M. Spaces and questions. GAFA 2000 (Tel Aviv, 1999). *Geom. Funct. Anal.* (2000), Special Volume, Part I, 118–161.
- [25] Higson, N., Lafforgue, V. and Skandalis, G., Counterexamples to the Baum-Connes conjecture, *Geom. Funct. Anal.*, **12**, (2002), no. 2, 330–354.
- [26] Higson, N., and Roe, J. *Analytic K-homology* Oxford Mathematical Monographs. Oxford Science Publications. Oxford University Press, Oxford, 2000.
- [27] Ji, L., Asymptotic dimension and the integral  $K$ -theoretic Novikov conjecture for arithmetic groups, *J. Differential Geom.*, **68**, (2004), no. 3, 535–544.
- [28] Kaminker, J., and Schochet, C.,  $K$ -theory and Steenrod homology: applications to the Brown-Douglas-Fillmore theory of operator algebras. *Trans. Amer. Math. Soc.* **227** (1977), 63–107.
- [29] Keesling, J. The one-dimensional Čech cohomology of the Higson compactification and its corona. *Topology Proc.* **19** (1994), 129–148.
- [30] Massey, William S. *Homology and cohomology theory. An approach based on Alexander-Spanier cochains* Monographs and Textbooks in Pure and Applied Mathematics, Vol. 46. Marcel Dekker, Inc., New York-Basel, 1978.
- [31] Madsen, I. and Milgram, R. J. The universal smooth surgery class. *Comment. Math. Helv.* **50** (1975), no. 3, 281–310.
- [32] Ozawa, N. Amenable actions and exactness for discrete groups, *C. R. Acad. Sci. Paris Sér. I Math.*, **330**, (2000), no. 8, 691–695.
- [33] Pedersen, E., Bounded and continuous control, *Novikov conjectures, index theorems and rigidity, Vol. 2 (Oberwolfach, 1993)*, London Math. Soc. Lecture Note Ser., **227**, 277–284, Cambridge Univ. Press, Cambridge, 1995.
- [34] Ranicki, A., *Algebraic L-theory and topological manifolds*, Cambridge Tracts in Mathematics, **102**, Cambridge University Press, Cambridge, 1992.
- [35] Ranicki, A., Additive  $L$ -theory,  $K$ -Theory, **3**, (1989), no. 2, 163–195.
- [36] Ranicki, A. On the Novikov conjecture *Novikov conjectures, index theorems and rigidity, Vol. 1 (Oberwolfach, 1993)*, 272–337, London Math. Soc. Lecture Note Ser., 226, Cambridge Univ. Press, Cambridge, 1995.
- [37] Roe, J. Coarse cohomology and index theory on complete Riemannian manifolds. *Mem. Amer. Math. Soc.* **104** (1993), no. 497.

- [38] Roe, J. *Lectures on coarse geometry* University Lecture Series, 31. American Mathematical Society, Providence, RI, 2003.
- [39] Rosenberg, J. Analytic Novikov for topologists, *Novikov conjectures, index theorems and rigidity, Vol. I (Oberwolfach, 1993)*, 338–372, London Math. Soc. Lecture Note Ser., 226, Cambridge Univ. Press, Cambridge, 1995.
- [40] Siegel, J. and Williams, F., Uniform bounds for isoperimetric problems *Proc. Amer. Math. Soc.* **107** (1989), no. 2, 459–464.
- [41] Skandalis, G., Tu, J. L., and Yu, G., The coarse Baum-Connes conjecture and groupoids *Topology* **41** (2002), no. 4, 807–834.
- [42] Taylor, J., A counterexample in shape theory, *Bull. Amer. Math. Soc.*, **81**, (1975), 629–632.
- [43] Taylor, L., Williams, B., Surgery spaces: formulae and structure. *Algebraic topology, Waterloo, 1978 (Proc. Conf., Univ. Waterloo, Waterloo, Ont., 1978)*, pp. 170–195, Lecture Notes in Math., **741**, Springer, Berlin, 1979.
- [44] Yu, G., The Novikov conjecture for groups with finite asymptotic dimension. *Ann. of Math. (2)* **147** (1998), no. 2, 325–355.

Received Month 200X.