

Handout 3

- **The Cantor set.** Let us start with the interval $C = [0, 1]$ and remove the middle third open interval $(\frac{1}{3}, \frac{2}{3})$. This leaves us with the set $C_1 = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$ formed of 2 closed subintervals. Having constructed $C_1 \supset C_2 \supset \dots \supset C_n$ where C_n is the union of 2^n subintervals each of length $\frac{1}{3^n}$, we construct C_{n+1} as follows: To obtain C_{n+1} we remove the middle third of each of the 2^n intervals that form C_n . This leaves us with a union of 2^{n+1} intervals each of length $\frac{1}{3^{n+1}}$.

Q1) Let $C = \bigcap_{n=1}^{\infty} C_n$. Why is C non-empty? Is it compact?

Q2) Show that every point in C is a limit point. Hence C is a perfect set.

Conclusion: From the homework (HW 2), we deduce that C is uncountable, since any perfect subset of \mathbb{R}^d is uncountable.

Q3) Show that C cannot contain any interval (a, b) .

Conclusion: As such, C is totally disconnected (it has no non-trivial connected subset) and nowhere dense (the interior of its closure is empty).

Q4) What is the total length of C_n ? What would be a reasonable definition of the length of C ?

-
- **Wish list for a measure function** Motivated by the above, it would be grand to have a measure function that tells us how big or small a subset of \mathbb{R}^d is. This would be a function from the set of subsets of \mathbb{R}^d into $[0, \infty]$, say $m : \mathcal{P}(\mathbb{R}^d) \rightarrow [0, \infty]$. We would like this function to satisfy the following properties:

- a) If E_1, E_2, \dots is a countable collection of disjoint subsets of \mathbb{R} , then

$$m(\cup_{n=1}^{\infty} E_n) = \sum_{n=1}^{\infty} m(E_n).$$

- b) If E is congruent to F (i.e. F can be obtained from E by applying rigid motions: translations, rotations, or a reflections) then we should have that $m(E) = m(F)$.
- c) $m([0, 1)^d) = 1$.

The bad news is that no such function can exist, and here's why (at least when $d = 1$). Let us define an equivalence relation between elements of $[0, 1)$ as follows: We say $x \sim y$ if $x - y$ is a rational number. Let N be the subset of $[0, 1]$ that contains exactly one element of each equivalence relation (the existence of this N requires invoking the axiom of choice). Now let $R = [0, 1) \cap \mathbb{Q}$, and for each $r \in R$ define the set

$$N_r = \{x + r : x \in N \cap [0, 1 - r]\} \cup \{x + r - 1 : x \in N \cap [1 - r, 1)\}.$$

(Basically N_r is just the translate of N by r units to the right, except that we move the part that sticks out of the interval $[0, 1)$ one unit to the left).

- Q5)** Show that $[0, 1)$ is the disjoint union of N_r for $r \in R$.
- Q6)** Show that if a measure function satisfying a), b) and c) above exists, then $m(N) = m(N_r)$ for every $r \in R$.
- Q7)** Arrive at a contradiction.

Remark: One might think that possibly relaxing condition a) to cover only *finitely* many disjoint sets E_n , i.e.

$$m(\cup_{n=1}^N E_n) = \sum_{n=1}^N m(E_n).$$

would resolve the contradiction. Unfortunately, the Banach-Tarski paradox (cf. Figure 1) tells us that this is not enough to resolve this issue.



Figure 1: Banach-Tarski tells us that we can split the unit ball in \mathbb{R}^3 into finitely many (actually 5 is sufficient) many disjoint pieces, apply rigid motions to those pieces and then reassemble them to obtain two copies of the unit ball.

Conclusion: The problem with the above wishlist is that we insisted on being able to measure *every* subset of \mathbb{R}^d . We have shown that this is impossible. The solution is to be content with a measure function that is defined on some but not all subsets. Such subsets will be called measurable subsets.

MATH 395 Notes

Faye Jackson

September 18, 2020

Proof of Q1. For notational convenience denote for $n \in \mathbb{N}_0$:

$$C_n = \bigcup_{i=1}^{2^n} [a_i^n, b_i^n]$$

So that inductively for $1 \leq i \leq 2^n$:

$$\begin{aligned} C_0 &= [0, 1] \\ [a_{2i-1}^{n+1}, b_{2i-1}^{n+1}] &= \left[a_i^n, \frac{2a_i^n + b_i^n}{3} \right] \\ [a_{2i}^{n+1}, b_{2i}^{n+1}] &= \left[\frac{a_i^n + 2b_i^n}{3}, b_i^n \right] \end{aligned}$$

Now lets tackle both of these questions!

- Note that $a_1^0 = 0$ will always lie at the edge of an interval because supposing $a_1^n = 0$ we know $a_1^{n+1} = a_{2 \cdot 1 - 1}^{n+1} = a_1^n = 0$. Therefore since:

$$0 \in [a_1^n, b_1^n] \subseteq C_n$$

for each $n \geq 0$ we must know that $0 \in C$. A similar argument shows that $1 \in C$.

- C is compact!!! Why? Note that for every $n \geq 0$ we have that C_n is a finite union of closed intervals, so each C_n is closed. Thus, $C = \bigcap_{n=0}^{\infty} C_n$ is closed. Furthermore since $C_0 = [0, 1]$ is closed and bounded, that is compact. Therefore since $C \subseteq C_0$ is a closed subset of a compact set, C must be compact.

Perfect! We win!



Proof of Q2. Fix some point $x \in C$. Then $x \in C_n$ for all $n \geq 0$, and so for each $n \geq 0$ there exists some $1 \leq i_n \leq 2^n$ so that $x \in [a_{i_n}^n, b_{i_n}^n]$. Suppose that $\varepsilon > 0$, then there is some $N \in \mathbb{N}$ so that $\frac{1}{3^N} < \varepsilon$. We claim that $a_{i_N}^N, b_{i_N}^N \in N_\varepsilon(x) \cap C$

- First we show that for all $n \geq 0$ and all $1 \leq i \leq 2^n$ we have a_i^n is in C . First note that $a_i^n \in [a_i^n, b_i^n] \subseteq C_n$, and thus for each $0 \leq m < n$ we must have $a_i^n \in C_n \subseteq C_m$. Inductively we will show that for $m \geq n$ if we let $j_n = i$ and $j_{m+1} = 2j_m - 1$ then:

$$a_i^n = a_{j_m}^m \in C_m$$

Note that it's trivial for $m = n$. Now suppose that $a_{j_m}^m = a_i^n$. Consider that:

$$a_{j_{m+1}}^{m+1} = a_{2j_m-1}^{m+1} = a_{j_m}^m = a_i^n$$

And so we must have that this works! Great.

- Now we show that for all $n \geq 0$ and all $1 \leq i \leq 2^n$ we have b_i^n is in C . First note that $b_i^n \in [a_i^n, b_i^n] \subseteq C_n$, and thus for each $0 \leq m < n$ we must have $b_i^n \in C_n \subseteq C_m$. Inductively we will show that for $m \geq n$ if we let $j_n = i$ and $j_{m+1} = 2j_m$ then:

$$b_i^n = b_{j_m}^m \in C_m$$

Note that it's trivial for $m = n$. Now suppose that $b_{j_m}^m = b_i^n$. Consider that:

$$b_{j_{m+1}}^{m+1} = b_{2j_m}^{m+1} = b_{j_m}^m = b_i^n$$


And so we must have that this works! Great.


- Now we show that for each $n \geq 0$ and each $1 \leq i \leq 2^n$ the interval $[a_i^n, b_i^n]$ has length $\frac{1}{3^n}$. Note first that:

$$b_1^0 - a_1^0 = 1 - 0 = 1 = \frac{1}{3^0}$$

Inductively for $1 \leq i \leq 2^n$ then we know that:

$$\begin{aligned} b_{2^{n-i}}^{n+1} - a_{2^{n-i}}^{n+1} &= \frac{2a_i^n + b_i^n}{3} - a_i^n = \frac{b_i^n - a_i^n}{3} = \frac{1}{3} \cdot \frac{1}{3^n} = \frac{1}{3^{n+1}} \\ b_{2^n}^{n+1} - a_{2^n}^{n+1} &= b_i^n - \frac{a_i^n + 2b_i^n}{3} = \frac{b_i^n - a_i^n}{3} = \frac{1}{3} \cdot \frac{1}{3^n} = \frac{1}{3^{n+1}} \end{aligned}$$

Now we're done, since in particular $a_{i_N}^N$ and $b_{i_N}^N$ are distinct, so for any ε neighborhood of x there are at least two points in $N_\varepsilon(x) \cap C$. Thus x is a limit point. 


Proof of Q3. Fix $a < b$. But then if we had two points $x, y \in (a, b)$ such that $x, y \in C$ and $y > x$. Note that we then know that there exists some $N \in \mathbb{N}$ so that $\frac{1}{3^N} < \varepsilon$. This means that x and y must lie in different intervals making up C_N , since these are disjoint. But then $(a, b) \cap C$ is not an interval, since $x, y \in C \cap (a, b)$ but there is some point z between x and y so that $z \notin C$. This necessarily means so then $(a, b) \neq C \cap (a, b)$, and so $(a, b) \not\subseteq C$. 

Proof of Q4. Note that the total length of C^n is:

$$\ell(C_n) = \frac{2^n}{3^n}$$

Since C_n is a union of 2^n disjoint intervals each of length 3^n . Note that for each $n \in \mathbb{N}$ we must conclude since $C \subseteq C_n$ we know:

$$\ell(C) \leq \ell(C_n) = \frac{2^n}{3^n} = \left(\frac{2}{3}\right)^n$$

Taking $n \rightarrow \infty$ we then can see that $\ell(C)$ should be zero. 

Proof of Q5. Fix $r, q \in R = [0, 1) \cap \mathbb{Q}$. We will first show that if $N_r \cap N_q \neq \emptyset$ then $r = q$, so by contrapositive the $\{N_r\}_{r \in R}$ are disjoint. Fix $y \in N_r \cap N_q$. There are four cases:

- $y = x_r + r$ and $y = x_q + q$ for some $x_r, x_q \in N$. Then $x_r - x_q = q - r$ by algebra, and so since $r, q \in \mathbb{Q}$ we have that $q - r \in \mathbb{Q}$ and so $x_r \sim x_q$. By the definition of N it follows that $x_r = x_q$. Therefore $x_r + r = x_r + q$, giving that $r = q$.
- $y = x_r + r - 1$ and $y = x_q + q - 1$ for some $x_r, x_q \in N$. Then $x_r - x_q = q - r$ by algebra, and so since $r, q \in \mathbb{Q}$ we have that $q - r \in \mathbb{Q}$ and so $x_r \sim x_q$. By

the definition of N it follows that $x_r = x_q$. Therefore $x_r + r - 1 = x_r + q - 1$, giving that $r = q$.

- $y = x_r + r - 1$ and $y = x_q + q$ for some $x_r, x_q \in N$. Then $x_r - x_q = q - r + 1 \in \mathbb{Q}$. Thus $x_r = x_q$. Therefore $q = r - 1$ by some quick algebra. This is clearly a contradiction! Why? Well $0 \leq r < 1$, and so $-1 \leq r - 1 < 0$, but we know $q \geq 0$!!! Oops!
- $y = x_r + r$ and $y = x_q + q - 1$ for some $x_r, x_q \in N$. Then $x_r - x_q = q - r - 1 \in \mathbb{Q}$. Thus $x_r = x_q$. Therefore $r = q - 1$ by some quick algebra. This is clearly a contradiction! Why? Well $0 \leq q < 1$, and so $-1 \leq q - 1 < 0$, but we know $r \geq 0$!!! Oops!

We want to show that:

$$[0, 1) = \bigcup_{r \in R} N_r$$

Let's go!

(\subseteq) Fix $y \in [0, 1)$. Then by definition there is some $x \in N$ so that $y \sim x$. Note that then $y - x \in \mathbb{Q}$. Further we have $0 \leq x, y < 1$ There are two cases:

- Suppose that $y - x \geq 0$. Now set $r := y - x$. First note that since $x \geq 0$ and $y < 1$ we know $y - x < 1 - 0 = 1$. Therefore $r \in \mathbb{Q} \cap [0, 1) = R$. We claim that $y \in N_r$. In particular note that $y = x + r$. All that remains to be shown is $x \in [0, 1 - r)$. We know since $x \in N$ that $x \in [0, 1)$, so $x \geq 0$ immediately. We merely need to show that $x < 1 - y + x$. This is simple, since $y < 1$ we know $1 - y > 0$. With this we must have that $x \in [0, 1) \cap N$, and so:

$$y \in \{x' + r \mid x' \in N \cap [0, 1 - r)\} \subseteq N_r$$

And so $y \in N_r$

- Suppose that $y - x < 0$. Set $r := y - x + 1$. Note then that $r < 1$. Since $0 \leq y$ we know $-x \leq y - x$, and then since $x < 1$ it follows that $-1 < -x \leq y - x$, and so $0 < r$. This shows since $r \in \mathbb{Q}$ that $r \in R = [0, 1) \cap \mathbb{Q}$. We claim that $y \in N_r$. Note in particular that $y = x + r - 1$ by algebra. We need merely show that $x \in [1 - r, 1)$. To

do this note that $y \geq 0$ so $y \leq 0$:

$$x \geq -y + x = 1 - y + x - 1 = 1 - (y - x + 1) = 1 - r$$

And we already know $x < 1$. Therefore:

$$y \in \{x' + r - 1 \mid x' \in N \cap [1 - r, 1)\} \subseteq N_r$$

And so $y \in N_r$!

Great! Since in either case $r \in R$, we must have that $y \in \bigcup_{r \in R} N_r$. This finishes this direction!

(\supseteq) This side follows fairly immediately. Fix $y \in \bigcup_{r \in R} N_r$. Then $y \in N_r$ for some $r \in R$. There are then two quick cases:

- We have that $y = x + r$ for some $x \in N \cap [0, 1 - r)$. Then note that since $r \geq 0$ we have:

$$\begin{aligned} 0 &\leq x < 1 - r \\ 0 &\leq r \leq x + r = y < 1 \end{aligned}$$

And thus $y \in [0, 1)$

- We have that $y = x + r - 1$ for some $x \in N \cap [1 - r, 1)$. Then note that since $1 - r \leq x < 1$ that $-r \leq x - 1 < 0$. Therefore since $r < 1$ we know:

$$0 \leq x + r - 1 < r < 1$$

With this we're done!

We've finished the proof that this is a disjoint union! Wow!



Proof of Q6. Fix some $r \in R$. We wish to show that $m(N_r) = m(N)$. First note that:

$$\begin{aligned} m(N_r) &= m(\{x + r \mid x \in N \cap [0, 1 - r)\} \cup \{x + r - 1 \mid x \in N \cap [1 - r, 1)\}) \\ &= m(\{x + r \mid x \in N \cap [0, 1 - r)\}) + m(\{x + r - 1 \mid x \in N \cap [1 - r, 1)\}) \end{aligned}$$

This follows from axiom (a) for our measure. But then by axiom (b) note that these are translations of $N \cap [0, 1 - r)$ and $N \cap [1 - r, 1)$ respectively so:

$$m(N_r) = m(N \cap [0, 1 - r)) + m(N \cap [1 - r, 1))$$

We need to now show that:

$$[0, 1) = [0, 1 - r) \cup [1 - r, 1)$$

This is fairly quick since we note that $r \in [0, 1)$

(\subseteq) Fix $x \in [0, 1)$. Then if $x < 1 - r$ we have $x \in [0, 1 - r)$. Otherwise we know $x \geq 1 - r$ and so $x \in [1 - r, 1)$.

(\supseteq) Fix $x \in [0, 1 - r)$. Then since $r \geq 0$ we know $x < 1 - r \leq 1$. Therefore $0 \leq x < 1$, and so $x \in [0, 1)$


In the other case, fix $x \in [1 - r, 1)$. Then we know since $r < 1$ that $0 < 1 - r \leq x$. Therefore since $0 < x < 1$ we have $x \in [0, 1)$.

Now consider that:

$$(N \cap [0, 1 - r)) \cup (N \cap [1 - r, 1)) = N \cap ([0, 1 - r) \cup [1 - r, 1)) = N \cap [0, 1) = N$$

The last equality holds since N is a subset of $[0, 1)$. Therefore:

$$m(N_r) = m(N \cap [0, 1 - r)) + m(N \cap [1 - r, 1)) = m(N)$$

And we are done! 

Proof of Q7. We wish to arrive at a contradiction. There are three quick cases:

- Suppose that $m(N) = 0$. Then since \mathbb{Q} is countable we know $R = \mathbb{Q} \cap [0, 1)$ is countable, giving us by axiom (a) and (c) that:

$$1 = m([0, 1)) = m\left(\bigcup_{r \in R} N_r\right) = \sum_{r \in R} m(N_r) = \sum_{r \in R} 0 = 0$$

This is a clear contradiction! Oops!

- Suppose that $m(N) > 0$. Note that R is countable and for $n \geq 2$ we have $0 < \frac{1}{n} < 1$ and so $\frac{1}{n} \in R$. Then using axiom (a), axiom (c), and the fact that $m(N)$ is positive we know that:

$$\begin{aligned}
 1 = m([0, 1)) &= m\left(\bigcup_{r \in R} N_r\right) \\
 &= \sum_{r \in R} m(N_r) \geq \sum_{n=2}^{\infty} m\left(N_{\frac{1}{n}}\right) \\
 &= \sum_{n=2}^{\infty} m(N) = \infty
 \end{aligned}$$

This is clearly true, since we know that $m(N) > 0$ doesn't go to zero, $\sum_{n=2}^{\infty} m(N)$ must diverge to infinity. This is an oops since $1 < \infty$

- Suppose that $m(N) = \infty$ Then since R is countable and $0 \in R = \mathbb{Q} \cap [0, 1)$ we know that by axiom (b) and axiom (c),

$$1 = m([0, 1)) = m\left(\bigcup_{r \in R} N_r\right) = \sum_{r \in R} m(N_r) \geq m(N_0) = m(N) = \infty$$

This cannot be true since $1 < \infty$. Oops!

With all three of these completed, we must conclude that $m(N)$ is undefined!!! Wow!
This is amazing ☺ 