

Math 19620 – Section 50 – Midterm 2

November 17, 2015

Information and Directions

- This exam will last 80 minutes.
- This is a closed-book exam.
- No electronic devices are allowed to be used during this exam.
- Make sure to show all details of your work.

Name: Zev Choroles

Problem	Points	Score
1	25	
2	25	
3	20	
4	20	
5	20	

Your score will be computed out of 100.

Problem 1 (25 points)

Consider this 3×5 matrix:

$$A = \begin{bmatrix} 1 & 2 & 1 & 5 & 3 \\ 4 & 9 & 0 & 17 & 9 \\ 2 & 5 & -2 & 7 & 3 \end{bmatrix}$$

Part 1 (10 points)

Find a basis of the kernel of A .

First, find $\text{RREF}(A)$.

$$\begin{bmatrix} 1 & 2 & 1 & 5 & 3 \\ 4 & 9 & 0 & 17 & 9 \\ 2 & 5 & -2 & 7 & 3 \end{bmatrix} \xrightarrow[\substack{(\text{row } 3) - 2(\text{row } 1)}]{\substack{(\text{row } 2) - 4(\text{row } 1)}} \begin{bmatrix} 1 & 2 & 1 & 5 & 3 \\ 0 & 1 & -4 & -3 & -3 \\ 0 & 1 & -4 & -3 & -3 \end{bmatrix}$$

$$\xrightarrow[\substack{(\text{row } 3) - 1(\text{row } 2)}]{(\text{row } 1) - 2(\text{row } 2)} \begin{bmatrix} 1 & 0 & 9 & 11 & 9 \\ 0 & 1 & -4 & -3 & -3 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{This is } \text{RREF}(A)$$

For each column without a pivot, we get a relation:

$$9 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + (-4) \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + (-1) \begin{bmatrix} 9 \\ -4 \\ 0 \end{bmatrix} + 0 \begin{bmatrix} 11 \\ -3 \\ 0 \end{bmatrix} + 0 \begin{bmatrix} 9 \\ -3 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$11 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + (-3) \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 0 \begin{bmatrix} 9 \\ -4 \\ 0 \end{bmatrix} + (-1) \begin{bmatrix} 11 \\ -3 \\ 0 \end{bmatrix} + 0 \begin{bmatrix} 9 \\ -3 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$9 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + (-3) \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 0 \begin{bmatrix} 9 \\ -4 \\ 0 \end{bmatrix} + 0 \begin{bmatrix} 11 \\ -3 \\ 0 \end{bmatrix} + (-1) \begin{bmatrix} 9 \\ -3 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\text{A basis of } \ker(A) \text{ is } \left\{ \begin{bmatrix} 9 \\ -4 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 11 \\ -3 \\ 0 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 9 \\ -3 \\ 0 \\ 0 \\ -1 \end{bmatrix} \right\}$$

Part 2 (10 points)

Find a basis of the image of A .

For each column of $\text{RREF}(A)$ with a pivot,
take the corresponding column of A .

A basis of $\text{im}(A)$ is $\left\{ \begin{bmatrix} 1 \\ 4 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 9 \\ 5 \end{bmatrix} \right\}$

Part 4 (5 points)

Complete the statement of the rank-nullity theorem: for any $m \times n$ matrix B , we have

$$\underline{\dim(\ker(B))} + \underline{\dim(\text{im}(B))} = \underline{n}$$

(fill in with mathematical expressions)

Use your work in this problem to find the specific values of each of these expressions for the matrix A .
Make sure the integers you write below line up exactly with the corresponding expression above.

$$\underline{3} + \underline{2} = \underline{5}$$

(fill in with integers)

Problem 2 (25 points)

Consider this 3×3 matrix and this basis of $\mathbb{R}^3$:

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 4 & -3 \\ 2 & 7 & -5 \end{bmatrix} \quad \mathcal{B} = \left\{ \vec{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \vec{v}_3 = \begin{bmatrix} 1 \\ 3 \\ 4 \end{bmatrix} \right\}$$

Part 1 (15 points)

Express the matrix A in the basis $\mathcal{B}$.

If B is the matrix A in the basis $\mathcal{B}$, then

$$B = S^{-1}AS \quad \text{where} \quad S = \begin{bmatrix} | & | & | \\ \vec{v}_1 & \vec{v}_2 & \vec{v}_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 3 & 4 \end{bmatrix}.$$

Need to find S^{-1} : take RREF of $[S | I_3]$.

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & 2 & 3 & 0 & 1 & 0 \\ 1 & 3 & 4 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & -1 & 1 & 0 \\ 0 & 2 & 3 & -1 & 0 & 1 \end{array} \right]$$

$$\rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & -1 & 2 & -1 & 0 \\ 0 & 1 & 2 & -1 & 1 & 0 \\ 0 & 0 & -1 & 1 & -2 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & -1 & 2 & -1 & 0 \\ 0 & 1 & 2 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 2 & -1 \end{array} \right]$$

$$\rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & -1 \\ 0 & 1 & 0 & 1 & -3 & 2 \\ 0 & 0 & 1 & -1 & 2 & -1 \end{array} \right] \quad \text{Thus } S^{-1} = \begin{bmatrix} 1 & 1 & -1 \\ 1 & -3 & 2 \\ -1 & 2 & -1 \end{bmatrix}.$$

$$B = S^{-1}AS = \begin{bmatrix} 1 & 1 & -1 \\ 1 & -3 & 2 \\ -1 & 2 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 2 & 4 & -3 \\ 2 & 7 & -5 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 3 & 4 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 1 & -1 \\ 1 & -3 & 2 \\ -1 & 2 & -1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 3 & 1 & 2 \\ 4 & 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \quad \text{This is } A \text{ expressed in the basis } \mathcal{B}.$$

Part 2 (10 points)

Let $\vec{w} = 5\vec{v}_1 + 7\vec{v}_2 + 3\vec{v}_3$, so that $[\vec{w}]_{\mathcal{B}} = \begin{bmatrix} 5 \\ 7 \\ 3 \end{bmatrix}$.

Express the vector $A\vec{w}$ in the basis $\mathcal{B}$, i.e. compute $[A\vec{w}]_{\mathcal{B}}$.

(You can do this directly, by computing $\vec{w}$, then computing $A\vec{w}$, then finding that vector's expression in the basis $\mathcal{B}$. There is also a much faster way using your answer from Part 1. Either is acceptable.)

Fast way: If B is the matrix A expressed in the basis $\mathcal{B}$, then $B[\vec{v}]_{\mathcal{B}} = [A\vec{v}]_{\mathcal{B}}$ for any vector $\vec{v} \in \mathbb{R}^3$.

Thus $\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 5 \\ 7 \\ 3 \end{bmatrix} = \begin{bmatrix} 7 \\ 3 \\ 5 \end{bmatrix}$ is equal to $[A\vec{w}]_{\mathcal{B}}$.

Slow way: $\vec{w} = 5\vec{v}_1 + 7\vec{v}_2 + 3\vec{v}_3 = 5\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + 7\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + 3\begin{bmatrix} 1 \\ 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 15 \\ 28 \\ 38 \end{bmatrix}$

$$A\vec{w} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 4 & -3 \\ 2 & 7 & -5 \end{bmatrix} \begin{bmatrix} 15 \\ 28 \\ 38 \end{bmatrix} = \begin{bmatrix} 15 \\ 28 \\ 36 \end{bmatrix}$$

If $\vec{v}$ is any vector in $\mathbb{R}^3$, then $[\vec{v}]_{\mathcal{B}} = S^{-1}\vec{v}$.

$$[A\vec{w}]_{\mathcal{B}} = \begin{bmatrix} 1 & 1 & -1 \\ 1 & -3 & 2 \\ -1 & 2 & 1 \end{bmatrix} \begin{bmatrix} 15 \\ 28 \\ 36 \end{bmatrix} = \begin{bmatrix} 7 \\ 3 \\ 5 \end{bmatrix}$$

Problem 3 (20 points)

Let V be the subspace of $\mathbb{R}^4$ with these vectors as a basis:

$$\left\{ \vec{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \vec{v}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix} \right\}$$

Part 1 (10 points)

Find an orthonormal basis of V .

Gram-Schmidt:

$$\vec{w}_1 = \frac{1}{\|\vec{v}_1\|} \vec{v}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \\ 0 \end{bmatrix}$$

$\vec{w}_2$ is formed by taking $\vec{v}_2 - (\vec{v}_2 \cdot \vec{w}_1) \vec{w}_1$ and dividing it by its length

$$\vec{v}_2 - (\vec{v}_2 \cdot \vec{w}_1) \vec{w}_1 = \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} - 0 \cdot \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} \text{ which has length } \sqrt{2}$$

$$\text{so } \vec{w}_2 = \begin{bmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \\ 0 \\ 0 \end{bmatrix}$$

$\vec{w}_3$ is formed by taking $\vec{v}_3 - (\vec{v}_3 \cdot \vec{w}_1) \vec{w}_1 - (\vec{v}_3 \cdot \vec{w}_2) \vec{w}_2$ and dividing it by its length

$$\vec{v}_3 - (\vec{v}_3 \cdot \vec{w}_1) \vec{w}_1 - (\vec{v}_3 \cdot \vec{w}_2) \vec{w}_2 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix} - \sqrt{2} \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \\ 0 \end{bmatrix} - 0 \begin{bmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} \text{ which has length } 1$$

$$\text{so } \vec{w}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}$$

Thus $\left\{ \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} \right\}$ is an orthonormal basis of V

Part 2 (10 points)

Find the matrix of $\text{proj}_V : \mathbb{R}^4 \rightarrow \mathbb{R}^4$.

The matrix of proj_V is QQ^T , where $Q = [\vec{v}_1 | \dots | \vec{v}_k]$

is a matrix whose columns are an orthonormal basis of V .

Thus, with $Q = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} & 0 \\ 1/\sqrt{2} & -1/\sqrt{2} & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$, we have

$$QQ^T = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} & 0 \\ 1/\sqrt{2} & -1/\sqrt{2} & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} & 0 & 0 \\ 1/\sqrt{2} & -1/\sqrt{2} & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Problem 4 (20 points)

Part 1 (10 points)

Compute the value of $\det(A)$, where

$$A = \begin{bmatrix} 3 & 6 & 3 \\ 4 & 10 & 0 \\ 2 & 0 & 8 \end{bmatrix}$$

We use Gauss-Jordan elimination.

$$\begin{bmatrix} 3 & 6 & 3 \\ 4 & 10 & 0 \\ 2 & 0 & 8 \end{bmatrix} \quad \text{This matrix has determinant} = \det(A)$$

$\left. \begin{array}{l} \text{ } \end{array} \right\} \text{(row 1)} \cdot \frac{1}{3} \quad \underline{\text{Multiplies determinant by } \frac{1}{3}}$

$$\begin{bmatrix} 1 & 2 & 1 \\ 4 & 10 & 0 \\ 2 & 0 & 8 \end{bmatrix} \quad \text{This matrix has determinant} = \frac{1}{3} \det(A)$$

$\left. \begin{array}{l} \text{ } \end{array} \right\} \begin{array}{l} \text{(row 2)} - 4(\text{row 1}) \\ \text{(row 3)} - 2(\text{row 1}) \end{array} \quad \underline{\text{Both leave determinant unchanged}}$

$$\begin{bmatrix} 1 & 2 & 1 \\ 0 & 2 & -4 \\ 0 & -4 & 6 \end{bmatrix} \quad \text{This matrix has determinant} = \frac{1}{3} \det(A)$$

$\left. \text{ } \right\} \text{(row 2)} \cdot \frac{1}{2} \quad \underline{\text{Multiplies determinant by } \frac{1}{2}}$

$$\begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & -2 \\ 0 & -4 & 6 \end{bmatrix} \quad \text{This matrix has determinant} = \frac{1}{6} \det(A)$$

$\left. \begin{array}{l} \text{ } \end{array} \right\} \begin{array}{l} \text{(row 1)} - 2(\text{row 2}) \\ \text{(row 3)} + 4(\text{row 2}) \end{array} \quad \underline{\text{Both leave determinant unchanged}}$

$$\begin{bmatrix} 1 & 0 & 5 \\ 0 & 1 & -2 \\ 0 & 0 & -2 \end{bmatrix} \quad \text{This matrix has determinant} = \frac{1}{6} \det(A)$$

$\left. \text{ } \right\} \text{(row 3)} \cdot \frac{-1}{2} \quad \underline{\text{Multiplies determinant by } \frac{-1}{2}}$

$$\begin{bmatrix} 1 & 0 & 5 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} \quad \text{This matrix has determinant} = -\frac{1}{12} \det(A)$$

$\left. \begin{array}{l} \text{ } \end{array} \right\} \begin{array}{l} \text{(row 1)} - 5(\text{row 3}) \\ \text{(row 2)} + 2(\text{row 3}) \end{array} \quad \underline{\text{Both leave determinant unchanged}}$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \text{This matrix has determinant} = -\frac{1}{12} \det(A)$$

But $\det(\text{identity matrix}) = 1$. Thus $1 = -\frac{1}{12} \det(A)$, so

$$\det(A) = -12$$

Part 2 (10 points)

Compute the value of $\det(B)$, where

$$B = \begin{bmatrix} 3 & 6 & 3 \\ 4 & 10 & 0 \\ 14 & 0 & 56 \end{bmatrix}$$

Note that B is the same as A , except that one of its rows has been multiplied by 7.

Since $\det$ is multilinear in the rows of a matrix,

$$\det \left(\begin{bmatrix} v_1 \\ v_2 \\ 7v_3 \end{bmatrix} \right) = 7 \det \left(\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \right)$$

$$\text{Thus } \det(B) = 7 \cdot \det(A) = 7 \cdot (-12) = -84$$

Problem 5 (20 points)

Consider the system of linear equations $A\vec{x} = \vec{b}$, where

$$A = \begin{bmatrix} 1 & 0 & -1 \\ 1 & 0 & -1 \\ 2 & 1 & 0 \\ 2 & 1 & 0 \end{bmatrix}, \quad \vec{b} = \begin{bmatrix} -1 \\ 1 \\ -3 \\ 3 \end{bmatrix}$$

Part 1 (15 points)

Find the least-squares solutions of $A\vec{x} = \vec{b}$.

The least-squares solutions are the exact solutions to

$$A^T A \vec{x} = A^T \vec{b}$$

$$A^T A = \begin{bmatrix} 1 & 1 & 2 & 2 \\ 0 & 0 & 1 & 1 \\ -1 & -1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \\ 1 & 0 & -1 \\ 2 & 1 & 0 \\ 2 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 10 & 4 & -2 \\ 4 & 2 & 0 \\ -2 & 0 & 2 \end{bmatrix}$$

$$A^T \vec{b} = \begin{bmatrix} 1 & 1 & 2 & 2 \\ 0 & 0 & 1 & 1 \\ -1 & -1 & 0 & 0 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \\ -3 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

So we want to find the solutions to $\begin{bmatrix} 10 & 4 & -2 \\ 4 & 2 & 0 \\ -2 & 0 & 2 \end{bmatrix} \vec{x} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$

Take RREF of augmented matrix:

$$\left[\begin{array}{ccc|c} 10 & 4 & -2 & 0 \\ 4 & 2 & 0 & 0 \\ -2 & 0 & 2 & 0 \end{array} \right] \rightsquigarrow \left[\begin{array}{ccc|c} 1 & 2/5 & -1/5 & 0 \\ 4 & 2 & 0 & 0 \\ -2 & 0 & 2 & 0 \end{array} \right] \rightsquigarrow \left[\begin{array}{ccc|c} 1 & 2/5 & -1/5 & 0 \\ 0 & 2/5 & 4/5 & 0 \\ 0 & 4/5 & 8/5 & 0 \end{array} \right]$$

$$\rightarrow \left[\begin{array}{ccc|c} 1 & 2/5 & -1/5 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 4/5 & 8/5 & 0 \end{array} \right] \rightsquigarrow \left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

This is RREF.

There is one free variable, corresponding to the column without a pivot.

Least-squares solutions to $A\vec{x} = \vec{b}$ are $\left\{ \begin{bmatrix} + \\ -2+ \\ + \end{bmatrix} : + \in \mathbb{R} \right\}$

Part 2 (5 points)

Find the vector in $\text{im}(A)$ that is closest to $\vec{b}$, otherwise known as $\text{proj}_{\text{im}(A)}(\vec{b})$.

(For extra credit up to an additional 5 points, give an intuitive explanation of the answer to this part.)

For any least-squares solution $\vec{x}$, the value of $A\vec{x}$ is $\text{proj}_{\text{im}(A)}(\vec{b})$. Indeed, the idea of a least-squares solution to $A\vec{x} = \vec{b}$ is to be a choice of $\vec{x}$ that makes the distance $\|\vec{b} - A\vec{x}\|$ as small as possible.

Thus, we can take any least-squares solution $\vec{x}$, and multiply it by A to get $\text{proj}_{\text{im}(A)}(\vec{b})$.

$$A\vec{x} = \begin{bmatrix} 1 & 0 & -1 \\ 1 & 0 & -1 \\ 2 & 1 & 0 \\ 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} + \\ -2+ \\ + \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} = \text{proj}_{\text{im}(A)}\left(\begin{bmatrix} -1 \\ 1 \\ -3 \\ 3 \end{bmatrix}\right).$$

$$\text{Since } \text{im}(A) = \text{span}\left(\begin{bmatrix} 1 \\ 1 \\ 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ -1 \\ 0 \\ 0 \end{bmatrix}\right) = \left\{ \begin{bmatrix} r \\ r \\ s \\ s \end{bmatrix} : r, s \in \mathbb{R} \right\},$$

the subspace of $\mathbb{R}^4$ comprised of the vectors whose 1st and 2nd coordinates are equal and whose 3rd and 4th coordinates are equal, the vector in $\text{im}(A)$ that is closest to $\vec{b} = \begin{bmatrix} -1 \\ 1 \\ -3 \\ 3 \end{bmatrix}$, a vector whose 1st and 2nd, and whose 3rd and 4th, coordinates are opposites, must be $\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$.

To choose any r or s other than $r=s=0$ would only make a vector $\begin{bmatrix} r \\ r \\ s \\ s \end{bmatrix}$ further from $\begin{bmatrix} -1 \\ 1 \\ -3 \\ 3 \end{bmatrix}$.