

Math 13100 – Section 58 – Final Exam

March 15, 2016

Information and Directions

- This exam will last 120 minutes.
- This is a closed-book exam.
- No electronic devices are allowed to be used during this exam.
- Partial credit is given for showing your calculations and explaining your thoughts.

Name: _____

SOLUTIONS

Problem	Points	Score
1	8	
2	8	
3	4	
4	16	
5	16	
6	16	
7	16	
8	8	
9	8	

Problem 1 (8 points)

Consider the inequality

$$-5 < 2x - 1 < 7$$

Part 1 (4 points)

Express the set of solutions of this inequality as an interval.

$$-5 < 2x - 1 < 7$$

$$-5 + 1 < 2x < 7 + 1$$

$$-4 < 2x < 8$$


$$-4/2 < 2x/2 < 8/2$$

$$-2 < x < 4$$

Solutions are the real numbers x in the interval $(-2, 4)$

Part 2 (4 points)

For which numbers a and b does $|x - a| < b$ have the same solutions as the above inequality?



The center of the interval $(-2, 4)$ is the number 1, and the ends are a distance of 3 away from the center. Thus $-2 < x < 4$ is equivalent to $|x - 1| < 3$.

Alternatively, you can remember that

$$|x - a| < b \iff a - b < x < a + b$$

and look for the numbers a and b such that
$$\begin{aligned} a - b &= -2 \\ a + b &= 4 \end{aligned}$$

You can solve by substitution, e.g., $a - b = -2$ implies $a = b - 2$,

so $a + b = 4$ becomes $(b - 2) + b = 4$, hence $2b = 6$, hence $b = 3$, and $a = b - 2 = 1$.

Problem 2 (8 points)

Define the functions

$$f(x) = x^2 + 1$$

$$g(x) = \frac{\sqrt{x-2}}{x-5}$$

Part 1 (4 points)

What is the natural domain of g ?

$\sqrt{x-2}$ is only defined when $x-2 \geq 0$,

and $\frac{1}{x-5}$ is only defined when $x-5 \neq 0$,

So $\frac{\sqrt{x-2}}{x-5}$ is only defined when $x \geq 2$, except $x = 5$

Written as a union of intervals, the domain is $[2, 5) \cup (5, \infty)$

Part 2 (4 points)

What is the value of $(f \circ g)(6)$?


$$\begin{aligned}(f \circ g)(6) &= f(g(6)) \\&= f\left(\frac{\sqrt{6-2}}{6-5}\right) \\&= f\left(\frac{\sqrt{4}}{1}\right) \\&= f(2) \\&= 2^2 + 1 \\&= 5\end{aligned}$$

Problem 3 (4 points)

Graph the function $f(x) = (x - 2)^2 - 1$.

For at least three different points on your graph, label the point with its x - and y -coordinates.

(Hint: take the graph of x^2 and translate it by the appropriate amounts horizontally and vertically.)



Problem 4 (16 points)

Part 1 (8 points)

Let $f(x)$ be a function. Give the precise, mathematical meaning of the statement

$$\lim_{x \rightarrow 5} f(x) = 2$$

For any $\epsilon > 0$, there exists some $\delta > 0$

such that $0 < |x - 5| < \delta \Rightarrow |f(x) - 2| < \epsilon$

Part 2 (8 points)

Give a complete proof, without appealing to theorems about continuity, that

$$\lim_{x \rightarrow 5} x^2 - 3x - 8 = 2$$

We want to show that for any $\epsilon > 0$, there exists a $\delta > 0$ such that

$0 < |x - 5| < \delta \Rightarrow |x^2 - 3x - 8 - 2| < \epsilon$. Factoring the right side,

$$|x^2 - 3x - 10| < \epsilon \iff |(x - 5)(x + 2)| < \epsilon$$

$$\iff |x - 5| \cdot |x + 2| < \epsilon$$

If we decide in advance to choose a $\delta < 1$, then by the triangle inequality, $|x + 2| = |(x - 5) + 7| \leq |x - 5| + |7| < \delta + 7 < 8$.

Thus, $|x - 5| \cdot (\text{something} < 8) < \epsilon$, hence $|x - 5| < \frac{\epsilon}{8}$.

Therefore, no matter what $\epsilon > 0$ is, if we choose $\delta = \min \{1, \frac{\epsilon}{8}\}$,

we can guarantee that $0 < |x - 5| < \delta \Rightarrow |x^2 - 3x - 10| < \epsilon$.

(Keep writing on the back of the page if you need more space.)

Problem 5 (16 points)

Part 1 (8 points)

Let c be a real number, and let f be a function. Give the precise definition of what it means to say

" f is continuous at c "

(Hint: there are three conditions that must be met.)

- $f(c)$ is defined
- $\lim_{x \rightarrow c} f(x)$ exists
- $\lim_{x \rightarrow c} f(x) = f(c)$

Part 2 (8 points)

Are there any discontinuities of the function

$$f(x) = \frac{x^2 - 5x + 6}{x - 3} ?$$

(That is, can you find any c where f is not continuous at c ?) If so, are they removable discontinuities?

Explain your answer or thought process.

Theorem A from section 1.6 says that a rational function is continuous at any real number c where its denominator is not zero. Thus, f is continuous at any $c \neq 3$. At any $c \neq 3$, we have $f(c) = \frac{(c-2)(c-3)}{(c-3)} = c-2$.

f is not continuous at 3 because $f(3)$ is not defined.

However, if we define $f(3) = 1$, then $f(x) = x-2$ for all x , and since $x-2$ is continuous at 3, this demonstrates that the discontinuity that f had was removable, since we could turn f into a continuous function by choosing a new definition of $f(3)$.

Problem 6 (16 points)

In this problem, define the function $f(x) = (x^2 - 7)^5$.

Part 1 (8 points)

Using the chain rule, compute the derivative of f .

$$\begin{aligned} g(x) &= x^5 & h(x) &= x^2 - 7 & f(x) &= (x^2 - 7)^5 = (g \circ h)(x) \\ g'(x) &= 5x^4 & h'(x) &= 2x \end{aligned}$$

$$\begin{aligned} \text{so } f'(x) &= g'(h(x)) \cdot h'(x) \\ &= 5(x^2 - 7)^4 \cdot (2x) \\ &= 10x(x^2 - 7)^4 \end{aligned}$$

Part 2 (8 points)

Find the equation of the tangent line to the graph of $y = f(x)$ at 3, i.e., the tangent line that goes through the point

$$(3, f(3)) = (3, (3^2 - 7)^5) = (3, 2^5) = (3, 32)$$

The tangent line to the graph of $y = f(x)$ at 3 is characterized by these two properties:

- It goes through $(3, f(3)) = (3, 32)$.
- Its slope is $f'(3) = 10 \cdot 3 \cdot (3^2 - 7)^4 = 30 \cdot 2^4 = 480$

By the point-slope formula, the equation of the line is

$$y - 32 = 480(x - 3)$$

$$y = 480x - 1408$$

Problem 7 (16 points)

Part 1 (8 points)

Using implicit differentiation, find $\frac{dy}{dx}$ where $x^2y^2 + 4xy = 12y$.

$$\frac{d}{dx}(x^2y^2 + 4xy) = \frac{d}{dx}(12y)$$

$$(2xy^2 + 4y) + (2x^2y + 4x - 12)\frac{dy}{dx} = 0$$

$$\frac{d}{dx}(x^2y^2) + \frac{d}{dx}(4xy) = \frac{d}{dx}(12y)$$

$$\frac{dy}{dx} = \frac{-2xy^2 - 4y}{2x^2y + 4x - 12}$$

product
rule +
chain rule

$$(2xy^2 + x^2 \cdot 2y \frac{dy}{dx}) + 4(y + x \frac{dy}{dx}) = 12 \frac{dy}{dx}$$

Part 2 (8 points)

Find the equation of the tangent line to the graph of

$$x^2y^2 + 4xy = 12y$$

that goes through the point (2, 1).

The tangent line to the graph at (2, 1) is characterized by:

- It goes through (2, 1)
- Its slope is $\frac{dy}{dx}$ when $x=2$, $y=1$:

$$\frac{-2(2)(1)^2 - 4(1)}{2(2)^2(1) + 4(2) - 12} = \frac{-8}{4} = -2$$

By the point-slope formula, the equation of the line is

$$y - 1 = (-2)(x - 2)$$

$$y = -2x + 5$$

Problem 8 (8 points)

Use derivatives to approximate the value of $\sqrt{16.1}$.

(Hint: you are being asked to approximate $f(16 + 0.1)$, where $f(x) = \sqrt{x}$. You should start by finding $f(16)$ and $f'(16)$.)

To approximate the value of $f(x+h)$, we use

$$\underset{\text{(approx)}}{f(x+h)} \approx f(x) + f'(x)h$$

In this problem, $f(x) = \sqrt{x} = x^{1/2}$, so $f'(x) = \frac{1}{2\sqrt{x}} = (1/2)x^{(-1/2)}$.

Thus, $f(16) = \sqrt{16} = 4$ and $f'(16) = \frac{1}{2\sqrt{16}} = \frac{1}{8}$,

$$\text{So } f(16 + 0.1) \approx 4 + \frac{1}{8} \cdot 0.1 = 4.0125$$

(The actual value is 4.0124805... so this is pretty close.)

Problem 9 (8 points)

Each edge of a cube is growing at a rate of 3 cm/sec. How quickly is the volume of the cube increasing when the edges of the cube are exactly 12 cm long?

(Hint: The volume of a cube whose edges have length x is equal to x^3 .)

We want to find the rate of change of volume, $\frac{dV}{dt}$

We know the relationship between volume and edge length: $V = x^3$

We know the rate of change of edge length: $\frac{dx}{dt} = 3 \text{ cm/sec}$

We know we're asking about the time at which edges $x = 12 \text{ cm}$

Taking the derivative $\frac{d}{dt}$ of the relationship $V = x^3$,

we get $\frac{dV}{dt} = 3x^2 \frac{dx}{dt}$, so

$$\frac{dV}{dt} = 3(12 \text{ cm})^2 (3 \text{ cm/sec}) = 1296 \text{ cm}^3/\text{sec}$$