

# Math 13200 – Section 58 – Final Exam Sample

June 4, 2016

## Information and Directions

- This exam will last 120 minutes.
- This is a closed-book exam.
- No electronic devices are allowed to be used during this exam.
- **Partial credit is given for showing your calculations and explaining your thoughts.**

Name: SOLUTIONS

| Problem | Points | Score |
|---------|--------|-------|
| 1       | 10     |       |
| 2       | 10     |       |
| 3       | 10     |       |
| 4       | 10     |       |
| 5       | 10     |       |
| 6       | 10     |       |
| 7       | 10     |       |
| 8       | 10     |       |
| 9       | 10     |       |
| 10      | 10     |       |

## Problem 1 (10 points)

### Part 1 (5 points)

What is the definition of the number  $e$ ?

$e$  is the unique real number with the property that  $\ln(e) = 1$ .

### Part 2 (5 points)

What is  $e^{\ln(2)+\ln(7)}$ ?

Because  $e^{a+b} = e^a \cdot e^b$ , and  $e^{\ln(x)} = x$ ,

we have  $e^{\ln(2)+\ln(7)} = e^{\ln(2)} \cdot e^{\ln(7)} = 2 \cdot 7 = 14$

Alternatively, since  $\ln(ab) = \ln(a) + \ln(b)$ ,

we have  $e^{\ln(2)+\ln(7)} = e^{\ln(14)} = 14$ .

## Problem 2 (10 points)

### Part 1 (5 points)

What is the equation of the tangent line to  $y = \cos(x)$  at  $x = \frac{\pi}{4}$ ?

• Point : goes through  $(\frac{\pi}{4}, \cos(\frac{\pi}{4})) = (\frac{\pi}{4}, \frac{1}{\sqrt{2}})$

• Slope : derivative is  $\frac{d}{dx} \cos(x) = -\sin(x)$ , so slope is  
 $-\sin(\frac{\pi}{4}) = -\frac{1}{\sqrt{2}}$

By the point-slope formula, the equation of the tangent line is

$$(y - \frac{1}{\sqrt{2}}) = (-\frac{1}{\sqrt{2}})(x - \frac{\pi}{4})$$

$$y = (-\frac{1}{\sqrt{2}})x + (\frac{\pi}{4\sqrt{2}} + \frac{1}{\sqrt{2}})$$

### Part 2 (5 points)

What is the equation of the tangent line to  $y = \ln(\sin(x))$  at  $x = \frac{\pi}{2}$ ?

• Point : goes through  $(\frac{\pi}{2}, \ln(\sin(\frac{\pi}{2}))) = (\frac{\pi}{2}, \ln(1)) = (\frac{\pi}{2}, 0)$

• Slope : derivative is  $\frac{d}{dx} \ln(\sin(x)) \stackrel{\text{chain rule}}{=} \frac{1}{\sin(x)} \cdot \frac{d}{dx}(\sin(x)) = \frac{\cos(x)}{\sin(x)} = \cot(x)$

$$\text{so slope is } \cot(\frac{\pi}{2}) = \frac{\cos(\frac{\pi}{2})}{\sin(\frac{\pi}{2})} = \frac{0}{1} = 0$$

By the point-slope formula, the equation of the tangent line is

$$(y - 0) = 0(x - \frac{\pi}{2})$$

$$y = 0$$

### Problem 3 (10 points)

#### Part 1 (5 points)

Find the derivative  $\frac{d}{dx} \sin^3(x) \cos(x)$ .

$$\begin{aligned} \frac{d}{dx} (\sin^3(x) \cos(x)) &\stackrel{\text{product rule}}{=} \frac{d}{dx} (\sin^3(x)) \cos(x) + \sin^3(x) \frac{d}{dx} (\cos(x)) \\ &\stackrel{\text{chain rule}}{=} (3 \sin^2(x) \cdot \frac{d}{dx} (\sin(x))) \cos(x) + \sin^3(x) \frac{d}{dx} (\cos(x)) \\ \frac{d}{dx} (\sin(x)) &= \cos(x) \\ \frac{d}{dx} (\cos(x)) &= -\sin(x) \\ &\stackrel{\text{simplifying}}{=} 3 \sin^2(x) \cos^2(x) - \sin^4(x) \end{aligned}$$

#### Part 2 (5 points)

Find the antiderivative  $\int \sin^3(x) \cos(x) dx$ .

u-substitution with  $u = \sin(x)$

therefore,  $du = \cos(x) dx$

$$\begin{aligned} \int \sin^3(x) \cos(x) dx &= \int (\sin(x))^3 \cos(x) dx = \int u^3 du = \frac{u^4}{4} + C \\ &= \frac{\sin^4(x)}{4} + C \end{aligned}$$

## Problem 4 (10 points)

### Part 1 (5 points)

Find the derivative  $\frac{d}{dx} e^x \sin(e^x)$ .

$$\frac{d}{dx} (e^x \sin(e^x)) \stackrel{\text{product rule}}{=} \frac{d}{dx}(e^x) \sin(e^x) + e^x \frac{d}{dx}(\sin(e^x))$$

$$\stackrel{\text{chain rule}}{=} \frac{d}{dx}(e^x) \sin(e^x) + e^x (\cos(e^x) \cdot \frac{d}{dx}(e^x))$$

$$\frac{d}{dx}(e^x) = e^x \stackrel{\text{chain rule}}{=} e^x \sin(e^x) + e^x (\cos(e^x) e^x)$$

$$\stackrel{\text{simplifying}}{=} e^x \sin(e^x) + e^{2x} \cos(e^x)$$

### Part 2 (5 points)

Find the antiderivative  $\int e^x \sin(e^x) dx$ .

$u$ -substitution with  $u = e^x$

$$\text{therefore, } du = e^x dx$$

$$\int e^x \sin(e^x) dx = \int \sin(e^x) e^x dx = \int \sin(u) du = -\cos(u) + C$$

$$= -\cos(e^x) + C$$

### Problem 5 (10 points)

For this problem, define the function

$$G(x) = \int_{13}^x \ln(t^2 + 1) dt$$

#### Part 1 (5 points)

What is  $G'(x)$ ?

1<sup>st</sup> FTC says that  $\frac{d}{dx} \left( \int_a^x f(t) dt \right) = f(x)$

Therefore  $G'(x) = \ln(x^2 + 1)$

#### Part 2 (5 points)

Is  $G$  concave up, concave down, or neither at the point  $x = 2$ ?

2<sup>nd</sup> Derivative Test:

If  $G''(x) > 0$ , then  $G$  is concave up at  $x$

If  $G''(x) < 0$ , then  $G$  is concave down at  $x$

(No conclusion if  $G''(x) = 0$ )

$$\frac{d}{dx} (\ln(x)) = \frac{1}{x}, \text{ so } G''(x) = \frac{d}{dx} G'(x) = \frac{d}{dx} \ln(x^2 + 1)$$

chain rule  $\Rightarrow \frac{1}{x^2 + 1} \cdot \frac{d}{dx} (x^2 + 1) = \frac{2x}{x^2 + 1}$

$$G''(2) = \frac{4}{5} > 0, \text{ so } G \text{ is concave up at } 2$$

**Problem 6 (10 points)**

Find the definite integral  $\int_1^{e^3} \frac{\ln(x)}{x} dx$ . For this problem, you must simplify fully.

$u$  - substitution with  $u = \ln(x)$

$$\text{therefore, } du = \frac{1}{x} dx$$

$$1 \rightsquigarrow \ln(1) = 0$$

$$e^3 \rightsquigarrow \ln(e^3) = 3 \cdot \ln(e) = 3 \cdot 1 = 3$$

$$\int_1^{e^3} \frac{\ln(x)}{x} dx = \int_1^{e^3} \ln(x) \cdot \frac{1}{x} dx = \int_0^3 u du = \left[ \frac{u^2}{2} \right]_0^3$$


$$= \left( \frac{9}{2} - 0 \right) = \frac{9}{2}$$

### Problem 7 (10 points)

Consider the region bounded by the lines  $x = 0$  and  $y = ex$ , and the curve  $y = e^x$ .

#### Part 1 (5 points)


Plot the region on this grid. For each intersection between the boundaries described above, label the point with its  $x$ - and  $y$ -coordinates. (Remember,  $e \approx 2.7$ .)



#### Part 2 (5 points)

Find the area of the region.

$$\text{Area} = \int_{\text{one side}}^{\text{other side}} (\text{length of slice at } x) \, dx$$




$$= \int_0^1 (e^x - ex) \, dx = \left[ e^x - e \frac{x^2}{2} \right]_0^1$$


$$= \left( e - \frac{e}{2} \right) - \left( 1 - 0 \right) = \frac{e}{2} - 1$$

### Problem 8 (10 points)

Consider the region bounded by the lines  $y = 0$ ,  $x = 0$ , and  $x = \sqrt{\pi}$ , and the curve  $y = \sin(x^2)$ . If we rotate the region around the  $y$ -axis, what is the volume of the resulting shape?



( $\sin(x^2)$  between 0 and  $\sqrt{\pi}$  looks approximately like  $\sin(x)$  does between 0 and  $\pi$ )



Shell at radius  $x$  has height  $\sin(x^2)$

area of shell at radius  $x = 2\pi r h = 2\pi x \sin(x^2)$

$$\text{Volume} = \int_{\text{inner radius}}^{\text{outer radius}} (\text{area of shell at radius } x) dx = \int_0^{\sqrt{\pi}} 2\pi x \sin(x^2) dx$$

$$\int_0^{\sqrt{\pi}} 2\pi x \sin(x^2) dx = \pi \int_0^{\sqrt{\pi}} \sin(x^2) \cdot 2x dx$$

$u$  - substitution with  $u = x^2$

$$\text{therefore, } du = 2x dx$$

$$0 \rightsquigarrow 0^2 = 0$$

$$\sqrt{\pi} \rightsquigarrow (\sqrt{\pi})^2 = \pi$$



$$= \pi \int_0^{\pi} \sin(u) du = \pi \left[ -\cos(u) \right]_0^{\pi}$$

$$= \pi (-(-1) - (-1)) = 2\pi$$


### Problem 9 (10 points)

Consider the region bounded by the line  $y = 2x$  and the curve  $y = x^2$ .

If a shape has this region as its base, and every cross-section of the shape which is perpendicular to the  $x$ -axis is a circle, what is the volume of the shape?



Slice at  $x$  looks like



Circle of diameter  $d$  has area  $\pi r^2 = \pi \left(\frac{d}{2}\right)^2 = \frac{\pi}{4} d^2$

$$\text{Volume} = \int_{\text{one side}}^{\text{other side}} (\text{area of slice at } x) dx = \int_0^2 \frac{\pi}{4} (2x - x^2)^2 dx =$$

$$\frac{\pi}{4} \int_0^2 4x^2 - 4x^3 + x^4 dx$$

$$\frac{\pi}{4} \int_0^2 4x^2 - 4x^3 + x^4 dx = \frac{\pi}{4} \left[ \frac{4x^3}{3} - x^4 + \frac{x^5}{5} \right]_0^2$$

$$= \frac{\pi}{4} \left( \left( \frac{32}{3} - 16 + \frac{32}{5} \right) - (0) \right)$$


$$= \frac{\pi}{4} \left( \frac{160}{15} - \frac{240}{15} + \frac{96}{15} \right)$$

$$= \frac{\pi}{4} \left( \frac{16}{15} \right)$$

$$= \frac{4\pi}{15}$$

### Problem 10 (10 points)

Suppose that I want to make a rectangle with the largest area I can, but it is required that the rectangle sits inside the triangle formed by  $y = 0$ ,  $y = -2x + 5$ , and  $y = 2x + 5$  (see the picture below).



What are the width and the height of the rectangle I should choose to get the maximum area?

$$\begin{aligned}\text{Area of the rectangle is } (\text{base}) \cdot (\text{height}) &= (2x)(-2x+5) \\ &= -4x^2 + 10x\end{aligned}$$

So we are looking for the maximum of the function  $A(x) = -4x^2 + 10x$  on the interval  $[0, \frac{5}{2}]$

Look at critical points:

Stationary points:  $A'(x) = -8x + 10$ , so  $A'(x) = 0$  when  $x = \frac{5}{4}$

Singular points: none,  $A'(x)$  always exists

End points:  $0, \frac{5}{2}$

$$A(0) = 0$$

$$A\left(\frac{5}{4}\right) = \frac{25}{4}$$

$$A\left(\frac{5}{2}\right) = 0$$

Since  $A\left(\frac{5}{4}\right) = \frac{25}{4}$  is biggest, maximum area occurs when  $x = \frac{5}{4}$

$$\text{width} = 2 \cdot \frac{5}{4} = \frac{5}{2} \quad \text{height} = -2\left(\frac{5}{4}\right) + 5 = \frac{5}{2}$$