

The companion section for classical groups

T. Hameister, B. C. Ngô

August 14, 2023

1 Introduction

Let G be a reductive group over k , and denote by \mathfrak{g} its Lie algebra. The Chevalley map

$$\chi: \mathfrak{g} \rightarrow \mathfrak{g} // G,$$

where $\mathfrak{g} // G := \text{Spec}(k[\mathfrak{g}]^G)$ denotes the invariant theoretic quotient of \mathfrak{g} by the adjoint action of G , is of fundamental importance in the construction of the Hitchin system [4]. In particular, for $\mathfrak{g} = \mathfrak{gl}_n$, χ sends a matrix to its characteristic polynomial.

In [7], Kostant exhibited a section of the Chevalley map for a general reductive group G under the assumption that the characteristic of k does not divide the order of the Weyl group. As explained in [8], this section can be used to construct sections of the Hitchin fibration and affine Springer fibers. However, Kostant's construction has two important detractors: on the one hand, it requires a bound on the characteristic of the field that depends on the group G , which is usually not optimal for classical groups, and on the other hand, it can be counter-intuitive for computations. To illustrate this latter point, consider the case $G = \text{GL}_3(k)$, in which case $\mathfrak{g} // G = \mathbb{A}^3$ is the 3-dimensional affine space. The Kostant section is the map sending

$$(a_1, a_2, a_3) \in \mathfrak{g} // G \quad \mapsto \quad \begin{pmatrix} \frac{a_1}{3} & \frac{a_1^2}{6} + \frac{a_2}{2} & -\frac{4a_1^3}{27} - \frac{a_1 a_2}{3} - a_3 \\ 1 & \frac{a_1}{3} & \frac{a_1^2}{6} + \frac{a_2}{2} \\ 0 & 1 & \frac{a_1}{3} \end{pmatrix} \in \mathfrak{g}$$

If you introduced this problem to an undergraduate student of linear algebra, of course, they would not give you the answer above; they might instead suggest the map:

$$(a_1, a_2, a_3) \in \mathfrak{g} // G \quad \mapsto \quad \begin{pmatrix} 0 & 0 & -a_3 \\ 1 & 0 & -a_2 \\ 0 & 1 & -a_1 \end{pmatrix} \in \mathfrak{g}$$

sending a characteristic polynomial to its *companion matrix*. The section to the Hitchin map that Hitchin constructed in [4] is not strictly the same as the one of [8] in the sense that he does not rely on the Kostant section but another section that feels more like a generalization of the companion matrix. Instead of the companion matrix, a map $\mathfrak{g} // G \rightarrow \mathfrak{g}$, we will construct a map $\mathfrak{g} // G \rightarrow [\mathfrak{g}/G]$, where $[\mathfrak{g}/G]$ is the quotient of \mathfrak{g} by the adjoint action of G in the sense of algebraic stack. This section will be called the companion section, which is free of any choice. The present note aims to explicitly construct the companion section for classical groups, including the symplectic and orthogonal groups and G_2 .

As applications of the companion sections, we will give elementary descriptions of affine Springer fibers and Hitchin fibers for classical groups similar to the description of the Hitchin fibers in the linear case

due to Beauville-Narasimhan-Ramanan. In addition to this elementary description of affine Springer fibers and Hitchin fibers for classical groups, using the companion section instead of the Kostant section allows us to save on the characteristic of the base field. Indeed, whereas the construction of the Kostant section requires the characteristic of the base field not to divide the order of the Weyl group, the existence of the companion section only requires that the characteristic of the base field is zero or greater than 2 for symplectic and orthogonal groups, and is zero or greater than 3 for the group G_2 .

The emphasis of this work is on providing case-by-case explicit formulas for the companion section for classical groups. It is also possible to construct the companion section uniformly. This will be the subject of our subsequent work.

2 Tensors defining classical groups

We will recall the standard definition of classical groups as the subgroup of the linear groups fixing certain tensors. This is very well known for symplectic and orthogonal groups but a bit less known for G_2 , which in a certain respect could qualify as a classical group as well.

Let V be a $2n$ -dimensional vector space over a base field k , V^* its dual vector space. The linear group $\mathrm{GL}(V)$ acts on the space $\wedge^2 V^*$ of alternating bilinear forms on V with an open orbit. An alternating bilinear form $\mu \in \wedge^2 V^*$ is considered non-degenerate if it lies in this open orbit. This is equivalent to requiring the induced map $\mu : V \rightarrow V^*$ to be an isomorphism. The stabilizer of such a non-degenerate alternating bilinear form is a symplectic group G . We note that $\mu \in \wedge^2 V^*$ is non-degenerate if $\wedge^n \mu \in \wedge^{2n} V^*$ is a non-zero vector of the 1-dimensional vector space $\wedge^{2n} V^*$ and as a result, G is contained in the special linear group $\mathrm{SL}(V)$. Then, a G -bundle over a k -scheme S consists of a locally free \mathcal{O}_S -module \mathcal{V} of rank $2n$ equipped with an alternating bilinear form $\wedge_S^2 \mathcal{V} \rightarrow \mathcal{O}_S$ which is non-degenerate fiberwise. We are more concerned with G -bundles than G itself, so choosing a specific non-degenerate alternating form $\mu \in \wedge^2 V^*$ is immaterial. We will write $G = \mathrm{Sp}_{2n}$.

Let V be a n -dimensional vector space over a base field k , V^* its dual vector space. The linear group $\mathrm{GL}(V)$ acts on the space $S^2 V^*$ of symmetric bilinear forms on V with an open orbit. A symmetric bilinear form $\mu \in S^2 V^*$ is considered non-degenerate if it lies in this open orbit. This is also equivalent to the induced map $\mu : V \rightarrow V^*$ being an isomorphism, which in turn is equivalent to the induced map $\wedge^n \mu : \wedge^n V \rightarrow \wedge^n V^*$ being an isomorphism of 1-dimensional vector spaces. We note that $\wedge^n V$ and $\wedge^n V^*$ are dual as vector spaces so that for every choice of a basis vector $\omega \in \wedge^n V$, we have a dual basis vector $\omega^* \in \wedge^n V^*$. A basis vector $\omega \in \wedge^n V$ is said to be compatible with μ if the equation $\wedge^n \mu(\omega) = \omega^*$ is satisfied. This equation has exactly two non-zero solutions $\omega \in \wedge^n V$, which differ by a sign. The stabilizer of a non-degenerate symmetric bilinear form $\mu \in S^2 V^*$ is an orthogonal group $O(\mu)$. The stabilizer of a pair (μ, ω) consisting of a non-degenerate symmetric bilinear form $\mu \in S^2 V^*$ and a compatible basis vector $\omega \in \wedge^n V$ is the special orthogonal group $\mathrm{SO}(\mu, \omega)$ which is the neutral component of $O(\mu)$. We note that $\mathrm{SO}(\mu, \omega) = O(\mu) \cap \mathrm{SL}(V)$ so that the special orthogonal group can also be defined as the stabilizer of a pair (μ, ω) as above but without requiring ω being compatible with μ . The stabilizer of any such pair is a special orthogonal group G . A G -bundle over a k -scheme S consists then in a locally free \mathcal{O}_S -module \mathcal{V} of rank n equipped a symmetric bilinear form $\wedge_S^2 \mathcal{V} \rightarrow \mathcal{O}_S$ which is non-degenerate fiberwise. As we are more concerned with G -bundles than G itself, choosing a specific non-degenerate symmetric form $\mu \in S^2 V^*$ is immaterial. We will write $G = \mathrm{SO}_n$.

There is a simple tensor definition of G_2 due to Engel [2]. Let V be a 7-dimensional vector space. The linear group $\mathrm{GL}(V)$ acts on the space $\wedge^3 V^*$ of non-degenerate trilinear forms on V with an open orbit. We will follow Hitchin's [5] in formulating the equation defining this open orbit. We will denote the contraction $\wedge^3 V^* \times V \rightarrow \wedge^2 V^*$ by μ_v for $\mu \in \wedge^3 V^*$ and $v \in V$. For $v_1, v_2 \in V$ and $\mu \in \wedge^3 V^*$, we then

have

$$\mu_{v_1} \wedge \mu_{v_2} \wedge \mu \in \wedge^3 V^*.$$

By choosing a non-zero vector ι of the determinant $\wedge^3 V$, μ gives rise to a symmetric bilinear form $\nu \in S^2 V^*$

$$\nu(v_1, v_2) = \langle \iota, \mu_{v_1} \wedge \mu_{v_2} \wedge \mu \rangle \quad (1)$$

which is non-degenerate if and only if μ lies in the open orbit of $\wedge^3 V^*$. We will say that μ is a non-degenerate 3-form on V . The stabilizer of a non-degenerate 3-form is a group $G_2 \times \mu_3(k)$ where $\mu_3(k)$ is the group of 3rd roots of unity in k ; We obtain the connected component, a group of type G_2 , by taking the intersection with $SL(V)$. A G_2 -bundle over a k -scheme S is thus a locally free \mathcal{O}_S -module \mathcal{V} of rank 7 equipped with an alternating trilinear form $\mu \in \wedge^3 \mathcal{V}^*$ which is non-degenerate fiberwise together with a trivialization of the determinant.

3 Spectral cover and the companion matrix

For all groups G discussed previously, including symplectic, special orthogonal, and G_2 , G is defined as a subgroup of GL_n fixing certain tensors. We call the inclusion $G \rightarrow GL_n$ the standard representation of G . We also have the induced inclusion of Lie algebras $\mathfrak{g} \rightarrow \mathfrak{gl}_n$ compatible with the adjoint actions of G and GL_n . We derive a morphism between the Chevalley invariant theoretic quotients

$$\mathfrak{c} = \mathfrak{g} // G \rightarrow \mathfrak{gl}_n // GL_n = \mathfrak{c}_n$$

which is a closed embedding for symplectic groups, odd special orthogonal groups, and G_2 , but not for even orthogonal groups. For GL_n , we have a spectral cover $\mathfrak{s}_n \rightarrow \mathfrak{c}_n$ which is a finite flat morphism of degree n so that $\mathcal{O}_{\mathfrak{s}_n}$ is a locally free $\mathcal{O}_{\mathfrak{c}_n}$ -module of rank n given with a canonical endomorphism $[x]$ which is the usual companion matrix. The main result of this work can be formulated as follows:

Theorem 3.1. *Let G be a symplectic group, odd special orthogonal group, or G_2 group and $G \rightarrow GL_n$ its standard representation. Let $\mathfrak{c} \rightarrow \mathfrak{c}_n$ be the induced map of Chevalley quotients which is a closed embedding in these cases. Then the restriction $\mathcal{O}_{\mathfrak{s}_n}$ to \mathfrak{c}*

$$\mathcal{V} = \mathcal{O}_{\mathfrak{c}} \otimes_{\mathcal{O}_{\mathfrak{c}_n}} \mathcal{O}_{\mathfrak{s}_n}$$

as locally free $\mathcal{O}_{\mathfrak{c}}$ -module affords canonical tensors so that it has a canonical G -reduction and the companion matrix for GL_n defines a canonical map $\mathfrak{g} // G \rightarrow [\mathfrak{g}/G]$ which is a section of the natural map $[\mathfrak{g}/G] \rightarrow \mathfrak{g} // G$. This statement remains valid for even orthogonal groups after replacing $\mathfrak{c} \times_{\mathfrak{c}_n} \mathfrak{s}_n$ by its normalization.

We prove the theorem by a case-by-case analysis. In particular, we will construct the explicit tensors required in each case.

3.1 Linear groups

We first recall how the companion matrix is connected to the universal spectral cover in the case GL_n . In this case, the Chevalley quotient $\mathfrak{g} // G$ is the n -dimensional affine space \mathbf{A}^n and the map $\chi : \mathfrak{g} \rightarrow \mathfrak{g} // G$ is given by the characteristic polynomial $\chi(\gamma) = (a_1(\gamma), \dots, a_n(\gamma))$ where $a_i(\gamma) = (-1)^i \text{tr}(\wedge^i \gamma)$. In this case we have $\mathfrak{c}_n = \text{Spec}(A_n)$ where $A_n = k[a_1, \dots, a_n]$. The spectral cover $\mathfrak{s}_n = \text{Spec}(B_n)$ where B_n is the A_n -algebra

$$B_n = A_n[x] / (x^n + a_1 x^{n-1} + \dots + a_n)$$

which is a free A_n -module of rank n as the images of $1, x, \dots, x^{n-1}$ form an A_n -basis of B_n . We also note that B_n is a regular k -algebra as it is isomorphic to the polynomial algebra of variables a_1, \dots, a_{n-1}, x . On the other hand, B_n is equipped with a A_n -linear operator $[x] : B_n \rightarrow B_n$ given by $b \mapsto bx$ which provides us with a A_n -point of $[\mathfrak{gl}_n/\mathrm{GL}_n]$. We have thus constructed a map $[x] : \mathfrak{c}_n \rightarrow [\mathfrak{gl}_n/\mathrm{GL}_n]$ which is a section to $\chi : [\mathfrak{gl}_n/\mathrm{GL}_n] \rightarrow \mathfrak{c}_n$. In term of matrices with respect to the A_n -basis of B_n given by $1, x, \dots, x^{n-1}$, $[x]$ is given by the usual companion matrix

$$x_\bullet = \begin{pmatrix} 0 & \cdots & -a_n \\ 1 & 0 & \cdots & -a_{n-1} \\ & \cdots & 0 & \\ 0 & & 1 & -a_1 \end{pmatrix} \in \mathfrak{gl}_n(A) \quad (2)$$

The companion matrix thus gives us a map $x_\bullet : \mathfrak{c} \rightarrow \mathfrak{g}$ in the case $G = \mathrm{GL}_n$ which is a section to the characteristic polynomial map. However, it is often more useful to think of $[x]$ as a map $[x] : \mathfrak{c} \rightarrow [\mathfrak{g}/G]$ in the case $G = \mathrm{GL}_n$.

There is an issue with using the companion matrix to construct a section to the Hitchin map as the companion map $x_\bullet : \mathfrak{c} \rightarrow \mathfrak{g}$ is not \mathbb{G}_m -equivariant. We note, however, that the stack-valued map $[x] : \mathfrak{c} \rightarrow [\mathfrak{g}/G]$ is almost \mathbb{G}_m -equivariant in the sense that after a base change by the isogeny $\mathbb{G}_m \rightarrow \mathbb{G}_m$ given by $t \mapsto t^2$, it becomes equivariant because of the identity

$$\mathrm{ad}(\mathrm{diag}(t^{n-1}, t^{n-3}, \dots, t^{1-n}))(\gamma) = t^{-2} \begin{pmatrix} 0 & 0 & \cdots & -t^{2n}a_n \\ 1 & 0 & \cdots & -t^{2n-2}a_{n-1} \\ & \cdots & & \\ 0 & & 1 & -t^2a_1 \end{pmatrix}. \quad (3)$$

This explains why we have a section to the Hitchin map after choosing a square root of the canonical bundle as in [4].

As we intend to use the companion matrix (2) to construct a canonical section to the Chevalley map $\chi : [\mathfrak{g}/G] \rightarrow \mathfrak{c}$ for classical groups, it is useful to further investigate the linear algebraic structure of B_n as an A_n -module. We have a symmetric A_n -bilinear map $\xi : B_n \otimes_{A_n} B_n \rightarrow A_n$ given by

$$\xi(b_1 \otimes_{A_n} b_2) = \mathrm{tr}_{B_n/A_n}(b_1 b_2)$$

thus an element $\xi \in S_{A_n}^2 B_n^*$. Because this element induces degenerate forms over the ramification locus of B_n over A_n , we need a correction term to get a symmetric bilinear form that is non-degenerate fiberwise.

This element defines a A_n -linear map also denoted by $\mu : B_n \rightarrow B_n^*$ where $B_n^* = \mathrm{Hom}_{A_n}(B_n, A_n)$. We note that the A_n -module B_n^* is naturally a B_n -module and $\mu : B_n \rightarrow B_n^*$ is B_n linear; thus, it is uniquely determined by the image of $1 \in B_n$ that we will also denote by $\mu \in B_n^*$. We will show that B_n^* is a free B_n -module of rank 1, construct a generator of B_n^* and find an explicit formula for $\mu \in B_n^*$ as a multiple of this generator.

Lemma 3.2. *Let us denote by v_0, \dots, v_{n-1} the basis of B_n given by the images of $1, x, \dots, x^{n-1}$ in B_n and v_0^*, \dots, v_{n-1}^* the dual basis of B_n^* . Then $\beta^* = v_{n-1}^*$ is a generator of B_n^* as a B_n -module. Let us denote $f' \in B_n = A_n[x]/(f)$ the image of the derivative*

$$nx^{n-1} + (n-1)a_1x^{n-2} + \cdots + a_{n-1} \in A_n[x]$$

of the universal polynomial $f = x^n + a_1x^{n-1} + \cdots + a_n \in A_n[x]$. Then we have $\mu = f'\beta^$.*

Proof. First, the discriminant d of the universal polynomial f , defined as the resultant between f and its derivative, is a nonzero element of the polynomial ring A_n . Indeed, d defines the ramification divisor of the finite flat covering $\mathfrak{s} \rightarrow \mathfrak{c}$, which is generically étale for there exist separable polynomials in $k'[x]$ of degree n with coefficients in any infinite field k' containing k . We denote $A' = A_n[d^{-1}]$ the localization of A_n obtained by inverting d , and $B' = B_n \otimes_{A_n} A'$. By construction, f' is an invertible element of B' . The trace map $\mathrm{tr}_{B'/A'} : B' \rightarrow A'$ of B' as free A' -module of rank n is now given by the Euler formula [9]

$$\mathrm{Tr} \left(\frac{x^k}{f'} \right) = \begin{cases} 0 & \text{if } k < n - 1 \\ 1 & \text{if } k = n - 1 \end{cases}$$

If v_0, \dots, v_{n-1} denote the basis of B' given by the images of $1, x, \dots, x^{n-1}$ in B_n and v_0^*, \dots, v_{n-1}^* the dual basis of $(B')^*$, then we derive from the Euler formula that the identities

$$\mu(v_i) = f' \left(v_{n-1-i}^* + \sum_{j < i} a'_{i,j} v_{n-1-j}^* \right) \quad (4)$$

hold in $B_n^* \otimes_{A_n} A'$ for some $a'_{i,j} \in A'$. In particular, we have $\mu(v_0) = f' v_{n-1}^*$. As the localization map $B_n^* \rightarrow B_n^* \otimes_{A_n} A'$ is injective, this identity also holds in B_n^* . It follows that $\mu = f' v_{n-1}^*$ as desired. \square

As a consequence, we have a canonical nondegenerate bilinear form $\beta^* : B_n \otimes_{A_n} B_n \rightarrow A_n$ which is symmetric with respect to which the A_n -linear operator $[x] : B_n \rightarrow B_n$ is self-adjoint. We will denote symbolically

$$\beta^* = f'^{-1} \mu.$$

Studying the \mathbb{G}_m action on the form β^* will also be useful in the sequel.

For $G = \mathrm{SL}_n$, the Lie algebra $\mathfrak{g} = \mathfrak{sl}_n$ is the space of traceless matrices. We have $\mathfrak{c} = \mathrm{Spec}(A)$ where $A = k[a_2, \dots, a_n]$. We note that for $a_1 = 0$, the companion matrix (2) is traceless and thus gives rise to a A -point on \mathfrak{sl}_n . The companion map $\gamma : \mathfrak{c} \rightarrow \mathfrak{g}$ induces a map $[\gamma] : \mathfrak{c} \rightarrow [\mathfrak{g}/G]$. The latter lays over the point of BG with values in A corresponding to the SL_n -bundle corresponding to rank n vector bundle B equipped with the trivialization of the determinant given by the basis $1, x, \dots, x^{n-1}$. The formula (3) shows that the map $[\gamma] : \mathfrak{c} \rightarrow [\mathfrak{g}/G]$ is equivariant with respect to the isogeny $\mathbb{G}_m \rightarrow \mathbb{G}_m$ given by $t \mapsto t^2$ for the diagonal matrix $\mathrm{diag}(t^{n-1}, t^{n-3}, \dots, t^{1-n})$ belonging to SL_n .

3.2 Symplectic groups

In the case $G = \mathrm{Sp}_{2n}$, we have $\mathfrak{c} = \mathrm{Spec}(A)$ with $A = k[a_2, \dots, a_{2n}]$. The spectral cover $\mathfrak{s} = \mathrm{Spec}(B)$ where

$$B = A[x]/(x^{2n} + a_2 x^{2n-2} + \dots + a_{2n})$$

is a free A -module of rank $2n$, is equipped with an involution $\tau : B \rightarrow B$ given $\tau(x) = -x$. The companion matrix (2) gives a A -linear endomorphism of B as a free A -module. For the companion matrix to produce a section to the Chevalley map $[\mathfrak{g}/G] \rightarrow \mathfrak{c}$ in the symplectic case, we need to construct a canonical nondegenerate symplectic form on the A -module B for which γ is anti-self-adjoint.

The standard representation $\mathrm{Sp}_{2n} \rightarrow \mathrm{GL}_{2n}$ induces a map on Chevalley bases $\mathfrak{c} \rightarrow \mathfrak{c}_{2n} = \mathrm{Spec}(A_{2n})$ where $A_{2n} = k[a_1, \dots, a_{2n}]$ which identifies \mathfrak{c} with the closed subscheme of \mathfrak{c}_{2n} defined by the ideal generated by $a_1, a_3, \dots, a_{2n-1}$. We have $B = A \otimes_{A_{2n}} B_{2n}$ where B_{2n} is the finite free A_{2n} -algebra defining the spectral covering of \mathfrak{c}_{2n} . If we denote $B^* = \mathrm{Hom}_A(B, A)$ then we have $B^* = A \otimes_{A_{2n}} B_{2n}^*$ where $B_{2n}^* = \mathrm{Hom}_{A_{2n}}(B_{2n}, A_{2n})$. The generator β_{2n}^* of the free B_{2n} -module B_{2n}^* defined in Lemma

3.2 then induces a generator β^* of B^* as a free B -module of rank one which can also be viewed as the bilinear form $\beta^* : B \otimes_A B \rightarrow A$ given by $b_1 \otimes_A b_2 = \text{tr}_{B/A}(f'^{-1}b_1b_2)$ after localization.

The bilinear form $\omega : B \otimes_A B \rightarrow A$

$$\omega(b_1, b_2) = \beta^*(b_1, \tau(b_2)) = \text{tr}_{B/A}(f'^{-1}b_1\tau(b_2))$$

with the second identity only making sense after localization of A making f' invertible, is a non-degenerate symplectic form for which $[x]$ is anti-self-adjoint. Indeed, we have

$$\omega(b_1, b_2) = -\omega(b_2, b_1)$$

because $\tau(f') = -f'$ for $f' \in A[x]$ is an odd polynomial as $f \in A[x]$ is an even polynomial. The equation $\omega(xb_1, b_2) + \omega(b_1, xb_2) = 0$ can be derived from $\tau(x) = -x$.

It follows that we have a morphism

$$[x] : \mathfrak{c} \rightarrow [\mathfrak{g}/G]$$

which deserves to be called the companion map for the symplectic group. To obtain a companion matrix $x_\bullet : \mathfrak{c} \rightarrow \mathfrak{g}$, it is enough to find a trivialization of the G -bundle associated with the non-degenerate symplectic form $\omega : B \otimes_A B \rightarrow A$. For most applications, particularly the Hitchin fibration, we only need the section $[x] : \mathfrak{c} \rightarrow [\mathfrak{g}/G]$.

3.3 Odd special orthogonal groups

In the case $G = \text{SO}_{2n+1}$, we have $\mathfrak{c} = \text{Spec}(A)$ with $A = k[a_2, a_4, \dots, a_{2n}]$. The spectral cover is defined as $\mathfrak{s} = \text{Spec}(B)$ where $B = A[x]/(f)$ with $f = xf_0$ and $f_0 = x^{2n} + a_2x^{2n-2} + \dots + a_{2n}$. B is a free A -module of rank $2n + 1$. As in the symplectic case, we will define a symmetric non-degenerate bilinear form $B \otimes_A B \rightarrow A$ for which the multiplication by x is anti-self-adjoint.

The standard representation $\text{SO}_{2n+1} \rightarrow \text{GL}_{2n+1}$ gives rise to a map $\mathfrak{c} \rightarrow \mathfrak{c}_{2n+1} = \text{Spec}(k[a_1, \dots, a_{2n+1}])$ which is a closed embedding defined by the ideal generated by $a_1, a_3, \dots, a_{2n+1}$. We have $B = A \otimes_{A_{2n+1}} B_{2n+1}$ where B_{2n+1} is the finite free A_{2n+1} -module of rank $2n + 1$ defining the spectral cover in the case GL_{2n+1} . We also have $B^* = A \otimes_{A_{2n+1}} B_{2n+1}^*$ where $B^* = \text{Hom}_A(B, A)$ and $B_{2n+1}^* = \text{Hom}_{A_{2n+1}}(B_{2n+1}, A_{2n+1})$. Following the discussion in the linear case B_{2n+1}^* is a free B_{2n+1} generated by the element $\beta_{2n+1}^* = (f')^{-1}\mu$ where μ is the trace form $\mu(b_1 \otimes_A b_2) = \text{tr}(b_1b_2)$. It induces a generator β^* of B^* as a B -module. We define the bilinear form $\omega : B \otimes_A B \rightarrow A$ by

$$\omega(b_1, b_2) = \beta^*(b_1, \tau(b_2)) = \text{tr}(f'^{-1}b_1\tau(b_2)). \quad (5)$$

The form bilinear form ω is a nondegenerate bilinear form because β^* is. It is symmetric because $\tau(f') = f'$ as f' is an even polynomial. The equation $\omega(xb_1, b_2) + \omega(b_1, xb_2)$ can be derived from the fact $\tau(x) = -x$.

By choosing a trivialization of the determinant, we obtain a companion map $[x] : \mathfrak{c} \rightarrow [\mathfrak{g}/G]$ for $G = \text{SO}_{2n+1}$.

3.4 Even special orthogonal groups

The case $G = \text{SO}_{2n}$ is slightly more difficult for the map $\mathfrak{c} \rightarrow \mathfrak{c}_{2n}$ induced by the standard representation of SO_{2n} is not a closed embedding. Indeed, we have $\mathfrak{c}_{2n} = \text{Spec}(A_{2n})$ where $A_{2n} = k[a_1, \dots, a_{2n}]$ but

$\mathfrak{c} = \text{Spec}(A)$ where $A = k[a_2, \dots, a_{2n-2}, p_n]$ where p_n is the Pfaffian satisfying $p_n^2 = a_{2n}$ does not lie in the image of $A_{2n} \rightarrow A$. If B_{2n} is the spectral cover of A_{2n} and $B = A \otimes_{A_{2n}} B_{2n}$ then we have

$$B = A[x]/(x^{2n} + a_2 x^{2n-2} + \dots + a_{2n-2} x^2 + p_n^2).$$

As indicated by Hitchin [4], the true spectral cover for even special orthogonal groups is not B but its blowup \tilde{B} along the singular locus defined by x . We have

$$\tilde{B} = A[x, p_{n-1}]/(xp_{n-1} - p_n, x^{2n-2} + a_2 x^{2n-4} + \dots + a_{2n-2} x^2 + p_{n-1}^2)$$

which is a free A -module of rank $2n$ and smooth as a k -algebra. We have an involution τ on B and \tilde{B} given by $\tau(x) = -x$ and $\tau(p_{n-1}) = -p_{n-1}$.

The dualizing sheaf $\omega_{\tilde{B}/A}$ is a free rank-one \tilde{B} -module, canonically isomorphic to \tilde{B} away from the ramification locus. As a \tilde{B} -submodule of $\text{Fr}(\tilde{B})$ it is generated by the inverse of the different $\mathfrak{D}_{\tilde{B}/A}$ which is given by the formula

$$\begin{aligned} \mathfrak{D}_{\tilde{B}/A} &= \det \begin{pmatrix} -p_{n-1} & f' \\ -x & 2p_{n-1} \end{pmatrix} \\ &= (n-1)x^{2(n-1)} + (n-2)a_2 x^{2(n-2)} + \dots + a_{2n-2} x^2 + p_{n-1}^2. \end{aligned}$$

In other words, the bilinear form $\tilde{B} \otimes_A \tilde{B} \rightarrow A$ given by

$$b_1 \otimes_A b_2 \mapsto \text{tr}_{\tilde{B}/A}(\mathfrak{D}_{\tilde{B}/A}^{-1} b_1 b_2)$$

is non-degenerate. As in the symplectic and odd special orthogonal cases, we now consider the sesquilinear bilinear form

$$\omega(b_1, b_2) = \text{tr}_{\tilde{B}/A}(\mathfrak{D}_{\tilde{B}/A}^{-1} b_1 \tau(b_2))$$

Then ω is a non-degenerate symmetric bilinear form because $\tau(\mathfrak{D}_{\tilde{B}/A}) = \mathfrak{D}_{\tilde{B}/A}$, and it satisfies

$$\omega(xb_1, b_2) = -\omega(b_1, xb_2).$$

After a choice of trivialization of the determinant of \tilde{B} as a free A -module of rank n , the multiplication by x gives rise to the companion section $[x] : \mathfrak{g} // G \rightarrow [\mathfrak{g}/G]$ for the odd special orthogonal group $G = \text{SO}_{2n+1}$.

3.5 The group G_2

In the case G_2 , the invariant quotient is $A = k[e, q]$ with $\deg(e) = 2$ and $\deg(q) = 6$. The spectral cover $\mathfrak{s} = \text{Spec}(B)$ of $\mathfrak{c} = \text{Spec}(A)$ given by

$$B = A[x]/(xf_0) \quad \text{for } f_0 = x^6 - ex^4 + \frac{e^2}{4}x^2 + q$$

is a reducible cover of A with two components corresponding to the quotient maps

$$B \rightarrow B' = A[x]/(f_0) \quad \text{and} \quad B \rightarrow A = A[x]/(x).$$

The cover $\mathfrak{s}' = \text{Spec}(B')$ of A is finite, flat of degree 6, and factors through two subcovers, of degrees 2 and 3, corresponding to the sub- A -algebras

$$A \subset A[y]/\left(y^3 - ey^2 + \frac{e^2}{4}y + q\right) \subset B' \quad \text{where } y = x^2$$

$$A \subset A[z]/(z^2 + q) \subset B' \quad \text{where } z = x \left(x^2 - \frac{e}{2} \right)$$

Let $\epsilon \in B[q^{-1}]^* := \text{Hom}_{A[q^{-1}]}(B[q^{-1}], A[q^{-1}])$ be dual to f_0 ; $\delta_i \in B[q^{-1}]^*$ be dual to x^i ; and $\eta_i \in B[q^{-1}]^*$ be dual to $x^i z$ for $i = 1, 2, 3$. Let tr_z denote the skew-symmetric bilinear form on B given by

$$\text{tr}_z(g, h) = \text{Tr}_{\text{Frac}(B)/\text{Frac}(A)} \left(\frac{g(x)h(-x)z}{f(x)} \right)$$

We will denote by ρ the 3-form on $B[q^{-1}]$ given by

$$\rho := \delta_1 \wedge \delta_2 \wedge \eta_3 + \delta_1 \wedge \eta_2 \wedge \delta_3 + \eta_1 \wedge \delta_2 \wedge \delta_3 - q \cdot \eta_0 \wedge \eta_1 \wedge \eta_2 + \epsilon \wedge \text{tr}_z \quad (6)$$

A priori, the 3-form above is valued in $A[q^{-1}]$. The next proposition tells us that it restricts to an element of $\bigwedge_A^3 B^*$.

Proposition 3.3. *Restricting the 3-form ρ to $B \rightarrow B[q^{-1}]$ induces a 3-form $\rho \in \bigwedge_A^3 B^*$. In other words, ρ takes values in A when restricted to B .*

Proof. Consider the A -basis of B given by

$$\{1, x^i, x^i z : i = 1, 2, 3\}.$$

This differs from the $A[q^{-1}]$ -basis

$$\{f_0, x^i, x^i z : i = 1, 2, 3\}$$

of $B[q^{-1}]$ only by scaling f_0 . As ρ is valued in A on the A -linear span of the latter basis, it suffices to check the contraction $\iota_1 \rho$ of ρ along $1 \in B$ is valued in A . We compute

$$\begin{aligned} \iota_1 \rho &= \eta_1 \wedge \eta_2 - \frac{e}{2} \eta_2 \wedge \eta_3 + \frac{1}{q} (\text{tr}_z - \delta_1 \wedge \delta_2 + \frac{e}{2} \delta_2 \wedge \delta_3 - i_{x^3 z} \text{tr}_z \wedge \epsilon + \frac{e}{2} i_{x z} \text{tr}_z \wedge \epsilon) \\ &= \eta_1 \wedge \eta_2 - \frac{e}{2} \eta_2 \wedge \eta_3 + \left[\frac{1}{q} (\text{tr}_z - \delta_1 \wedge \delta_2 + \frac{e}{2} \delta_2 \wedge \delta_3) - \iota_1 \text{tr}_z \wedge \epsilon \right] \end{aligned}$$

Rewriting the latter in terms of a dual basis ξ_i , $i = 0, \dots, 6$ for the A -basis $\{x^i : i = 0, \dots, 6\}$ of B , we see that the expression in square brackets above is

$$\iota_1 \rho = e_3 \wedge e_6 + e_4 \wedge e_5 - \frac{3e}{2} e_5 \wedge e_6$$

whose image lies in A . □

As the previous proposition illustrates, working with the form ρ requires significantly more computational effort. As such, the latter two results will be checked primarily with computer algebra packages. These computations were done in Macaulay2; explicit code for each calculation is referred to in Appendix A.

Proposition 3.4. *Let ν be the bilinear form associated to ρ as in equation (1) and let $\omega \in S_A^2 B^*$ be the symmetric, nondegenerate form given by the formula (5). Then, $\nu = -2^4 3^2 \omega$.*

Proposition 3.5. *The form ρ is compatible with the endomorphism $[x]$, in the sense that*

$$\rho(xb_1, b_2, b_3) + \rho(b_1, xb_2, b_3) + \rho(b_1, b_2, xb_3) = 0.$$

As such, the form ρ together with a trivialization of the determinant gives a map $[x]: \mathfrak{c} \rightarrow [\mathfrak{g}_2/G_2]$.

4 Special components

In the previous section, we gave explicit formulas for the tensors defining the reduction of the vector bundle $\mathcal{O}_{c \times_{c_n} s_n}$ to G so that the companion section for GL_n induces the companion section for classical group G . These explicit formulas may feel like miracles, especially in the G_2 case where a computer algebra system is needed. In this section, we will derive them from the geometry of spectral covers, which makes the construction more conceptual, especially in the G_2 case. In subsequent work, we use this approach to construct the companion section uniformly.

4.1 Special form and component associated with a subcover

Let A be a k -algebra, B a finite flat A -algebra of degree n generated by one element $b \in B$, and $A' \subset B$ an A -subalgebra of B such that A' is finite flat of degree m over A generated by one element $a' \in A'$ and B is a finite flat A' -algebra of degree d generated by b . Under these assumptions, we have $B \simeq A[x]/P(x)$ where $P(x)$ is the characteristic polynomial of the A -linear $b : B \rightarrow B$ defined as the multiplication by b . Similarly we have $A' \simeq A[x]/(P_1(x))$ where $P_1(x)$ is the characteristic polynomial of the A -linear operator $a' : A' \rightarrow A'$, and $B \simeq A'[x]/P_2(x)$ where $P_2(x)$ is the characteristic polynomial of the A' -linear operator $b : B \rightarrow B$.

Assuming that the characteristic of k is greater than d , we want to construct an alternating d -form

$$\omega_{A'} : \wedge_A^d B \rightarrow A$$

supported on a special component of $\mathrm{Spec}(S_A^d B)$ isomorphic to $\mathrm{Spec}(A')$. We explain what this means. As far as we know, the concept of non-degeneracy for d -forms is not yet defined for $d \geq 3$ and thus we can prove it only for $d = 1$ or $d = 2$. However, we expect that the form we construct is non-degenerate for a reasonable definition of this concept. As to the special component, $\wedge_A^d B$ is a module over the ring of symmetric tensors $(\otimes_A^d B)^{\mathfrak{S}_d}$. We will construct a surjective homomorphism of A -algebras $(\otimes_A^d B)^{\mathfrak{S}_d} \rightarrow A'$ which realizes $\mathrm{Spec}(A')$ as an irreducible component of $\mathrm{Spec}((\otimes_A^d B)^{\mathfrak{S}_d})$ if B is generically étale over A and A' is a domain.

The homomorphism of A -algebras $(\otimes_A^d B)^{\mathfrak{S}_d} \rightarrow A'$ is constructed as follows. Let $P_2(x) = x^d + a'_1 x^{d-1} + \dots + a'_d$ be the characteristic polynomial of the A' -linear map $b : B \rightarrow B$. Then we have

$$B = A'[x]/(x^d + a'_1 x^{d-1} + \dots + a'_d).$$

We consider the polynomial ring $R = k[x_1, \dots, x_d]$ and the subring S of invariant polynomials under the symmetric group \mathfrak{S}_d . We have

$$S = k[x_1, \dots, x_d]^{\mathfrak{S}_d} = k[\alpha_1, \dots, \alpha_d]$$

with

$$\alpha_i = (-1)^i \sum_{1 \leq j_1 < \dots < j_i \leq d} \alpha_{j_1} \dots \alpha_{j_i}.$$

Since R and S are regular, and R is a finite generated S -module, R is a finite flat S -algebra of degree $d!$. We consider the homomorphism of algebras $S \rightarrow A'$ given by $\alpha_i \mapsto a'_i$ and the base change $A' \otimes_S R$ which is a finite flat A' -algebra of degree $d!$ equipped with an action of \mathfrak{S}_d . We have $(A' \otimes_S R)^{\mathfrak{S}_d} = A'$. Moreover, for every $i \in \{1, \dots, d\}$ we have a homomorphism of A' -algebras $B \rightarrow R \otimes_S A'$ given by $x \mapsto x_i$ which together give rise to a surjective homomorphism of A' -algebras $\otimes_{A'}^d B \rightarrow A' \otimes_S R$, which is \mathfrak{S}_d -equivariant. We derive a \mathfrak{S}_d -equivariant surjective homomorphism of A -algebras

$$\bigotimes_A^d B \rightarrow \bigotimes_{A'}^d B \rightarrow A' \otimes_S R. \quad (7)$$

By taking the \mathfrak{S}_d -invariant, we obtain the desired homomorphism of algebras

$$S_A^d B = \left(\bigotimes_A^d B \right)^{\mathfrak{S}_d} \rightarrow A',$$

which is surjective because taking \mathfrak{S}_d -invariants is an exact functor under the characteristic assumption.

We will now construct a special d -form on B

$$\omega_{A'} : \bigwedge_A^d B \rightarrow A$$

supported on the special component. As above, we have a surjective homomorphism of algebras \mathfrak{S}_d -equivariant surjective homomorphisms of A -algebras $\bigotimes_A^d B \rightarrow \bigotimes_{A'}^d B \rightarrow A' \otimes_S R$ which induces a surjective A -linear maps of the alternating parts $\bigwedge_A^d B \rightarrow \bigwedge_{A'}^d B \rightarrow A' \otimes_S R^{\text{sgn}}$ where R^{sgn} is the direct factor of R as S -module in which \mathfrak{S}_d acts as the sign character. It is known that R^{sgn} is a free S -module generated by $\prod_{1 \leq i < j \leq d} (x_i - x_j)$. We thus obtains a surjective A -linear map $\bigwedge_A^d B \rightarrow A'$. By composing it with the generator of $\text{Hom}_A(A', A)$ constructed in 3.2 we obtain the special d -form $\omega_{A'} : \bigwedge_A^d B \rightarrow A$ which is supported by the special by construction.

Let us discuss the non-degeneracy of the special d -form $\omega_{A'} : \bigwedge_A^d B \rightarrow A$. For $d = 1$, this follows from lemma 3.2. We can check that it is also non-degenerate everywhere for $d = 2$. For $d \geq 3$, we don't know a general definition of non-degeneracy but it easy to see that the special form $\omega_{A'}$ is everywhere non-zero. In dimension 6 and 7 where the definition of non-degeneracy is available, we will check that the special d -form is everywhere non-degenerate by direct calculation.

4.2 Sp_{2n} case

We recall in the case $G = \text{Sp}_{2n}$, we have $\mathfrak{c} = \text{Spec}(A)$ with $A = k[a_2, \dots, a_{2n}]$. The spectral cover $\mathfrak{s} = \text{Spec}(B)$ where

$$B = A[x]/(x^{2n} + a_2 x^{2n-2} + \dots + a_{2n})$$

is a free A -module of rank $2n$, is equipped with an involution $\tau : B \rightarrow B$ given $\tau(x) = -x$. We consider the subalgebra A' of B consisting of elements fixed under τ

$$A' = A[y]/(y^n + a_2 y^{n-1} + \dots + a_{2n}).$$

We then have $B = A'[x]/(x^2 - y)$.

The construction of the special form and special component in 4.1 gives rise to an alternating form

$$\omega_{A'} : \bigwedge_A^2 B \rightarrow A$$

supported in the special component $\mathfrak{c}' = \text{Spec}(A')$ of $(\mathfrak{s} \times_{\mathfrak{c}} \mathfrak{s}) // \mathfrak{S}_2$ where $\mathfrak{s} = \text{Spec}(B)$ and $\mathfrak{c} = \text{Spec}(A)$. The homomorphism (7) $\text{Sym}_A^2(B) \rightarrow A'$ can be explicitly computed elements of the form:

$$b \otimes_A 1 + 1 \otimes_A b \mapsto \text{tr}_{B/A'}(b).$$

In particular, $x \otimes_A 1 + 1 \otimes_A x$ be long to the kernel of $\text{Sym}_A^2(B) \rightarrow A'$, and in fact on can verify that it is a generator of the kernel. Since $x \otimes_A 1 + 1 \otimes_A x$ annihilates $\omega_{A'}$ we have

$$\omega_{A'}(x b_1, b_2) + \omega_{A'}(b_1, x b_2) = 0$$

for every $b_1, b_2 \in B$. By lemma ??, the 2-form $\omega_{A'}$ is everywhere non-degenerate. We can also see by explicit calculation that the form $\omega_{A'}$ is the same as the 2-form we constructed in subsection 3.2 by means of the Euler formula.

4.3 G_2 case

In the case G_2 , the invariant quotient is $A = k[e, q]$ with $\deg(e) = 2$ and $\deg(q) = 6$. The spectral cover $\mathfrak{s} = \text{Spec}(B)$ with

$$B = A[x]/(xf_0) \quad \text{for } f_0 = x^6 - ex^4 + \frac{e^2}{4}x^2 + q$$

is a reducible cover of A with two components corresponding to the quotient maps

$$B \rightarrow B' = A[x]/(f_0) \quad \text{and} \quad B \rightarrow A$$

We will define a canonical 3-form on B out of a 3-form and a 2-form on B' associated to subalgebras

$$\begin{aligned} A \subset A' = A[z]/(z^2 + q) = k[e, y] &\subset B' = A'[x]/(x^3 - \frac{e}{2}x - z) \\ A \subset A'' = A[y]/\left(y^3 - ey^2 + \frac{e^2}{4}y + q\right) = k[e, z] &\subset B' = A''[x]/(x^2 - y). \end{aligned}$$

Since both A' and A'' are regular algebras, they are finite flat A -modules of rank 2 and 3, respectively, whereas B are finite flat A' -module and A'' -module of rank 3 and 2, respectively. The construction of the special form associated with a subcover gives rise to

$$\omega_{A'} : \wedge_A^3 B' \rightarrow A \quad \text{and} \quad \omega_{A''} : \wedge_A^2 B' \rightarrow A$$

supported on the special components $\mathfrak{c}' = \text{Spec}(A')$ and $\mathfrak{c}'' = \text{Spec}(A'')$ of $\mathfrak{s}^{\times 3} // \mathfrak{S}_3$ and $\mathfrak{s}^{\times 2} // \mathfrak{S}_2$, respectively. By arguing as in the symplectic case, we see that $\omega_{A'}$ is annihilated by $x \otimes_A 1 \otimes_A 1 + 1 \otimes_A x \otimes_A 1 + 1 \otimes_A 1 \otimes_A x$ and $\omega_{A''}$ by $x \otimes_A 1 + 1 \otimes_A x$. It follows that as alternating forms, they satisfy the relations:

$$\begin{aligned} \omega_{A'}(xb_1, b_2, b_3) + \omega_{A'}(b_1, xb_2, b_3) + \omega_{A'}(b_1, b_2, xb_3) &= 0 \\ \omega_{A''}(xb_1, b_2) + \omega_{A''}(b_1, xb_2) &= 0 \end{aligned}$$

for all $b_1, b_2, b_3 \in B$.

The form $\omega_{A'}$ agrees with the restriction of the form ρ calculated by Macaulay 2 when restricted to $B' \rightarrow B$, with the inclusion given by multiplication by x : Indeed, the restriction of ρ takes value 1 on each of:

$$z \wedge x \wedge x^2, \quad x \wedge zx \wedge x^2, \quad 1 \wedge x \wedge zx^2$$

and $-q$ on $z \wedge zx \wedge zx^2$. This exactly detects the coefficient of z when these wedges are written in terms of the A' basis $1 \wedge x \wedge x^2$ for $\wedge_A^3 B'$, which matches $\omega_{A'}$ since the generator of $\text{Hom}_A(A', A)$ as an A'' module detects the coefficient of A' .

We now build a 3-form on B out of the 3-form $\omega_{A'}$ and 2-form $\omega_{A''}$ on B' . Since $B = A[x]/(xf_0)$, $B' = A[x]/(f_0)$ we have exact sequences of free A -modules

$$0 \rightarrow A \rightarrow B \rightarrow B' \rightarrow 0 \quad \text{and} \quad 0 \rightarrow B' \rightarrow B \rightarrow A \rightarrow 0$$

where the map $A \rightarrow B$ in the first sequence is given by $1 \mapsto f_0$ and the map $B' \rightarrow B$ in the second sequence is given by $1 \mapsto x$. It follows an exact sequences

$$0 \rightarrow A \oplus B' \rightarrow B \rightarrow Q \rightarrow 0 \quad \text{and} \quad 0 \rightarrow B \rightarrow A \oplus B' \rightarrow Q \rightarrow 0$$

where $Q = A/(q) = B'/(x)$. It follows an exact sequence

$$0 \rightarrow \wedge_A^3 B^* \rightarrow \wedge_A^3 (B')^* \oplus \wedge_A^2 (B')^* \rightarrow \wedge^2 (B')^*/(q) \rightarrow 0$$

where the map $\wedge_A^2 (B')^* \rightarrow \wedge^2 (B')^*/(q)$ is the reduction modulo q , and the map $\wedge_A^3 (B')^* \rightarrow \wedge^2 (B')^*/(q)$ is obtained by the composition

$$\wedge_A^3 (B')^* \rightarrow \wedge_A^3 B^* \rightarrow \wedge_A^2 B^* \rightarrow \wedge_A^2 (B')^* \rightarrow \wedge_A^2 (B')^*/(q)$$

where the first map is induced by the projection $B \rightarrow B'$, the second is given by contraction with f_0 , the third map is induced by the inclusion $B' \rightarrow B$ sending $1 \mapsto x$, and the final map is the quotient map. Since $q \wedge^2 (B')^* \simeq \wedge^2 (B')^*$ is a free, rank 1 module over the special component of $S_A^2(B')$, there is a unique generator as an A'' module. The 3-form $\omega_{A'}$ and the 2-form $\omega_{A''}$ do not have the same image in $\wedge_A^2 (B')^*/(q)$; however, the form $z\omega_{A''}$ is and it gives a generator for the A'' submodule of 2-forms compatible with $\omega_{A'}$. The pair $(\omega_{A'}, z\omega_{A''})$ comes from an element of $\wedge_A^3 B^*$ which agrees with the 3-form calculated by Macaulay2.

5 Lattice description of affine Springer fibers of classical groups

Let us recall Kazhdan-Lusztig's definition [6] of affine Springer fibers. Let G be a split reductive group defined over a field k and \mathfrak{g} its Lie algebra. Let $F = k((\varpi))$ the field of Laurent formal series and $\mathcal{O} = k[[\varpi]]$ its ring of integers. Let $\gamma \in \mathfrak{g}(F)$ be a regular semisimple element. The affine Springer fiber associated with γ is an ind-scheme defined over k whose set of k -points is

$$\mathcal{M}_\gamma(k) = \{g \in G(F)/G(\mathcal{O}) \mid \text{ad}(g)^{-1}\gamma \in \mathfrak{g}(\mathcal{O})\}.$$

We note that \mathcal{M}_γ is non-empty only if the image $a \in \mathfrak{c}(F)$ lies in $\mathfrak{c}(\mathcal{O})$ where $\mathfrak{c} = \mathfrak{g} // G$ is the invariant theoretic quotient of \mathfrak{g} by the adjoint action of G . As argued in [8], using the Kostant section, we can define an affine Springer fiber \mathcal{M}_a depending only on a instead of γ , which is isomorphic to \mathcal{M}_γ .

For $G = \text{GL}_n$, the affine Springer fiber \mathcal{M}_a has a well-known lattice description. In this case, $\mathfrak{c} = \mathbb{A}^n$. If $a = (a_1, \dots, a_n) \in \mathcal{O}_n$, we form the finite flat \mathcal{O} -algebra

$$B_a = \mathcal{O}[x]/(f_a)$$

where $f_a = x^n + a_1 x^{n-1} + \dots + a_n$ by the base change from the universal spectral cover. As $\gamma \in \mathfrak{g}(F)$ is a regular semisimple element, $B_a \otimes_{\mathcal{O}} F$ is finite and étale over F . We have a well-known lattice description of the affine Springer fiber \mathcal{M}_a in this case. The set $\mathcal{M}_a(k)$ consists then of lattices \mathcal{V} in the n -dimensional vector space $V = B_a \otimes F$ which are also B_a -modules.

For computational purposes, it is desirable to have a lattice description of affine Springer fibers for classical groups, which is as simple as in the linear case. This is possible due to the construction of the companion matrix, and in fact, this was our original motivation.

In the cases we have investigated in the paper, i.e., symplectic, special orthogonal, and G_2 , we have constructed a finite, flat spectral cover $\mathfrak{s} = \text{Spec}(B)$ of the invariant theoretic quotient $\mathfrak{c} = \text{Spec}(A)$ which is étale over the regular semisimple locus of \mathfrak{c} . The degree $d = \deg(B/A)$ is the degree of the standard representation which is $2n$ for Sp_{2n} , $2n+1$ for SO_{2n+1} , $2n$ for SO_{2n} and 7 for G_2 . In the case $\text{SO}(2n)$, we must consider the normalization \tilde{B} of B . In each of these cases, we constructed a form ω , which is

- a non-degenerate symplectic form $\omega : B \times B \rightarrow A$ satisfying $\omega(xb_1, b_2) + \omega(b_1, xb_2) = 0$ for Sp_{2n}

- a non-degenerate symmetric form $\omega : B \times B \rightarrow A$ satisfying $\omega(xb_1, b_2) + \omega(b_1, xb_2) = 0$ for SO_{2n+1}
- a non-degenerate symmetric form $\omega : \tilde{B} \times \tilde{B} \rightarrow A$ satisfying $\omega(xb_1, b_2) + \omega(b_1, xb_2) = 0$ for SO_{2n}
- a non-degenerate alternating form $\omega : B \times B \times B \rightarrow A$ satisfying

$$\omega(xb_1, b_2, b_3) + \omega(b_1, xb_2, b_3) + \omega(b_1, b_2, xb_3) = 0$$

for G_2

We also constructed a trivialization of the determinant $\bigwedge_A^d B = A$ in all these cases.

For every $a \in \mathfrak{c}(\mathcal{O}) \cap \mathfrak{c}^{rs}(F)$, we construct a finite flat \mathcal{O} -algebra B_a by base change from the spectral cover $\mathfrak{s} \rightarrow \mathfrak{c}$. Because $a \in \mathfrak{c}^{rs}(F)$, the generic fiber $V_a = B_a \otimes_{\mathcal{O}} F$ is a finite étale F -algebra of degree d . By pulling back ω , we get a form ω_a which is a non-degenerate alternating F -bilinear form on V_a in the symplectic case, a non-degenerate symmetric F -bilinear form on V_a in the orthogonal case, and a non-degenerate alternating F -trilinear form on V_a in the G_2 case. Moreover, it extends to a non-degenerate form valued in \mathcal{O} on B_a in Sp_{2n} , SO_{2n+1} and G_2 cases and on \tilde{B}_a in the SO_{2n} -case.

Theorem 5.1. *The set of k -points of the affine Springer fiber \mathcal{M}_a is the set of \mathcal{O} -lattices \mathcal{V} of V_a , which are B_a -modules, such that the restriction of ω_a has value in \mathcal{O} and such that $\deg(\mathcal{V} : B_a) = 0$ in Sp_{2n} , SO_{2n+1} , G_2 cases and $\deg(\mathcal{V} : \tilde{B}_a) = 0$ in the SO_{2n} case.*

6 Application to the Hitchin fibration

Let X be a smooth, projective curve over an algebraically closed field k and let G be a reductive group over k with Lie algebra \mathfrak{g} . Fix a line bundle L on X such that either $\deg(L) > 2g - 2$ or $L = K$ is the canonical bundle. Hitchin [4] introduced the moduli space \mathcal{M} on X to be the moduli stack of Higgs bundles whose k points are given by the set of Higgs bundles

$$\mathcal{M}(k) = \{(E, \phi) : E \rightarrow X \text{ is a } G \text{ bundle, } \phi \in \Gamma(X, \mathrm{ad}(E) \otimes L)\}$$

More succinctly, \mathcal{M} is the mapping stack $\mathcal{M} = \mathrm{Maps}(X, [\mathfrak{g}_L/G])$ where $\mathfrak{g}_L = \mathfrak{g} \wedge^{\mathbb{G}_m} L$ is the twisted bundle of Lie algebras on X .

Recall that under mild hypotheses on the characteristic of k ($\mathrm{char}(k) > 2$ for $G = \mathrm{SO}_n$ and Sp_{2n} and $\mathrm{char}(k) > 3$ for $G = G_2$), the Chevalley isomorphism shows

$$\mathfrak{g} // G \simeq \mathfrak{t} // W \simeq \mathbb{A}^n$$

is an affine space with \mathbb{G}_m action by weights d_1, \dots, d_n . Let

$$\mathcal{A} = \mathrm{Maps}(X, \mathfrak{g}_L // G) \simeq \otimes_{i=1}^n \Gamma(X, L^{\otimes d_i})$$

Hitchin studied the space \mathcal{M} through the fibration that now bears his name:

$$h : \mathcal{M} \rightarrow \mathcal{A}, \quad (E, \phi) \mapsto \mathrm{char}(\phi)$$

where $\mathrm{char}(\phi)$ is given by composition with the quotient map $[\mathfrak{g}/G] \rightarrow \mathfrak{g} // G$. Let \mathcal{M}_a denote the fiber of the map h over a point $a \in \mathcal{A}$. In the case that $G = \mathrm{GL}_n$, $d_i = i$ and $\mathrm{char}(\phi) = \sum_i a_i x^i$ is the characteristic polynomial of ϕ , whose coefficients are then sections $a_i \in \Gamma(X, L^{\otimes i})$.

The companion section $[x] : \mathfrak{g} // G \rightarrow [\mathfrak{g}/G]$ can be used to construct an explicit section to the Hitchin map after extracting a square root of L . This section in many cases is almost the same as the section

constructed by Hitchin [4] and [3], but can be different from the section constructed in [8] which is based on the Kostant section. In every case, the Higgs bundle constructed from the companion section will be built out of the structural sheaf of the spectral curve. Note that the following assumes basic \mathbb{G}_m equivariance properties of the relevant forms. For example, in the case of $G = \mathrm{Sp}_{2n}$, we have constructed a canonical alternating form $\omega: \wedge_A^2 B \rightarrow A$ which satisfies $\omega(\lambda\xi) = \lambda^{1-2n}\omega(\xi)$ for any $\lambda \in \mathbb{G}_m$ and $\xi \in \wedge_A^2 B$.

- for $G = \mathrm{GL}_n$, and $a \in \mathcal{A}$ we have a spectral cover $p_a: Y_a \rightarrow X$ embedded in the total space $|L|$ of L . We then associate with a the Higgs bundle $E_a = p_{a*}\mathcal{O}_{Y_a}$ and the Higgs fields $\phi: E_a \rightarrow E_a \otimes L$ given by the structure of \mathcal{O}_{Y_a} as $\mathcal{O}_{|L|}$ -module.
- for $G = \mathrm{Sp}_{2n}$, and $a \in \mathcal{A}$, we have a spectral cover $p_a: Y_a \rightarrow X$ embedded in the total space $|L|$ of L . If $E_a = p_{a*}\mathcal{O}_{Y_a}$ then we have a canonical symplectic form $\wedge^2 E_a \rightarrow L^{\otimes(1-2n)}$. If L' is a square root of L then $E'_a = E_a \otimes L'^{\otimes 1-2n}$ will be equipped with a canonical symplectic form with value in \mathcal{O}_X and also equipped with a Higgs fields derived from the the structure of \mathcal{O}_{Y_a} as a $\mathcal{O}_{|L|}$ -module.
- for $G = \mathrm{SO}_{2n+1}$, and $a \in \mathcal{A}$, we have a spectral cover $p_a: Y_a \rightarrow X$ embedded in the total space $|L|$ of L . If $E_a = p_{a*}\mathcal{O}_{Y_a}$ then we have a canonical non-degenerate symmetric form $S^2 E_a \rightarrow L^{\otimes(-2n)}$ so that the vector bundle $E'_a = E_a \otimes L^{\otimes n}$ affords a canonical no-degenerate symmetric form with value in \mathcal{O}_X , and also equipped with a Higgs fields derived from the the structure of \mathcal{O}_{Y_a} as a $\mathcal{O}_{|L|}$ -module. It also affords a trivialization of the determinant depending on the choice of a square root of L .
- for $G = \mathrm{SO}_{2n}$, and $a \in \mathcal{A}$, we have a spectral cover $p_a: Y_a \rightarrow X$ embedded in the total space $|L|$ of L . Using the normalization of the universal spectral cover, we obtain a partial normalization \tilde{Y}_a of Y_a . If $E_a = p_{a*}\mathcal{O}_{\tilde{Y}_a}$ then we have a canonical non-degenerate symmetric form $S^2 E_a \rightarrow L^{\otimes(2-2n)}$ so that the vector bundle $E'_a = E_a \otimes L^{\otimes 1-n}$ affords a canonical no-degenerate symmetric form with value in \mathcal{O}_X , and also equipped with a Higgs fields derived from the the structure of \mathcal{O}_{Y_a} as a $\mathcal{O}_{|L|}$ -module. It also affords a canonical trivialization of the determinant depending on the choice of a square root of L .
- for $G = G_2$, and $a \in \mathcal{A}$, we have a spectral cover $p_a: Y_a \rightarrow X$ embedded in the total space $|L|$ of L . If $E_a = p_{a*}\mathcal{O}_{\tilde{Y}_a}$ then we have a canonical non-degenerate 3-form $\wedge^3 E_a \rightarrow L^{-9}$ so that the vector bundle $E'_a = E_a \otimes L^{\otimes 3}$ affords a canonical non-degenerate 3-form with value in \mathcal{O}_X , and also equipped with a Higgs fields derived from the the structure of \mathcal{O}_{Y_a} as a $\mathcal{O}_{|L|}$ -module. It also affords a canonical trivialization of the determinant depending on the choice of a square root of L .

In [8], it is shown that over a large open subset of \mathcal{A} , there is a close connection between Hitchin fibers and affine Springer fibers given by the Product Formula. More precisely, let $\mathfrak{D} = \bigcup_{\alpha} \mathfrak{t}^{\alpha} // W$ be the divisor consisting of the union of the image of each root hyperplane in \mathfrak{t} ; in particular, the complement of \mathfrak{D} in \mathfrak{c} is the regular, semisimple locus \mathfrak{c}^{rs} . Fix $a \in \mathcal{A}$ such that $a(X) \not\subset \mathfrak{D}$, and let $U \subset X$ be the preimage of \mathfrak{c}^{rs} in X . Given trivialization of the line bundle D on some neighborhood of each point $v \in X \setminus U$, we have a map

$$\prod_{v \in X \setminus U} \mathcal{M}_{x,a} \rightarrow \mathcal{M}_a.$$

from the product of affine Springer fibers at the points $x \in X \setminus U$ to the Hitchin fiber, which consists

of gluing with the companion section restricted to U . It induces a universal homeomorphism

$$\prod_{\gamma \in X \setminus U} \mathcal{M}_{\gamma, a} \wedge \Pi_{\gamma} \mathcal{P}_{\gamma}(J_a) \mathcal{P}_a \rightarrow \mathcal{M}_a.$$

The groups $\mathcal{P}_{\gamma}(J_a)$ and \mathcal{P}_a are discussed in detail in [8]; we will not describe them here. This is proved in [8] under the assumption that $\pi_0(\mathcal{P}_a)$ is finite, and by Bouthier and Cesnavicius in [1] under the only assumption that $a(X) \notin \mathfrak{D}$.

A Computer algebra code and G_2 computations

In this appendix, we give the computer code used to compute the 3-form ρ in Section 3.5.

A.1 Construction of ρ

To construct ρ , we will use the connection between nondegenerate alternating 3-forms and cross products. Let V be a vector space with a nondegenerate, symmetric bilinear form ν .

Definition A.1. A cross product on (V, ν) is a bilinear map

$$c: V \otimes V \rightarrow V$$

satisfying the following three properties for all $v_1, v_2 \in V$:

1. (Skew symmetry) $c(v_1, v_2) = -c(v_2, v_1)$;
2. (Orthogonality) $\nu(c(v_1, v_2), v_1) = 0$;
3. (Normalization) $\nu(c(v_1, v_2), c(v_1, v_2)) = \det \begin{pmatrix} \nu(v_1, v_1) & \nu(v_1, v_2) \\ \nu(v_1, v_2) & \nu(v_2, v_2) \end{pmatrix}$

The data of a cross product on (V, ν) is equivalent to the data of a nondegenerate 3-form on V whose associated symmetric bilinear form (see equation (1)) is a scalar multiple of ν . Indeed, to a cross product c , one associates the 3-form

$$\rho(v_1, v_2, v_3) = \nu(c(v_1, v_2), v_3) \tag{8}$$

while for any non-degenerate 3-form ρ , there is a unique cross product c satisfying equation (8).

Now, consider the free, rank 7 A -module B as in Section 3.5 equipped with the symmetric, nondegenerate form ω defined by the formula

$$\omega(g_1, g_2) = \text{tr}_{B/A} \left(\frac{g_1 \tau(g_2)}{f'} \right)$$

as in the SO_7 case. Here, $\tau(x) = -x$ is the natural involution on B , and the trace is taken after inverting f' in A . To construct a 3-form on B which is nondegenerate over every k point of A , it suffices to construct a cross product

$$c: B \otimes_A B \rightarrow B$$

for (B, ω) . Moreover, the equation

$$\rho(xg_1, g_2, g_3) + \rho(g_1, xg_2, g_3) + \rho(g_1, g_2, xg_3) = 0$$

is equivalent to the condition

$$c(xg_1, g_2) + c(g_1, xg_2) = xc(g_1, g_2). \quad (9)$$

To simplify computations further, we note that any form $c: B \otimes_A B \rightarrow B$ satisfying the conditions of Definition A.1 and equation (9) can be recovered from its trace:

$$tc: B \otimes_A B \rightarrow A, \quad (g_1, g_2) \mapsto \text{tr}_{B/A}(c(g_1, g_2))$$

Indeed, if we express

$$c(x^i, x^j) = \sum_{l=0}^6 c_{i,j}^{(l)} x^l$$

then $c_{i,j}^{(6)} = tc(x^i, x^j)$ and

$$\text{tr}_{B/A}(x^l c(x^i, x^j)) = \sum_{r=0}^l \binom{l}{r} tc(x^{i+r}, x^{j+l-r})$$

can be expressed in terms of $c_{i,j}^{(m)}$ for $6-l \leq m \leq 6$. This allows us to recover the coefficients $c_{i,j}^{(l)}$ by downward induction on l .

This idea is implemented in the following Macaulay2 code. There is a one-dimensional solution space, which is specialized at a particular point to give the form stated in equation (6). Note that it is immediate from the computer calculation that the form ρ is valued in B and satisfies the conclusion of Proposition 3.5.

```
S=QQ[e,q];
F=frac(S);
R=F[p_(0,0) .. p_(6,6)]; -- ring with p_(i,j)=tc(x^i,x^j), 0\leq i,j\leq 6

-- The following three commands define tc(x^i,x^j) for i or j between 7 and 12 using
the relation x^7-e*x^5+e^4/4*x^3+q*x=0.
for l from 0 to 5 do [for k from 0 to 6 do p_(k,7+l)=e*p_(k,5+l)-(1/4)*e^2*p_(k,3+l)
-q*p_(k,1+l)];
for l from 0 to 5 do [for k from 0 to 6 do p_(7+l,k)=e*p_(5+l,k)-(1/4)*e^2*p_(3+l,k)
-q*p_(1+l,k)];
for l from 0 to 5 do [for k from 7 to 12 do p_(k,7+l)=e*p_(k,5+l)-(1/4)*e^2*p_(k,3+l)
-q*p_(k,1+l)];

-- I encodes orthogonality:
I = ideal(flatten for a from 0 to 6 list for k from 0 to 6 list sum(0..k,j->
binomial(k,j)*p_(k+j,a+k-j)));

-- J encodes skew symmetry:
J = ideal( flatten for a from 0 to 6 list for b from 0 to 6 list p_(a,b)+p_(b,a) );

-- The following encodes the normalization condition:
B=R[x]/(x^7-e*x^5+(1/4)*e^2*x^3+q*x);
-- determinant of norms of x^i,x^j:
f = (i,j) -> coefficient(x^6,(-1)^i*x^(2*i))*coefficient(x^6,(-1)^j*x^(2*j))
```



```

- coefficient(x^6,(-1)^j*x^(i+j))*coefficient(x^6,(-1)^j*x^(i+j));
-- norm of c(x^i,x^j):
g = (i,j) -> coefficient(x^6, (p_(i,j)*(x^6-e*x^4+(1/4)*e^2*x^2+q)+sum(0..1,l->
binomial(1,l)*p_(i+1,j+1-l))*(x^5-e*x^3+(1/4)*e^2*x)+sum(0..2,l->binomial(2,l)
*p_(i+1,j+2-l))*(x^4-e*x^2+(1/4)*e^2)+sum(0..3,l->binomial(3,l)*p_(i+1,j+3-l))
*(x^3-e*x)+sum(0..4,l->binomial(4,l)*p_(i+1,j+4-l))*(x^2-e)+sum(0..5,l->binomial(5,l)
*p_(i+1,j+5-l))*(x)+sum(0..6,l->binomial(6,l)*p_(i+1,j+6-l))) * (p_(i,j)*
((-x)^6-e*(-x)^4+(1/4)*e^2*(-x)^2+q)+sum(0..1,l->binomial(1,l)*p_(i+1,j+1-l))*
((-x)^5-e*(-x)^3+(1/4)*e^2*(-x))+sum(0..2,l->binomial(2,l)*p_(i+1,j+2-l))*
((-x)^4-e*(-x)^2+(1/4)*e^2)+sum(0..3,l->binomial(3,l)*p_(i+1,j+3-l))*((-x)^3-e*(-x))
+sum(0..4,l->binomial(4,l)*p_(i+1,j+4-l))*((-x)^2-e)+sum(0..5,l->binomial(5,l)
*p_(i+1,j+5-l))*(-x)+sum(0..6,l->binomial(6,l)*p_(i+1,j+6-l))) );
-- K encodes the normalization condition:
K = ideal(flatten for i from 0 to 6 list for j from 0 to 6 list f(i,j)-g(i,j));

```

```

Q=R/(I+J+K); -- imposing the relations on our ring of variables
Q2=Q/ideal(p_(6,3)-1,p_(6,4),p_(6,5)-5*e/2); -- specializes to our particular form rho

```

```

-- Computation of c from tc:
P=Q2[x]/(x^7-e*x^5+e^2/4*x^3+q);
C=table(for k from 0 to 6 list k, for k from 0 to 6 list k, (i,j) -> (p_(i,j)*(x^6
-e*x^4+(1/4)*e^2*x^2+q)+sum(0..1,l->binomial(1,l)*p_(i+1,j+1-l))*(x^5-e*x^3+(1/4)
*e^2*x)+sum(0..2,l->binomial(2,l)*p_(i+1,j+2-l))*(x^4-e*x^2+(1/4)*e^2)+sum(0..3,
l->binomial(3,l)*p_(i+1,j+3-l))*(x^3-e*x)+sum(0..4,l->binomial(4,l)*p_(i+1,j+4-l))
*(x^2-e)+sum(0..5,l->binomial(5,l)*p_(i+1,j+5-l))*(x)+sum(0..6,l->binomial(6,l)*
p_(i+1,j+6-l)))); -- This is the matrix for c with respect to the basis x^i, i=0,...,6
netList C -- displays C

```

A.2 Nondegeneracy of ρ

Let ρ be the form computed in the previous section, stated explicitly in equation (6). Note that since we specialized to a particular form in the previous section, it is not yet clear that this form is nondegenerate. For this, we produce the following code in Macaulay2 to explicitly compute the associated bilinear form as in Proposition 3.4. The following uses some basic operations on permutations from the package `SpechtModule` authored by Jonathan Niño in Macaulay2.

```

T=permutations {0,1,2,3,4,5,6};
n = (v,w) -> sum(0..7!-1, k-> permutationSign(T_k)*coefficient(x^6,v*(C_((T_k)_0))_((T_k)_1))
*coefficient(x^6,w*(C_((T_k)_2))_((T_k)_3))*coefficient(x^6,(-x)^((T_k)_4)
*(C_((T_k)_5))_((T_k)_6)) );
S=table(for k from 0 to 6 list k, for k from 0 to 6 list k, (i,j) -> n((-x)^i,(-x)^j);
netList S

```

References

- [1] Alexis Bouthier and Kestutis Cesnavicius. “Torsors on loop groups and the Hitchin fibration”. In: *arXiv preprint arXiv:1908.07480* (2019).

- [2] Friedrich Engel. *Ein neues, dem linearen Komplexe analoges Gebilde*. 1900.
- [3] Cristina Draper Fontanals. “Notes on G2: The Lie algebra and the Lie group”. In: *Differential Geometry and its Applications* 57 (2018), pp. 23–74.
- [4] Nigel Hitchin. “Stable bundles and integrable systems”. In: *Duke mathematical journal* 54.1 (1987), pp. 91–114.
- [5] Nigel Hitchin. “The geometry of three-forms in six dimensions”. In: *Journal of Differential Geometry* 55.3 (2000), pp. 547–576.
- [6] David Kazhdan and George Lusztig. “Fixed point varieties on affine flag manifolds”. In: *Israel Journal of Mathematics* 62.2 (1988), pp. 129–168.
- [7] Bertram Kostant. “Lie group representations on polynomial rings”. In: *American Journal of Mathematics* 85.3 (1963), pp. 327–404.
- [8] B. C. Ngo. “Le lemme fondamental pour les algèbres de Lie”. In: *Publications mathématiques de l’IHÉS* 111.1 (2010), pp. 1–169.
- [9] Jean-Pierre Serre. *Local fields*. Vol. 67. Springer Science & Business Media, 2013.